

Introductory Algebra

Andrew Gloag
Anne Gloag

adapted by
James Sousa

Say Thanks to the Authors

Click <http://www.ck12.org/saythanks>

(No sign in required)

To access a customizable version of this book, as well as other interactive content, visit www.ck12.org

CK-12 Foundation is a non-profit organization with a mission to reduce the cost of textbook materials for the K-12 market both in the U.S. and worldwide. Using an open-content, web-based collaborative model termed the **FlexBook®**, CK-12 intends to pioneer the generation and distribution of high-quality educational content that will serve both as core text as well as provide an adaptive environment for learning, powered through the **FlexBook Platform®**.

Copyright © 2012 CK-12 Foundation, www.ck12.org

The names “CK-12” and “CK12” and associated logos and the terms “**FlexBook®**” and “**FlexBook Platform®**” (collectively “CK-12 Marks”) are trademarks and service marks of CK-12 Foundation and are protected by federal, state, and international laws.

Any form of reproduction of this book in any format or medium, in whole or in sections must include the referral attribution link <http://www.ck12.org/saythanks> (placed in a visible location) in addition to the following terms.

Except as otherwise noted, all CK-12 Content (including CK-12 Curriculum Material) is made available to Users in accordance with the Creative Commons Attribution/Non-Commercial/Share Alike 3.0 Unported (CC BY-NC-SA) License (<http://creativecommons.org/licenses/by-nc-sa/3.0/>), as amended and updated by Creative Commons from time to time (the “CC License”), which is incorporated herein by this reference.

Complete terms can be found at <http://www.ck12.org/terms>.

Printed: August 22, 2012

flexbook
next generation textbooks



AUTHORS

Andrew Gloag

Anne Gloag

Contents

1	Real Numbers Operations	1
1.1	Integers and Rational Numbers	2
1.2	Addition of Rational Numbers Learning Objectives	10
1.3	Subtraction of Rational Numbers	17
1.4	Multiplication of Rational Numbers	24
1.5	The Distributive Property	33
1.6	Division of Rational Numbers	40
1.7	Square Roots and Real Numbers	46
2	Expressions and Equations	53
2.1	Variable Expressions	54
2.2	Order of Operations	61
2.3	One-Step Equations	70
2.4	Two-Step Equations	79
2.5	Multi-Step Equations	88
2.6	Equations with Variables on Both Sides	96
2.7	Ratios and Proportions	104
3	Graphing Linear Equations	113
3.1	The Coordinate Plane	114
3.2	Graphs of Linear Equations	119
3.3	Graphing Using Intercepts	126
3.4	Slope and Rate of Change	136
3.5	Graphs Using Slope-Intercept Form	145
3.6	Direct Variation Models	154
3.7	Problem-Solving Strategies - Graphs	162
4	Determining Linear Equations	169
4.1	Linear Equations in Slope-Intercept Form	170
4.2	Linear Equations in Point-Slope Form	179
4.3	Linear Equations in Standard Form	186
4.4	Equations of Parallel and Perpendicular Lines	193
5	Introduction to Functions and Modeling	201
5.1	Functions as Rules and Tables	202
5.2	Functions as Graphs	210
5.3	Linear Function Graphs	224
5.4	Fitting a Line to Data	231
5.5	Predicting with Linear Models	240
5.6	Problem Solving Strategies: Use a Linear Model	248

6	Linear Inequalities	256
6.1	Solving One-Step Inequalities	257
6.2	Multi-Step Inequalities	269
6.3	Compound Inequalities	278
6.4	Absolute Value Equations	290
6.5	Absolute Value Inequalities	300
6.6	Linear Inequalities in Two Variables	305
7	Systems of Equations and Inequalities	318
7.1	Linear Systems by Graphing	319
7.2	Solving Linear Systems by Substitution	330
7.3	Solving Linear Systems by Elimination through Addition or Subtraction	339
7.4	Solving Systems of Equations by Multiplication	345
7.5	Special Types of Linear Systems	354
7.6	Systems of Linear Inequalities	364
8	Exponents and Polynomials	379
8.1	Exponent Properties Involving Products	380
8.2	Exponent Properties Involving Quotients	388
8.3	Zero, Negative, and Fractional Exponents	395
8.4	Scientific Notation	402
8.5	Addition and Subtraction of Polynomials	412
8.6	Multiplication of Polynomials	422
8.7	Special Products of Polynomials	433
8.8	Division of Polynomials	439
9	Introduction to Factoring	445
9.1	Polynomial Equations in Factored Form	446
9.2	Factoring Quadratic Expressions	456
9.3	Factoring Special Products	464
9.4	Factoring Polynomials Completely	474

CHAPTER

1**Real Numbers Operations****Chapter Outline**

- 1.1 INTEGERS AND RATIONAL NUMBERS**
 - 1.2 ADDITION OF RATIONAL NUMBERS LEARNING OBJECTIVES**
 - 1.3 SUBTRACTION OF RATIONAL NUMBERS**
 - 1.4 MULTIPLICATION OF RATIONAL NUMBERS**
 - 1.5 THE DISTRIBUTIVE PROPERTY**
 - 1.6 DIVISION OF RATIONAL NUMBERS**
 - 1.7 SQUARE ROOTS AND REAL NUMBERS**
-

1.1 Integers and Rational Numbers

Learning Objectives

- Graph and compare integers.
- Classify and order rational numbers.
- Find opposites of numbers.
- Find absolute values.
- Compare fractions to determine which is bigger.


Graph and Compare Integers

Integers are the counting numbers ($1, 2, 3, \dots$), the negative counting numbers ($-1, -2, -3, \dots$) and zero. There are an infinite number of integers. Examples of integers are $0, 3, 76, -2, -11, 995, \dots$ and you may know them by the name **whole numbers**. When we represent integers on the number line they fall exactly on the whole numbers.

Example 1

Compare the numbers 2 and -5

First, we will plot the two numbers on a number line.



We can compare integers by noting which is the **greatest** and which is the **least**. The **greatest** number is farthest to the right, and the **least** is farthest to the left.

In the diagram above, we can see that 2 is farther to the right on the number line than -5 , so we say that 2 is greater than -5 . We use the symbol “ $>$ ” to mean “greater than”.

Solution


$$2 > -5$$

Example 2

A frog is sitting perfectly on top of number 7 on a number line. The frog jumps randomly to the left or right, but always jumps a distance of exactly 2. Describe the set of numbers that the frog may land on, and list all the possibilities for the frog’s position after exactly 5 jumps.

Solution

We will graph the frog’s position, and also indicate what a jump of 2 looks like. We see that one possibility is that the frog lands on 5. Another possibility is that it lands on 9. It is clear that the frog will always land on an **odd number**.



After one jump the frog could be on either the 9 or the 5 (but not on the 7). After two jumps the frog could be on 11, 7 or 3. By counting the number of times the frog jumps to the right or left, we may determine where the frog lands.

After five jumps, there are many possible locations for the frog. There is a systematic way to determine the possible locations by how many times the frog jumped right, and by how many times the frog jumped left.

RRRRR = 5 jumps right	location = $7 + (5 \cdot 2) = 17$
RRRRL = 4 jumps right, 1 jump left	location = $7 + (3 \cdot 2) = 13$
RRLLL = 3 jumps right, 2 jumps left	location = $7 + (1 \cdot 2) = 9$
RLLLL = 2 jumps right, 3 jumps left	location = $7 - (1 \cdot 2) = 5$
RLLLL = 1 jump right, 4 jumps left	location = $7 - (3 \cdot 2) = 1$
LLLLL = 5 jumps left	location = $7 - (5 \cdot 2) = -3$

These are the possible locations of the frog after exactly five jumps. Notice that the order does not matter: three jumps right, one left and one right is the same as four jumps to the right and one to the left.

Classifying Rational Numbers

When we divide an integer by another integer (*not zero*) we get what we call a **rational number**. It is called this because it is the **ratio** of one number to another. For example, if we divide one integer a by a second integer b the rational number we get is $\frac{a}{b}$, provided that b is not zero. When we write a rational number like this, the top number is called the **numerator**. The bottom number is called the **denominator**. You can think of the rational number as a fraction of a cake. If you cut the cake into b slices, your share is a of those slices.

For example, when we see the rational number $\frac{1}{2}$, we imagine cutting the cake into two parts. Our share is one of those parts. Visually, the rational number $\frac{1}{2}$ looks like this.



With the rational number $\frac{3}{4}$, we cut the cake into four parts and our share is three of those parts. Visually, the rational number $\frac{3}{4}$ looks like this.



The rational number $\frac{9}{10}$ represents nine slices of a cake that has been cut into ten pieces. Visually, the rational number $\frac{9}{10}$ looks like this.



Proper fractions are rational numbers where the numerator (the number on the top) is less than the denominator (the number on the bottom). A proper fraction represents a number less than one. With a proper fraction you always end up with less than a whole cake!

Improper fractions are rational numbers where the numerator is greater than the denominator. Improper fractions can be rewritten as a mixed number – an integer plus a proper fraction. An improper fraction represents a number greater than one.

Equivalent fractions are two fractions that give the same numerical value when evaluated. For example, look at a visual representation of the rational number $\frac{2}{4}$.



You can see that the shaded region is identical in size to that of the rational number one-half $\frac{1}{2}$. We can write out the prime factors of both the numerator and the denominator and cancel matching factors that appear in both the numerator **and** denominator.

$$\left(\frac{2}{4}\right) = \left(\frac{\cancel{2} \cdot 1}{\cancel{2} \cdot 2 \cdot 1}\right) \quad \text{We then re-multiply the remaining factors.} \quad \left(\frac{2}{4}\right) = \left(\frac{1}{2}\right)$$

This process is called **reducing** the fraction, or writing the fraction in lowest terms. Reducing a fraction does not change the value of the fraction. It just simplifies the way we write it. When we have canceled all common factors, we have a fraction in its **simplest form**.

Example 3

Classify and simplify the following rational numbers

a) $\left(\frac{3}{7}\right)$

b) $\left(\frac{9}{3}\right)$

c) $\left(\frac{50}{60}\right)$

a) 3 and 7 are both prime – there is no simpler form for this rational number so...

Solution

$\frac{3}{7}$ is already in its simplest form.

b) $9 = 3 \cdot 3$ and 3 is prime. We rewrite the fraction as: $\left(\frac{9}{3}\right) = \left(\frac{\cancel{3} \cdot 3 \cdot 1}{\cancel{3} \cdot 1}\right)$. $9 > 3$ so...

Solution

$\frac{9}{3}$ is an improper fraction and simplifies to $\frac{3}{1}$ or simply 3.

c) $50 = 5 \cdot 5 \cdot 2$ and $60 = 5 \cdot 3 \cdot 2 \cdot 2$. We rewrite the fraction thus: $\frac{50}{60} = \left(\frac{\cancel{5} \cdot 5 \cdot \cancel{2} \cdot 1}{\cancel{5} \cdot 5 \cdot \cancel{2} \cdot 2 \cdot 1}\right)$. $50 < 60$ so...

Solution

$\frac{50}{60}$ is a proper fraction and simplifies to $\frac{5}{6}$.

Order Rational Numbers

Ordering rational numbers is simply a case of arranging numbers in order of increasing value. We write the numbers with the least (most negative) first and the greatest (most positive) last.

Example 4

Put the following fractions in order from least to greatest: $\frac{1}{2}, \frac{3}{4}, \frac{2}{3}$

Solution

$$\frac{1}{2} < \frac{2}{3} < \frac{3}{4}$$

Simple fractions are easy to order—we just know, for example, that one-half is greater than one quarter, and that two thirds is bigger than one-half. But how do we compare more complex fractions?

With simple fractions, it is easy to order them. Think of the example above. We know that one-half is greater than one quarter, and we know that two thirds is bigger than one-half. With more complex fractions, however we need to find a better way to compare.

Example 5

Which is greater, $\frac{3}{7}$ or $\frac{4}{9}$?


In order to determine this we need to find a way to rewrite the fractions so that we can better compare them. We know that we can write equivalent fractions for both of these. If we make the denominators in our equivalent fractions the same, then we can compare them directly. We are looking for the lowest common multiple of each of the denominators. This is called finding the **lowest common denominator** (LCD).

The lowest common multiple of 7 and 9 is 63. Our fraction will be represented by a shape divided into 63 sections. This time we will use a rectangle cut into 9 by 7 = 63 pieces:

7 divides into 63 nine times so:

$$\left(\frac{3}{7}\right) = \frac{9}{9} \left(\frac{3}{7}\right) = \left(\frac{27}{63}\right)$$


Note that multiplying by $\frac{9}{9}$ is the same as multiplying by 1. Therefore, $\frac{27}{63}$ is an equivalent fraction to $\frac{3}{7}$. Here it is shown visually.




9 divides into 63 seven times so:

$$\left(\frac{4}{9}\right) = \frac{7}{7} \left(\frac{4}{9}\right) = \left(\frac{28}{63}\right)$$

$\frac{28}{63}$ is an equivalent fraction to $\frac{4}{9}$. Here it is shown visually.



By writing the fractions over a **common denominator** of 63, you can easily compare them. Here we take the 28 shaded boxes out of 63 (from our image of $\frac{4}{9}$ above) and arrange them in a way that makes it easy to compare with our representation of $\frac{3}{7}$. Notice there is one little square “left over”.



Solution


Since $\frac{28}{63}$ is greater than $\frac{27}{63}$, $\frac{4}{9}$ is greater than $\frac{3}{7}$.

Remember

To compare rational numbers re-write them with a **common denominator**.

Find the Opposites of Numbers

Every number has an opposite. On the number line, a number and its opposite are *opposite* each other. In other words, they are the same distance from zero, but they are on opposite sides of the number line.



By definition, the opposite of zero is zero.

Example 6

Find the value of each of the following.

- a) $3 + (-3)$
- b) $5 + (-5)$
- c) $(-11.5) + (11.5)$
- d) $\frac{3}{7} + \frac{-3}{7}$

Each of the pairs of numbers in the above example are **opposites**. The opposite of 3 is (-3) , the opposite of 5 is (-5) , the opposite of (-11.5) is 11.5 and the opposite of $\frac{3}{7}$ is $\frac{3}{7}$.

Solution

The value of each and every sum in this problem is 0.

Example 7

Find the opposite of each of the following:

- a) 19.6
- b) $-\frac{4}{9}$
- c) x
- d) xy^2
- e) $(x - 3)$

Since we know that opposite numbers are on opposite sides of zero, we can simply multiply each expression by -1 . This changes the sign of the number to its opposite.

a) Solution

The opposite of 19.6 is -19.6 .

b) Solution

The opposite of $-\frac{4}{9}$ is $\frac{4}{9}$.

c) Solution

The opposite of x is $-x$.

d) Solution

The opposite of xy^2 is $-xy^2$.

e) Solution

The opposite of $(x - 3)$ is $-(x - 3) = 3 - x$.

Note: With the last example you must multiply the **entire expression** by -1 . A **common mistake** in this example is to assume that the opposite of $(x - 3)$ is $(x + 3)$. **DO NOT MAKE THIS MISTAKE!**

Find absolute values

When we talk about absolute value, we are talking about distances on the number line. For example, the number 7 is 7 units away from zero. The number -7 is also 7 units away from zero. The absolute value of a number is the

distance it is from zero, so the absolute value of 7 and the absolute value of -7 are both 7.

We **write** the absolute value of -7 like this $|-7|$. We **read** the expression $|x|$ like this “the absolute value of x .”

- Treat absolute value expressions like parentheses. If there is an operation inside the absolute value symbols evaluate that operation first.
- The absolute value of a number or an expression is **always** positive or zero. It cannot be negative. With absolute value, we are only interested in how far a number is from zero, and not the direction.

Example 8

Evaluate the following absolute value expressions.

a) $|5 + 4|$

b) $3 - |4 - 9|$

c) $|-5 - 11|$

d) $-|7 - 22|$

Remember to treat any expressions inside the absolute value sign as if they were inside parentheses, and evaluate them first.

Solution

a)

$$\begin{aligned} |5 + 4| &= |9| \\ &= 9 \end{aligned}$$

b)

$$\begin{aligned} 3 - |4 - 9| &= 3 - |-5| \\ &= 3 - 5 \\ &= -2 \end{aligned}$$

c)

$$\begin{aligned} |-5 - 11| &= |-16| \\ &= 16 \end{aligned}$$

d)

$$\begin{aligned} -|7 - 22| &= -|-15| \\ &= -(15) \\ &= -15 \end{aligned}$$


- **Integers** (or **whole numbers**) are the counting numbers (1, 2, 3 ...), the negative counting numbers (−1, −2, −3 ...), and zero.
- A **rational number** is the **ratio** of one integer to another, like $\frac{a}{b}$ or $\frac{3}{5}$. The top number is called the **numerator** and the bottom number (which can not be zero) is called the **denominator**.
- **Proper fractions** are rational numbers where the numerator is less than the denominator.
- **Improper fractions** are rational numbers where the numerator is greater than the denominator.
- Equivalent fractions are two fractions that give the same numerical value when evaluated.
- To **reduce** a fraction (write it in **simplest form**) write out all prime factors of the numerator and denominator, cancel common factors, then recombine.
- To compare two fractions it helps to write them with a **common denominator**: the same integer on the bottom of each fraction.
- The **absolute value** of a number is the distance it is from zero on the number line. The absolute value of a number or expression will always be positive or zero.
- Two numbers are **opposites** if they are the same distance from zero on the number line and on opposite sides of zero. The opposite of an expression can be found by multiplying **the entire expression** by -1 .

Review Questions

1. The tick-marks on the number line represent evenly spaced integers. Find the values of a, b, c, d and e .



2. Determine what fraction of the whole each shaded region represents.



3. Place the following sets of rational numbers in order, from least to greatest.

- (a) $\frac{1}{2}, \frac{1}{3}, \frac{1}{4}$
 (b) $\frac{11}{12}, \frac{12}{11}, \frac{13}{10}$
 (c) $\frac{39}{60}, \frac{49}{80}, \frac{59}{100}$
 (d) $\frac{7}{11}, \frac{8}{13}, \frac{12}{19}$

4. Find the simplest form of the following rational numbers.

- (a) $\frac{22}{44}$
 (b) $\frac{4}{9}$
 (c) $\frac{12}{27}$
 (d) $\frac{315}{420}$

5. Find the opposite of each of the following.

- (a) 1.001
 (b) $(5 - 11)$
 (c) $(x + y)$
 (d) $(x - y)$

6. Simplify the following absolute value expressions.

- (a) $11 - |-4|$
 (b) $|4 - 9| - |-5|$

- (c) $|-5 - 11|$
- (d) $7 - |22 - 15 - 19|$
- (e) $-|-7|$
- (f) $|-2 - 88| - |88 + 2|$

Review Answers

- 1. $a = -3; b = 3; c = 9; d = 12; e = 15$
- 2. $a = \frac{1}{3}; b = \frac{7}{12}; c = \frac{22}{35}$
- 3. (a) $\frac{1}{4} < \frac{1}{3} < \frac{1}{2}$
(b) $\frac{11}{12} < \frac{12}{11} < \frac{13}{10}$
(c) $\frac{59}{100} < \frac{49}{80} < \frac{39}{60}$
(d) $\frac{8}{13} < \frac{12}{19} < \frac{7}{11}$
- 4. (a) $\frac{1}{2}$
(b) $\frac{1}{3}$
(c) $\frac{3}{3}$
(d) $\frac{3}{4}$
- 5. (a) -1.001
(b) $6 - (x + y)$
(c) $(y - x)$
- 6. (a) 7
(b) 0
(c) 16
(d) -5
(e) -7
(f) 0

1.2 Addition of Rational Numbers Learning Objectives

Learning Objectives

- Add using a number line.
- Add rational numbers.
- Identify and apply properties of addition.
- Solve real-world problems using addition of fractions.

Add Using a Number Line

In Lesson one, we learned how to represent numbers on a number line. When we perform addition on a number line, we start at the position of the first number, and then move to the right by the number of units shown in the sum.

Example 1

Represent the sum $2 + 3$ on a number line.

We start at the number 2, and then move 3 to the right. We end at the number 5.

Solution

$$2 + 3 = 5$$




Example 2

Represent the sum $-2 + 3$ on a number line.

We start at the number -2 , and then move 3 to the right. We thus end at $+1$.

Solution

$$-2 + 3 = 1$$




Example 3

Represent the sum $2 - 3$ on a number line.

We are now faced with a subtraction. When subtracting a number, an equivalent action is **adding a negative number**. Either way, we think of it, we are moving to the left. We start at the number 2, and then move 3 to the left. We end at -1 .

Solution

$$2 - 3 = -1$$



If they have the same sign : Add the absolute value and use the common sign. This means a **positive + positive = positive** and a **negative + negative = negative**


$$3 + 5 = 8 \quad -3 + (-5) = -8$$

If they have different signs : Subtract the smaller absolute value from the larger absolute value. The answer will have the sign of the number with the larger absolute value.

$$-3 + 5 = 2 \quad 3 + (-5) = -2$$

The money analogy Using a money analogy might be helpful when adding signed numbers. When you add signed numbers you can think of a positive number as gaining money and a negative number as a loss of money. For example, $-3 + 5$ means you lose 3 dollars and then gain 5 dollars. This leaves you with 2 dollars so the answer is positive 2. $3 + (-5)$ means you gain 3 dollars and then lose 5 dollars. As a result, you owe 2 dollars so the result is -2.

We can use the number line as a rudimentary way of adding fractions. The enlarged number line below has a number of common fractions marked. The markings on a ruler or a tape measure follow the same pattern. The two shaded bars represent the lengths $\frac{1}{3}$ and $\frac{1}{4}$.



To find the difference between the two fractions look at the difference between the two lengths. You can see the red is $\frac{1}{12}$ longer than the yellow. You could use this as an estimate of the difference.

$$\text{equation} = \frac{1}{3} - \frac{1}{4} = \frac{1}{12}$$

To find the sum of the two fractions, we can lay them end to end. You can see that the sum $\frac{1}{3} + \frac{1}{4}$ is a little over one half.



Adding Rational Numbers

We have already seen the method for writing rational numbers over a common denominator. When we add two fractions we need to ensure that the denominators match before we can determine the sum.

Rules

1. Add or subtract the numerators. 2. Keep the denominators the same. 3. Simplify if possible

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

If the

1. Factor each denominator to build the LCD. The LCD is the least common multiple of the denominators. 2. Rewrite each fraction as an equivalent fraction that has the LCD for its denominator. 3. Add or subtract the numerators. The denominators stay the same. 4. Simplify if possible.

While our goal to add fractions is to determine the **least common denominator**, the following procedure will work. However, it will more than likely require additional simplifying.

$$\frac{a}{c} + \frac{b}{d} = \frac{a \cdot d}{c \cdot d} + \frac{b \cdot c}{d \cdot c} = \frac{a \cdot d + b \cdot c}{c \cdot d}$$

Example 4

Simplify $\frac{3}{5} + \frac{1}{6}$

To combine these fractions, we need to rewrite them over a common denominator. We are looking for the **lowest common denominator** (LCD). We need to identify the **lowest common multiple** or **least common multiple** (LCM) of 5 and 6. That is the smallest number that both 5 and 6 divide into without remainder.

- The lowest number that 5 and 6 both divide into without remainder is 30. The LCM of 5 and 6 is 30, so the lowest common denominator for our fractions is also 30.

We need to rewrite our fractions as new **equivalent fractions** so that the denominator in each case is 30.

If you think back to our idea of a cake cut into a number of slices, $\frac{3}{5}$ means 3 slices of a cake that has been cut into 5 pieces. You can see that if we cut the same cake into 30 pieces (6 times as many) we would need 18 slices to have an equivalent share, since $18 = 3 \times 6$.

$\frac{3}{5}$ is equivalent to $\frac{18}{30}$



By a similar argument, we can rewrite the fraction $\frac{1}{6}$ as a share of a cake that has been cut into 30 pieces. If we cut it into 5 times as many pieces we require 5 times as many slices.

$\frac{1}{6}$ is equivalent to $\frac{5}{30}$



Now that both fractions have the same common denominator, we can add the fractions. If we add our 18 smaller pieces of cake to the additional 5 pieces you see that we get a total of 23 pieces. 23 pieces of a cake that has been cut into 30 pieces means that our answer is.

Solution



$$\frac{3}{5} + \frac{1}{6} = \frac{18}{30} + \frac{5}{30} = \frac{23}{30}$$

You should see that when we have fractions with a common denominator, we **add the numerators** but we **leave the denominators alone**. Here is this information in algebraic terms.

When adding fractions: $\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$

Example 5

Simplify $\frac{14}{11} + \frac{1}{9}$

The lowest common denominator in this case is 99. This is because the lowest common multiple of 9 and 11 is 99. So we write equivalent fractions for both $\frac{14}{11}$ and $\frac{1}{9}$ with denominators of 99.

11 divides into 99 nine times so $\frac{14}{11}$ is equivalent to $\frac{14 \cdot 9}{11 \cdot 9} = \frac{126}{99}$

We can multiply the numerator and denominator by 9 (or by any number) since $9/9 = 1$ and 1 is the multiplicative identity.

9 divides into 99 eleven times so $\frac{1}{9}$ is equivalent to $\frac{1 \cdot 11}{9 \cdot 11} = \frac{11}{99}$.

Now we simply add the numerators.

Solution

$$\frac{14}{11} + \frac{1}{9} = \frac{126}{99} + \frac{11}{99} = \frac{137}{99}$$

Example 6

Simplify

$$-\frac{1}{12} + \frac{2}{9}$$

The least common denominator in this case is 36. This is because the LCM of 12 and 9 is 36. We now proceed to write the equivalent fractions with denominators of 36.

12 divides into 36 three times so $-\frac{1}{12}$ is equivalent to $\frac{-1 \cdot 3}{12 \cdot 3} = \frac{-3}{36}$.

9 divides into 36 four times so $\frac{2}{9}$ is equivalent to $\frac{2 \cdot 4}{9 \cdot 4} = \frac{8}{36}$.

Solution

$$-\frac{1}{12} + \frac{2}{9} = \frac{-3}{36} + \frac{8}{36} = \frac{5}{36}$$

You can see that we quickly arrive at an equivalent fraction by multiplying the numerator and the denominator by the same non-zero number.

Example 7

Simplify $-\frac{2}{15} + \frac{2}{25}$

The least common denominator is 75. This is because the LCM of 15 and 25 is 75. We now proceed to write the equivalent fractions with the denominator of 75.

Solution


$$\frac{-2}{3 \cdot 5} + \frac{3}{5 \cdot 5} = \frac{-2 \cdot 5}{3 \cdot 5 \cdot 5} + \frac{3 \cdot 3}{5 \cdot 5 \cdot 3} = \frac{-10+6}{75} = -\frac{4}{75}$$

Identify and Apply Properties of Addition

The three mathematical properties which involve addition are the **commutative**, **associative**, and the **additive identity properties**.

- **Commutative property** When two numbers are added, the sum is the same even if the order of the items being added changes.

Example $3 + 2 = 2 + 3$



On a number line this means move 3 units to the right then 2 units to the right. The commutative property says this is equivalent of moving 2 units to the right then 3 units to the right. You can see that they are both the same, as they both end at 5.

- **Associative Property** When three or more numbers are added, the sum is the same regardless of how they are grouped.

Example $(2 + 3) + 4 = 2 + (3 + 4)$

- **Additive Identity Property** The sum of any number and zero is the original number.

Example $5 + 0 = 5$

Example 8

Nadia and Peter are building sand castles on the beach. Nadia built a castle two feet tall, stopped for ice-cream and then added one more foot to her castle. Peter built a castle one foot tall before stopping for a sandwich. After his sandwich, he built up his castle by two more feet. Whose castle is the taller?

Solution

Nadia's castle is $(2 + 1)$ feet tall. Peter's castle is $(1 + 2)$ feet tall. According to the **Commutative Property of Addition**, the two castles are the same height.

Example 9

Nadia and Peter each take candy from the candy jar. Peter reaches in first and grabs one handful. He gets seven pieces of candy. Nadia grabs with both hands and gets seven pieces in one hand and five in the other. The following day Peter gets to go first. He grabs with both hands and gets five pieces in one hand and six in the other. Nadia, grabs all the remaining candy, six pieces, in one hand. In total, who got the most candy?

Solution

On day one, Peter gets 7 candies, and on day two he gets $(5 + 6)$ pieces. His total is $7 + (5 + 6)$. On day one, Nadia gets $(7 + 5)$ pieces. On day two, she gets 6. Nadia's total is therefore $(7 + 5) + 6$. According to the **Associative Property of Addition** they both received exactly the same amount.

Solve Real-World Problems Using Addition

Example 10

Peter is hoping to travel on a school trip to Europe. The ticket costs \$2400. Peter has several relatives who have pledged to help him with the ticket cost. His parents have told him that they will cover half the cost. His grandma Zenoviea will pay one sixth, and his grandparents in Florida will send him one fourth of the cost. What fraction of the cost can Peter count on his relatives to provide?

The first thing we need to do is extract the relevant information. Here is what Peter can count on.

$\left(\frac{1}{2}\right)$	From parents
$\left(\frac{1}{6}\right)$	From grandma
$\left(\frac{1}{4}\right)$	From grandparents in Florida

Here is our problem. $\frac{1}{2} + \frac{1}{6} + \frac{1}{4}$

To determine the sum, we first need to find the LCD. The LCM of 2, 6 and 4 is 12. This is our LCD.

2 divides into 12 six times :	$\frac{1}{2} = \frac{6 \cdot 1}{6 \cdot 2} = \frac{6}{12}$
6 divides into 12 two times :	$\frac{1}{6} = \frac{2 \cdot 1}{2 \cdot 6} = \frac{2}{12}$
4 divides into 12 six times :	$\frac{1}{4} = \frac{3 \cdot 1}{3 \cdot 4} = \frac{3}{12}$
So an equivalent sum for our problem is	$\frac{6}{12} + \frac{2}{12} + \frac{3}{12} = \frac{(6+2+3)}{12} = \frac{11}{12}$

Solution

Peter can count on eleven-twelfths of the cost of the trip (\$2,200 out of \$2,400).

Lesson Summary

- To add fractions, rewrite them over the **lowest common denominator (LCD)**. The lowest common denominator is the **lowest (or least) common multiple (LCM)** of the two denominators.
- When **adding fractions**: $\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$
- The fractions $\frac{a}{b}$ and $\frac{a \cdot c}{b \cdot c}$ are **equivalent** when $c \neq 0$
- The **additive properties** are:
 - Commutative property** the sum of two numbers is the same even if the order of the items to be added changes.

Ex: $2 + 3 = 3 + 2$


- **Associative Property** When three or more numbers are added, the sum is the same regardless of how they are grouped.

Ex: $(2 + 3) + 4 = 2 + (3 + 4)$

- **Additive Identity Property** The sum of any number and zero is the original number.

Review Question

- Write the sum that the following moves on a number line represent.



- Add the following rational numbers and write the answer in its **simplest form**.

- $\frac{3}{7} + \frac{2}{7}$
- $\frac{3}{10} + \frac{1}{5}$
- $\frac{5}{16} + \frac{5}{12}$
- $-\frac{3}{8} + \frac{9}{16}$
- $\frac{8}{25} + \frac{7}{10}$
- $\frac{1}{6} + \left(-\frac{1}{4}\right)$
- $\frac{7}{15} + \left(-\frac{2}{9}\right)$
- $\frac{5}{19} + \frac{2}{27}$

- Which property of addition does each situation involve?

- Whichever order your groceries are scanned at the store, the total will be the same.
- However many shovel-loads it takes to move 1 ton of gravel the number of rocks moved is the same.

- Nadia, Peter and Ian are pooling their money to buy a gallon of ice cream. Nadia is the oldest and gets the greatest allowance. She contributes half of the cost. Ian is next oldest and contributes one third of the cost. Peter, the youngest, gets the smallest allowance and contributes one fourth of the cost. They figure that this will be enough money. When they get to the check-out, they realize that they forgot about sales tax and worry there will not be enough money. Amazingly, they have exactly the right amount of money. What fraction of the cost of the ice cream was added as tax?

Review Answers

- $3 + 3 + 1 - 8 = -1$
 - $1 - 2 + 7 + 1 = 7$
- $\frac{5}{7}$
 - $\frac{1}{2}$
 - $\frac{35}{48}$
 - $\frac{3}{16}$
 - $\frac{51}{50}$
 - $-\frac{1}{12}$
 - $\frac{11}{45}$
 - $\frac{173}{513}$
- Commutative and Associative
 - Associative
- $\frac{1}{12}$ is added as tax.


1.3 Subtraction of Rational Numbers

Learning objectives

- Find additive inverses.
- Subtract rational numbers.
- Evaluate change using a variable expression.
- Solve real world problems using fractions.

Find Additive Inverses

The **additive inverse** of a number is simply **opposite** of the number. (see section 2.1.4). Here are opposites on a number line.



When we think of additive inverses we are really talking about the **opposite process** (or **inverse process**) of **addition**. In other words, the process of **subtracting** a number is the same as adding the **additive inverse** of that number. When we add a number to its additive inverse, we get zero as an answer.

$$(6) + (-6) = 0$$

-6 is the additive inverse of 6 .

$$(279) + (-279) = 0$$

-279 is the additive inverse of 279 .

$$(x) + (-x) = 0$$

$-x$ is the additive inverse of x .

For any real numbers a and b ,

$$\mathbf{a - b = a + (-b) \text{ and } a - (-b) = a + b}$$

(To subtract, add the opposite, or additive inverse, of the number being subtracted.)

In general, this rule states we can convert every subtraction problem to an addition problem. We can then follow the rules for adding integers. Evaluate change using a variable expression.

Example 1

Rewrite each subtraction problem as an addition problem. Then determine the sum.

a. $5 - 3$ b. $-6 - 7$ c. $8 - (-2)$ d. $-9 - (-4)$

Solution

a. $5 + (-3) = 2$ b. $-6 + (-7) = -13$ c. $8 + 2 = 10$ d. $-9 + 4 = -5$

The method for subtracting fractions (as you should have assumed) is just the same as addition. We can use the idea of an additive inverse to relate the two processes. Just like in addition, we are going to need to write each of the rational numbers over a common denominator.

Example 2Simplify $\frac{1}{3} - \frac{1}{9}$

The lowest common multiple of 9 and 3 is 9. Our common denominator will be nine. We will not alter the second fraction because the denominator is already nine.

3 divides into 9 three times $\frac{1}{3} = \frac{1 \cdot 3}{3 \cdot 3} = \frac{3}{9}$. In other words $\frac{3}{9}$ is an **equivalent fraction** to $\frac{1}{3}$.

Our sum becomes $\frac{3}{9} - \frac{1}{9}$

Remember that when we add fractions with a common denominator, we **add** the **numerators** and the **denominator is unchanged**. A similar relationship holds for subtraction, only that we subtract the numerators.

When subtracting fractions $\frac{a}{c} - \frac{b}{c} = \frac{a-b}{c}$

Solution


$$\frac{1}{3} - \frac{1}{9} = \frac{2}{9}$$

Two-ninths is the simplest form for our answer. So far we have only dealt with examples where it is easy to find the least common multiple of the denominators. With larger numbers, it is not so easy to be certain that we have the **least common denominator** (LCD). We need a more systematic method. In the next example, we will use the method of **prime factors** to find the least common denominator.

Example 3Simplify $\frac{29}{90} - \frac{13}{126}$

This time we need to find the lowest common multiple (LCM) of 90 and 126. To find the LCM, we first find the prime factors of 90 and 126. We do this by continually dividing the number by factors until we cannot divide any further. You may have seen a factor tree before:

The factor tree for 90 looks like this:




$$90 = 9 \cdot 10$$

$$9 = 3 \cdot 3$$

$$10 = 5 \cdot 2$$

$$90 = 3 \cdot 3 \cdot 5 \cdot 2$$

The factor tree for 126 looks like this:



$$\begin{aligned}
 126 &= 9 \cdot 14 \\
 9 &= 3 \cdot 3 \\
 14 &= 7 \cdot 2 \\
 126 &= 3 \cdot 3 \cdot 7 \cdot 2
 \end{aligned}$$

The LCM for 90 and 126 is made from the **smallest possible collection of primes** that enables us to construct either of the two numbers. We take only enough of each prime to make the number with the highest number of factors of that prime in its factor tree.

TABLE 1.1:

Prime	Factors in 90	Factors in 126	We Take
2	1	1	1
3	2	2	2
5	1	0	1
7	0	1	1

So we need: one 2, two 3's, one 5 and one 7. In other words: $2 \cdot 3 \cdot 3 \cdot 5 \cdot 7 = 630$

- The lowest common multiple of 90 and 126 is 630. The LCD for our calculation is 630.

90 divides into 630 seven times (notice that 7 is the only factor in 630 that is missing from 90) $\frac{29}{90} = \frac{7 \cdot 29}{7 \cdot 90} = \frac{203}{630}$

126 divides into 630 five times (notice that 5 is the only factor in 630 that is missing from 126) $\frac{13}{126} = \frac{5 \cdot 13}{5 \cdot 126} = \frac{65}{630}$

Now we complete the problem.

$$\frac{29}{90} - \frac{13}{126} = \frac{203}{630} - \frac{65}{630} = \frac{(203 - 65)}{630} = \frac{138}{630} \left\{ \text{remember, } \frac{a}{c} - \frac{b}{c} = \frac{a-b}{c} \right\}$$

This fraction **simplifies**. To be sure of finding the **simplest form** for $\frac{138}{630}$ we write out the numerator and denominator as **prime factors**. We already know the prime factors of 630. The prime factors of 138 are $138 = 2 \cdot 3 \cdot 23$.

$$\frac{138}{630} = \frac{2 \cdot 3 \cdot 23}{2 \cdot 3 \cdot 3 \cdot 5 \cdot 7} \quad \text{one factor of 2 and one factor of 3 cancels}$$

Solution

$$\frac{27}{90} - \frac{13}{126} = \frac{23}{105}$$

Example 4

Simplify $\frac{2}{7} - \frac{-3}{4}$

Solution

The least common denominator of 4 and 7 is 28.

$$\frac{2 \cdot 4}{7 \cdot 4} + \frac{-3 \cdot 7}{4 \cdot 7} = \frac{-8}{28} + \frac{-21}{28} = \frac{8 + (-21)}{28} = -\frac{13}{28}$$

Example 5

$$-\frac{3}{12} - \frac{5}{24}$$

The least common denominator of 12 and 24 is 24.

$$\frac{-3 \cdot 2}{12 \cdot 2} + \frac{-5}{24} = \frac{-6 + -5}{24} = -\frac{11}{24}$$

Example 6

A property management firm is acquiring parcels of land in order to build a small community of condominiums. It has bought three adjacent plots of land. The first is four-fifths of an acre, the second is five-twelfths of an acre, and the third is nineteen-twentieths of an acre. The firm knows that it must allow one-sixth of an acre for utilities and a small access road. How much of the remaining land is available for development?

The first thing we need to do is extract the relevant information. Here are the relevant fractions.

$$\frac{4}{5}, \frac{5}{12}, \frac{1}{6}$$

$$\text{and } \frac{19}{20}$$

The plots of land that the firm has acquired.

The amount of land that the firm has to give up.

This sum will determine the amount of land available for development.

$$\frac{4}{5} + \frac{5}{12} + \frac{19}{20} - \frac{1}{6}$$

$$5 = 5$$

$$12 = 2 \cdot 2 \cdot 3$$

$$20 = 2 \cdot 2 \cdot 5$$

$$6 = 2 \cdot 3$$

We need to find the LCM of 5, 12, 20 and 6.

one 5

two 2's, one 3

two 2's, one 5

one 3

The smallest set of primes that encompasses all of these is 2, 2, 3, 5. Our LCD is thus $2 \cdot 2 \cdot 3 \cdot 5 = 60$

Now we can convert all fractions to a common denominator of 60. To do this, we multiply by the factors of 60 that are missing in the denominator we are converting. For example, 5 is missing two 2's and a 3. This results in $2 \cdot 2 \cdot 3 = 12$.

$$\begin{aligned} \frac{4}{5} &= \frac{12 \cdot 4}{12 \cdot 5} = \frac{48}{60} \\ \frac{5}{12} &= \frac{5 \cdot 5}{5 \cdot 12} = \frac{25}{60} \\ \frac{19}{20} &= \frac{3 \cdot 19}{3 \cdot 20} = \frac{57}{60} \\ \frac{1}{6} &= \frac{10 \cdot 1}{10 \cdot 6} = \frac{10}{60} \end{aligned}$$

Our converted sum can be rewritten as: $\frac{48}{60} + \frac{25}{60} + \frac{57}{60} - \frac{10}{60} = \frac{(48+25+57-10)}{60} = \frac{120}{60}$

Next, we need to reduce this fraction. We can see immediately that the numerator is twice the denominator. This fraction reduces to $\frac{2}{1}$ or simply two. One is sometimes called the *invisible denominator*, as every whole number can be thought of as a rational number whose denominator is one.

Solution

The property firm has two acres available for development.

Evaluate Change Using a Variable Expression

When we write algebraic expressions to represent a real quantity, the difference between two values is the **change** in that quantity.

Example 7


The speed of a train increases according to the expression $\text{speed} = 1.5t$ where speed is measured in meters per second, and “ t ” is the time measured in seconds. Find the change in the speed between $t = 2$ seconds and $t = 6$ seconds.

This function represents a train that is stopped when time equals zero ($\text{speed} = 0 \times 0.25$). As the stopwatch ticks, the train’s speed increases in a linear pattern. We can make a table of what the train’s speed is at every second.

TABLE 1.2:

Time (seconds)	Speed (m/s)
0	0
1	1.5
2	3
3	4.5
4	6
5	7.5
6	9

We can even graph this function. The graph of speed vs. time is shown here.



We wish to find the change in speed between $t = 2$ seconds and $t = 6$ seconds. There are several ways to do this. We could look at the table, and read off the speeds at 2 seconds (3 m/s) and 6 seconds (9 m/s). Or we could determine the speeds at those times by using the graph.

Another way to find the change would be to substitute the two values for t into our expression for speed. First, we will substitute $t = 2$ into our expression. To indicate that the speed we get is the speed at time = 2 seconds, we denote it as $\text{speed}(2)$.

$$\text{speed}(2) = 1.5(2) = 3$$

Next, we will substitute $t = 6$ into our expression. This is the speed at 6 seconds, so we denote it as $\text{speed}(6)$


$$\text{speed}(6) = 1.5(6) = 9$$

The change between $t = 2$ and $t = 6$ is $\text{speed}(6) - \text{speed}(2) = 9 - 3 = 6$ m/s.

The speed change is **positive**, so the change is an **increase**.

Solution

Between the two and six seconds the train's speed increases by 6 m/s.

Example 8

The intensity of light hitting a detector when it is held a certain distance from a bulb is given by this equation.

$$\text{Intensity} = 3/(\text{dist})^2$$

Where (dist) is the distance measured in **meters**, and intensity is measured in **lumens**. Calculate the change in intensity when the detector is moved from two meters to three meters away.

We first find the values of the intensity at distances of two and three meters.

$$\begin{aligned}\text{Intensity}(2) &= \frac{3}{(2)^2} = \frac{3}{4} \\ \text{Intensity}(3) &= \frac{3}{(3)^2} = \frac{3}{9} = \frac{1}{3}\end{aligned}$$

The **difference** in the two values will give the **change** in the intensity. We move **from** two meters **to** three meters away.

$$\text{Change} = \text{Intensity}(3) - \text{Intensity}(2) = \frac{1}{3} - \frac{3}{4}$$

To find the answer, we will need to write these fractions over a common denominator.

The LCM of 3 and 4 is 12, so we need to rewrite each fraction with a denominator of 12:

$$\begin{aligned}\frac{1}{3} &= \frac{4 \cdot 1}{4 \cdot 3} = \frac{4}{12} \\ \frac{3}{4} &= \frac{3 \cdot 3}{3 \cdot 4} = \frac{9}{12}\end{aligned}$$

Our change is given by this equation.

$$\frac{4}{12} - \frac{9}{12} = \frac{(4-9)}{12} = -\frac{5}{12}$$

A negative indicates that the intensity is reduced.

Solution

When moving the detector from two meters to three meters the intensity falls by $\frac{5}{12}$ lumens.

Lesson Summary

- **Subtracting** a number is the same as adding the **opposite** (or **additive inverse**) of the number.
- When **subtracting fractions**: $\frac{a}{c} - \frac{b}{c} = \frac{a-b}{c}$
- The number one is sometimes called the **invisible denominator**, as every whole number can be thought of as a rational number whose denominator is one.
- The **difference** between two values is the **change** in that quantity.

Review Questions

1. Subtract the following rational numbers. Be sure that your answer is in the **simplest form**.
 - (a) $\frac{5}{12} - \frac{9}{18}$
 - (b) $\frac{2}{3} - \frac{1}{4}$
 - (c) $\frac{3}{4} - \frac{1}{3}$
 - (d) $\frac{15}{11} - \frac{9}{7}$
 - (e) $\frac{2}{13} - \frac{-1}{11}$
 - (f) $\frac{7}{27} - \frac{19}{39}$
 - (g) $\frac{6}{11} - \frac{3}{27}$
 - (h) $\frac{13}{64} - \frac{7}{40}$
 - (i) $\frac{11}{70} - \frac{11}{30}$
2. Consider the equation $y = 3x + 2$. Determine the change in y between $x = 3$ and $x = 7$.
3. Consider the equation $y = \frac{2}{3}x + \frac{1}{2}$. Determine the change in y between $x = 1$ and $x = 2$.
4. The time taken to commute from San Diego to Los Angeles is given by the equation $\text{time} = \frac{120}{\text{speed}}$ where *time* is measured in **hours** and *speed* is measured in **miles per hour** (mph). Calculate the change in time that a rush hour commuter would see when switching from traveling by bus to train. The bus averages 40 mph to a new high speed train which averages 90 mph.

Review Answers

1.
 - (a) $\frac{-1}{12}$
 - (b) $\frac{5}{12}$
 - (c) $\frac{5}{12}$
 - (d) $\frac{6}{77}$
 - (e) $\frac{35}{143}$
 - (f) $-\frac{80}{351}$
 - (g) $\frac{9}{22}$
 - (h) $\frac{320}{9}$
 - (i) $\frac{-22}{105}$
2. Change = +12
3. Change = $-\frac{1}{3}$
4. The journey time would decrease by $1\frac{2}{3}$ hours.


1.4 Multiplication of Rational Numbers

Learning Objectives

- Multiply by negative one.
- Multiply rational numbers.
- Identify and apply properties of multiplication.
- Solve real-world problems using multiplication.

Multiplying Numbers by Negative One

Whenever we multiply a number by negative one we change the sign of the number. In more mathematical words, multiplying by negative one maps a number onto its opposite. The number line below shows the process of multiplying negative one by the numbers three and negative one.



- When we multiply a number by negative one the absolute value of the new number is the same as the absolute value of the old number. Both numbers are the same distance from zero.
- The product of a number, x , and negative one is $-x$. This does not mean that $-x$ is necessarily less than zero. If x itself is negative then $-x$ is a positive quantity because a negative times a negative is a positive.
- When we multiply an expression by negative one remember to multiply the **entire expression** by negative one.

Example 1

Multiply the following by negative one.

- 79.5
- π
- $(x + 1)$
- $|x|$

a) Solution

$$79.5 \cdot (-1) = -79.5$$

b) Solution

$$\pi \cdot (-1) = -\pi$$

c) **Solution**

$$(x + 1) \cdot (-1) = -(x + 1) = -x - 1$$

d) **Solution**

$$|x| \cdot (-1) = -|x|$$

Note that in the last case the negative sign does **not** distribute into the absolute value. Multiplying the **argument** of an absolute value equation (the term between the absolute value symbol) does not change the absolute value. $|x|$ is always positive. $|-x|$ is always positive. $-|x|$ is always negative.

Whenever you are working with expressions, you can check your answers by substituting in numbers for the variables. For example you could check part *d* of example one by letting $x = -3$.

$$|-3| \neq -|3| \text{ since } |-3| = 3 \text{ and } -|3| = -3.$$

To Multiply two numbers, multiply their absolute values.

The sign of the answer is

1. **POSITIVE** if both numbers have the same sign

$$-5 * -5 = 25$$

$$5 * 5 = 25$$

2. **NEGATIVE** if the numbers have opposite signs

$$-5 * 5 = -25$$

$$5 * -5 = -25$$

Example 2

$$\text{a. } -1 \cdot -10 = 10 \quad \text{b. } 12 \cdot -5 = -60 \quad \text{c. } -15 \cdot 5 = -75 \quad \text{d. } 7 \cdot 8 = 56$$

To Multiply Fractions or Rational Numbers

Cross Simplify Multiply the numerators together Multiply the denominators together Simplify if possible.

Example 3

$$\text{Simplify } \frac{1}{3} \cdot \frac{2}{5}$$

One way to solve this is to think of money. For example, we know that *one third of sixty dollars* is written as $\frac{1}{3} \cdot \$60$. We can read the above problem as *one-third of two-fifths*. Here is a visual picture of the fractions **one-third** and **two-fifths**.



Notice that *one-third of two-fifths* looks like the *one-third* of the shaded region in the next figure.



Here is the intersection of the two shaded regions. The whole has been divided into five pieces width-wise and three pieces height-wise. We get two pieces out of a total of fifteen pieces.

Solution

$$\frac{1}{3} \cdot \frac{2}{5} = \frac{2}{15}$$

Example 4

Simplify $\frac{3}{7} \cdot \frac{4}{5}$

We will again go with a visual representation.



We see that the whole has been divided into a total of $7 \cdot 5$ pieces. We get $3 \cdot 4$ of those pieces.

Solution

$$\frac{3}{7} \cdot \frac{4}{5} = \frac{12}{35}$$

When multiplying rational numbers, the numerators multiply together and the denominators multiply together.

When multiplying fractions $\frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd}$

Even though we have shown this result for the product of two fractions, this rule holds true when multiplying multiple fractions together.

Example 5

Multiply the following rational numbers

- $\frac{1}{2} \cdot \frac{3}{4}$
- $-\frac{2}{5} \cdot \frac{5}{9}$
- $-\frac{1}{3} \cdot -\frac{2}{7} \cdot \frac{2}{5}$
- $\frac{1}{2} \cdot \frac{2}{3} \cdot \frac{3}{4} \cdot \frac{4}{5}$

a) Solution

$$\frac{1}{2} \cdot \frac{3}{4} = \frac{1 \cdot 3}{2 \cdot 4} = \frac{3}{8}$$

b) **Solution** With this problem, we can cancel the fives.

$$-\frac{2}{5} \cdot \frac{5}{9} = \frac{-2 \cdot 5}{5 \cdot 9} = -\frac{2}{9}$$

c) **Solution** With this problem, multiply **all the numerators** and **all the denominators**.

$$\frac{-1}{3} \cdot \frac{-2}{7} \cdot \frac{2}{5} = \frac{-1 \cdot -2 \cdot 2}{3 \cdot 7 \cdot 5} = \frac{4}{105}$$

d) **Solution** With this problem, we can cancel any factor that appears as both a numerator **and** a denominator since any number divided by itself is one, according to the Multiplicative Identity Property.

$$\frac{1}{\cancel{2}} \times \frac{\cancel{2}}{\cancel{3}} \times \frac{\cancel{3}}{\cancel{4}} \times \frac{\cancel{4}}{\cancel{5}} = \frac{1}{5}$$

With multiplication of fractions, we can either simplify before we multiply or after. The next example uses factors to help simplify before we multiply.

Example 6

Evaluate and simplify $\frac{12}{25} \cdot \frac{35}{42}$

We can see that 12 and 42 are both multiples of six, and that 25 and 35 are both factors of five. We write the product again, but put in these factors so that we can cancel them prior to multiplying.

$$\frac{12}{25} \cdot \frac{35}{42} = \frac{6 \cdot 2}{25} \cdot \frac{35}{6 \cdot 7} = \frac{6 \cdot 2 \cdot 5 \cdot 7}{5 \cdot 5 \cdot 6 \cdot 7} = \frac{2}{5}$$

Solution


$$\frac{12}{25} \cdot \frac{35}{42} = \frac{2}{5}$$

Identify and Apply Properties of Multiplication

The four mathematical properties which involve multiplication are the **Commutative**, **Associative**, **Multiplicative Identity** and **Distributive Properties**.

- **Commutative property** When two numbers are multiplied together, the product is the same regardless of the order in which they are written:

Example $4 \cdot 2 = 2 \cdot 4$



We can see a geometrical interpretation of **The Commutative Property of Multiplication** to the right. The Area of the shape ($length \times width$) is the same no matter which way we draw it.

- **Associative Property** When three or more numbers are multiplied, the product is the same regardless of their grouping

Example $(2 \cdot 3) \cdot 4 = 2 \cdot (3 \cdot 4)$

- **Multiplicative Identity Property:** The product of one and any number is that number.

Example $5 \cdot 1 = 5$.

- **Distributive property** The multiplication of a number and the sum of two numbers is equal to the first number times the second number plus the first number times the third number.

Example: $4(6 + 3) = 4 \cdot 6 + 4 \cdot 3$

Example 7

Nadia and Peter are raising money by washing cars. Nadia is charging \$3 per car, and she washes five cars in the first morning. Peter charges \$5 per car (including a wax). In the first morning, he washes and waxes three cars. Who has raised the most money?

Solution

Nadia raised $5 \cdot \$3$. Peter raised $3 \cdot \$5$. According to **The Commutative Property of Multiplication**, they both raised the **same amount** of money.

Example 8

Andrew is counting his money. He puts all his money into \$10 piles. He has one pile. How much money does Andrew have?

Solution

The amount of money in each pile is \$10. The number of piles is one. The total amount of money is $\$10 \cdot 1$. According to **The Multiplicative Identity Property**, Andrew has a total of \$10.

Example 8

A gardener is planting vegetables for the coming growing season. He wishes to plant potatoes and has a choice of a single 8×7 meter plot, or two smaller plots of 3×7 meters and 5×7 meter. Which option gives him the largest area for his potatoes?



Solution

In the first option, the gardener has a total area of (8×7) .

Since $8 = (3 + 5)$ we have $(3 + 5) \cdot 7$ square meter, which equals $(3 \cdot 7) + (5 \cdot 7)$.

In the second option, the total area is $(3 \cdot 7) + (5 \cdot 7)$ square meters.

According to **The Distributive Property** both options give the gardener the same area to plant potatoes

Solve Real-World Problems Using Multiplication



Example 9

In the chemistry lab there is a bottle with two liters of a 15% solution of hydrogen peroxide (H_2O_2). John removes one-fifth of what is in the bottle, and puts it in a beaker. He measures the amount of H_2O_2 and adds twice that amount of water to the beaker. Calculate the following measurements.'

- a) The amount of H_2O_2 left in the bottle.
 - b) The amount of diluted H_2O_2 in the beaker.
 - c) The concentration of the H_2O_2 in the beaker.
- a) To determine the amount of H_2O_2 left in the bottle, we first determine the amount that was removed. That amount was $\frac{1}{5}$ of the amount in the bottle (2 liters).

$$\text{Amount removed} = \frac{1}{5} \cdot 2 \text{ liters} = \frac{2}{5} \text{ liter (or 0.4 liters)}$$

$$\text{Amount remaining} = 2 - \frac{2}{5} = \frac{10}{5} - \frac{2}{5} = \frac{8}{5} \text{ liter (or 1.6 liters)}$$

Solution

There is 1.6 liters left in the bottle.

b) We determined that the amount of the 15% H_2O_2 solution removed was $\frac{2}{5}$ liter. The amount of water added was twice this amount.

$$\text{Amount of water} = 2 \cdot \frac{2}{5} = \frac{4}{5} \text{ liter.}$$

$$\text{Total amount} = \frac{4}{5} + \frac{2}{5} = \frac{6}{5} \text{ liter (or 1.2 liters)}$$

Solution

There are 1.2 liters of diluted H_2O_2 in the beaker.

c) The new concentration of H_2O_2 can be calculated.

Initially, with $\frac{2}{5}$ of undiluted H_2O_2 there is 15% of $\frac{2}{5}$ liters of pure H_2O_2 :

$$\text{Amount of pure } H_2O_2 = 0.15 \cdot \frac{2}{5} = 0.15 \cdot 0.4 = 0.06 \text{ liter of pure } H_2O_2.$$

After dilution, this H_2O_2 is dispersed into 1.2 liters of solution. The concentration = $\frac{0.06}{1.2} = 0.05$.

To convert to a percent we multiply this number by 100.


Solution

The final of diluted H_2O_2 in the bottle is 5%.

Example 10

Anne has a bar of chocolate and she offers Bill a piece. Bill quickly breaks off $\frac{1}{4}$ of the bar and eats it. Another friend, Cindy, takes $\frac{1}{3}$ of what was left. She splits the remaining candy bar into two equal pieces which she shares with a third friend, Dora. How much of the candy bar does each person get?

First, let's look at this problem visually.



Anne starts with a full candy bar.

Bill breaks off $\frac{1}{4}$ of the bar.

Cindy takes $\frac{1}{3}$ of what was left.

Dora gets half of the remaining candy bar.

We can see that the candy bar ends up being split four ways. The sum of each piece is equal to one.

Solution

Each person gets exactly $\frac{1}{4}$ of the candy bar.

We can also examine this problem using rational numbers. We keep a running total of what fraction of the bar remains. Remember, when we read a fraction followed by *of* in the problem, it means we multiply by that fraction.

We start with one full bar of chocolate

Bill breaks off of the bar

Bill removes $\frac{1}{4} \cdot 1 = \frac{1}{4}$ of the whole bar.

Cindy takes $\frac{1}{3}$ of what is left

Cindy removes $\frac{1}{3} \cdot \frac{3}{4} = \frac{1}{4}$ of a whole bar.

Anne and Dora get two equal pieces

The total we begin with is 1.

We multiply the amount of bar(1) by $\frac{1}{4}$

The bar remaining is $1 - \frac{1}{4} = \frac{4}{4} - \frac{1}{4} = \frac{3}{4}$

We multiply the amount of bar $\left(\frac{3}{4}\right)$ by $\frac{1}{3}$

The bar remaining is $\frac{3}{4} - \frac{1}{4} = \frac{2}{4} = \frac{1}{2}$

Dora gets $\frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4}$ of a whole bar.

Anne gets the remaining $\frac{1}{2} - \frac{1}{4} = \frac{2}{4} - \frac{1}{4} = \frac{1}{4}$

Solution

Each person gets exactly $\frac{1}{4}$ of the candy bar.

Extension: If each piece that is left is 3oz, how much did the original candy bar weigh?

Lesson Summary

- When multiplying an expression by negative one, remember to multiply the **entire expression** by negative one.
- To multiply fractions $\frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd}$
- The **multiplicative properties** are:
- **Commutative property** the product of two numbers is the same whichever order the items to be multiplied are written.

Ex: $2 \cdot 3 = 3 \cdot 2$

- **Associative Property:** When three or more numbers are multiplied, the sum is the same regardless of how they are grouped.

Ex: $(2 \cdot 3) \cdot 4 = 2 \cdot (3 \cdot 4)$

- **Multiplicative Identity Property** The product of any number and one is the original number.

Ex: $2 \cdot 1 = 2$

- **Distributive property** The multiplication of a number and the sum of two numbers is equal to the first number times the second number plus the first number times the third number.

Ex: $4(2 + 3) = 4(2) + 4(3)$

Review Questions

1. Multiply the following by negative one.

- (a) 25
- (b) -105
- (c) x^2
- (d) $(3 + x)$
- (e) $(3 - x)$

2. Multiply the following rational numbers, write your answer in the **simplest form**.

- (a) $\frac{5}{12} \times \frac{9}{10}$
- (b) $\frac{2}{3} \times \frac{1}{4}$
- (c) $\frac{3}{4} \times \frac{1}{3}$
- (d) $\frac{15}{11} \times \frac{9}{7}$
- (e) $\frac{1}{13} \times \frac{1}{11}$
- (f) $\frac{7}{27} \times \frac{9}{14}$
- (g) $\left(\frac{3}{5}\right)^2$
- (h) $\frac{1}{11} \times \frac{22}{21} \times \frac{7}{10}$
- (i) $\frac{12}{15} \times \frac{35}{13} \times \frac{10}{2} \times \frac{26}{36}$

3. Three monkeys spend a day gathering coconuts together. When they have finished, they are very tired and fall asleep.

The following morning, the first monkey wakes up. Not wishing to disturb his friends, he decides to divide the coconuts into three equal piles. There is one left over, so he throws this odd one away, helps himself to his share, and goes home.

A few minutes later, the second monkey awakes. Not realizing that the first has already gone, he too divides the coconuts into three equal heaps. He finds one left over, throws the odd one away, helps himself to his fair share, and goes home.

In the morning, the third monkey wakes to find that he is alone. He spots the two discarded coconuts, and puts them with the pile, giving him a total of twelve coconuts. How many coconuts did the first and second monkey take? [**Extension:** solve by working backward]

Review Answers

1.
 - (a) -25
 - (b) 105
 - (c) $-x^2$
 - (d) $-(x + 3)$ or $-x - 3$
 - (e) $(x - 3)$
2.
 - (a) $\frac{3}{8}$
 - (b) $\frac{1}{6}$
 - (c) $\frac{1}{4}$

(d) $\frac{135}{77}$

(e) $\frac{1}{143}$

(f) $\frac{1}{6}$

(g) $\frac{27}{125}$

(h) $\frac{1}{15}$

(i) $\frac{10}{9}$

3. The first monkey takes eight coconuts. The second monkey takes five coconuts.

1.5 The Distributive Property

Learning Objectives

- Apply the distributive property.
- Identify parts of an expression.
- Solve real-world problems using the distributive property.

Introduction

At the end of the school year, an elementary school teacher makes a little gift bag for each of his students. Each bag contains one class photograph, two party favors and five pieces of candy. The teacher will distribute the bags among his 28 students. How many of each item does the teacher need?

Apply the Distributive Property

When we have a problem like the one posed in the introduction, **The Distributive Property** can help us solve it. To begin, we can write an expression for the contents of each bag.

$$\text{Contents} = (\text{photo} + 2 \text{ favor} + 5 \text{ candy})$$

$$\text{Contents} = (p + 2f + 5c) \quad \text{We may even use single letter variables to write an expression.}$$

We know that the teacher has 28 students, therefore we can write the following expression for the number of items that the teacher will need.

$$\text{Items} = 28 \cdot (p + 2f + 5c) \quad \text{28 times the individual contents of each bag.}$$

We generally omit any multiplication signs that are not strictly necessary.

$$\text{Items} = 28(p + 2f + 5c)$$

The Distributive Property of Multiplication means that when faced with a term multiplying other terms inside parentheses, the outside term multiplies with each of the terms inside the parentheses.



$$28(p + 2f + 5c)$$

$$= 28(p + 2f + 5c) = 28(p) + 28(2f) + 28(5c) = 28p + 56f + 140c$$

So the teacher needs 28 class photos, 56 party favors and 140 pieces of candy.

The Distributive Property works when we have numbers inside the parentheses. You can see this by looking at a simple problem and considering the **Order of Operations**.

Example 1

Determine the value of $11(2+6)$ using both Order of Operations and the Distributive Property.

First, we consider the problem with the Order of Operations – PEMDAS dictates that we evaluate the amount inside the parentheses first.

Solution

$$11(2+6) = 11(8) = 88$$

Next we will use the Distributive Property. We multiply the 11 by each term inside the parentheses.

Solution

$$11(2+6) = 11(2) + 11(6) = 22 + 66 = 88$$

Example 2

Determine the value of $11(2-6)$ using both the Order of Operations and the Distributive Property.

First, we consider the Order of Operations and evaluate the amount inside the parentheses first.

Solution

$$11(2-6) = 11(-4) = -44$$

Next, the Distributive Property.

Solution

$$11(2-6) = 11(2) + 11(-6) = 22 - 66 = -44$$

Note When applying the Distributive Property you **MUST** take note of any **negative signs!**

Example 3

Use the Distributive Property to determine the following.

a) $11(2x+6)$

b) $7(3x-5)$

c) $\frac{2}{7}(3y^2-11)$

d) $\frac{2x}{7}\left(3y^2-\frac{11}{xy}\right)$

a) Simply multiply each term by 11.

Solution

$$11(2x+6) = 22x + 66$$

b) Note the negative sign on the second term.

Solution

$$7(3x - 5) = 21x - 35$$

$$c) \frac{2}{7}(3y^2 - 11) = \frac{2}{7}(3y^2) + \frac{2}{7}(-11) = \frac{6y^2}{7} - \frac{22}{7}$$

Solution

$$\frac{2}{7}(3y^2 - 11) = \frac{6y^2 - 22}{7}$$

$$d) \frac{2x}{7}\left(3y^2 - \frac{11}{xy}\right) = \frac{2x}{7}(3y^2) + \frac{2x}{7}\left(\frac{-11}{xy}\right) = \frac{6x^2y - 22x}{7xy}$$

Solution

$$\frac{2x}{7}\left(3y^2 - \frac{11}{xy}\right) = \frac{6xy^3 - 22}{7y}$$

Identify Expressions That Involve the Distributive Property

The Distributive Property often appears in expressions, and many times it does not involve parentheses as grouping symbols. In Lesson 1.2, we saw how the fraction bar acts as a grouping symbol. The following example involves using the Distributive Property with fractions.

Example 4

Simplify the following expressions.

a) $\frac{2x+8}{4}$

b) $\frac{9y-2}{3}$

c) $\frac{z+6}{2}$

Even though these expressions are not written in a form we usually associate with the Distributive Property, the fact that the numerator of fractions should be treated as if it were in parentheses makes this a problem that the Distributive Property can help us solve.

a) $\frac{2x+8}{4}$ can be re-written as $\frac{1}{4}(2x+8)$.

We can then proceed to distribute the $\frac{1}{4}$.

$$\frac{1}{4}(2x+8) = \frac{2x}{4} + \frac{8}{4} = \frac{2x}{2 \cdot 2} + \frac{4 \cdot 2}{4}$$

Solution

$$\frac{2x+8}{4} = \frac{x}{2} + 2$$

b) $\frac{9y-2}{3}$ can be re-written as $\frac{1}{3}(9y-2)$.

We can then proceed to distribute the $\frac{1}{3}$.

$$\frac{1}{3}(9y-2) = \frac{9y}{3} - \frac{2}{3} = \frac{\cancel{3} \cdot 3x}{\cancel{3}} - \frac{2}{3}$$

Solution

$$\frac{9y-2}{3} = 3y - \frac{2}{3}$$

c) $\frac{z+6}{2}$ can be re-written as $\frac{1}{2}(z+6)$.

We can then proceed to distribute the $\frac{1}{2}$.

$$\frac{1}{2}(z+6) = \frac{z}{2} + \frac{6}{2}$$

Solution

$$\frac{z+6}{2} = \frac{z}{2} + 3$$

Solve Real-World Problems Using the Distributive Property

The Distributive Property is one of the most common mathematical properties to be seen in everyday life. It crops up in business and in geometry. Anytime we have two or more groups of objects, the Distributive Property can help us solve for an unknown.



Example 5

An octagonal gazebo is to be built as shown right. Building code requires five foot long steel supports to be added along the base and four foot long steel supports to be added to the roof-line of the gazebo. What length of steel will be required to complete the project?

Each side will require two lengths, one of five and four feet respectively. There are eight sides, so here is our equation.

$$\text{Steel required} = 8(4 + 5) \text{ feet.}$$

We can use the distributive property to find the total amount of steel:

$$\text{Steel required} = 8 \times 4 + 8 \times 5 = 32 + 40 \text{ feet.}$$

Solution

A total of 72 feet of steel is required for the project.

**Example 6**

Each student on a field trip into a forest is to be given an emergency survival kit. The kit is to contain a flashlight, a first aid kit, and emergency food rations. Flashlights cost \$12 each, first aid kits are \$7 each and emergency food rations cost \$2 per day. There is \$500 available for the kits and 17 students to provide for. How many days worth of rations can be provided with each kit?

The unknown quantity in this problem is the number of days' rations. This will be x in our expression. Each kit will contain the following items.

- 1 · \$12 flashlight.
- 1 · \$7 first aid kit.
- x · \$2 daily rations.

The number of kits = 17, so the total cost is equal to the following equation.

$$\text{Total cost} = 17(12 + 7 + 2x)$$

We can use the Distributive Property on this expression.

$$17(12 + 7 + 2x) = 204 + 119 + 34x$$

We know that there is \$500 available to buy the kits. We can substitute the cost with the money available.

$$204 + 119 + 34x = 500$$

The sum of the numbers on the left equal to the money available

$$323 + 34x = 500$$

Subtract 323 from both sides

$$-323 - 323$$

$$34x = 177$$

Divide both sides by 34

$$x = 5.20588\dots$$

Since this represents the number of daily rations that can be bought, we must **round to the next lowest whole number**. We wouldn't have enough money to buy a sixth day of supplies.

Solution

Five days worth of emergency rations can be purchased for each survival kit.

Lesson Summary

- **Distributive Property** The multiplication of a number and the sum of two numbers is equal to the first number times the second number plus the first number times the third number.

Ex: $4 \times (6 + 3) = 4 \times 6 + 4 \times 3$

- When applying the Distributive Property you **MUST** take note of any **negative signs!**

Review Questions

1. Use the Distributive Property to simplify the following expressions.

- $(x + 4) - 2(x + 5)$
- $\frac{1}{2}(4z + 6)$
- $(4 + 5) - (5 + 2)$
- $(x + 2 + 7)$
- $y(x + 7)$
- $13x(3y + z)$

2. Use the Distributive Property to remove the parentheses from the following expressions.

- $\frac{1}{2}(x - y) - 4$
- $0.6(0.2x + 0.7)$
- $6 + (x - 5) + 7$
- $6 - (x - 5) + 7$
- $4(m + 7) - 6(4 - m)$
- $-5(y - 11) + 2y$

3. Use the Distributive Property to simplify the following fractions.

- $\frac{8x+12}{4}$
- $\frac{9x+12}{3}$
- $\frac{11x+12}{2}$
- $\frac{3y+2}{6}$
- $-\frac{6z-2}{3}$
- $\frac{7-6p}{3}$

4. A bookshelf has five shelves, and each shelf contains seven poetry books and eleven novels. How many of each type of book does the bookcase contain?

5. Amar is making giant holiday cookies for his friends at school. He makes each cookie with 6 oz of cookie dough and decorates them with macadamia nuts. If Amar has 5 lbs of cookie dough (1 lb = 16 oz) and 60 macadamia nuts, calculate the following.

- How many (**full**) cookies he can make?
- How many macadamia nuts he can put on each cookie, if each is to be identical?

Review Answers

- $-x - 6$
 - $2z + 3$
 - 2
 - $x + 9$
 - $xy + 7y$

- (f) $39xy + 13xz$
2. (a) $\frac{x}{2} - \frac{y}{2} - 4$
(b) $0.12x + 0.42$
(c) $x + 8$
(d) $18 - x$
(e) $10m + 4$
(f) $55 - 3y$
3. (a) $2x + 3$
(b) $3x + 4$
(c) $\frac{11x}{2} + 6$
(d) $\frac{y}{2} + \frac{1}{3}$
(e) $\frac{2}{3} - 2z$
(f) $\frac{7}{3} - 2p$
4. The bookshelf contains 35 poetry books and 55 novels.
5. (a) Amar can make 13 cookies (2 oz leftover).
(b) Each cookie has 4 macadamia nuts (8 left over).

1.6 Division of Rational Numbers

Learning Objectives

- Find multiplicative inverses.
- Divide rational numbers.
- Solve real-world problems using division.

Introduction – Identity elements

An **identity element** is a number which, when combined with a mathematical operation on a number, leaves that number unchanged. For addition and subtraction, the **identity element** is **zero**.

$$\begin{aligned}2 + 0 &= 2 \\ -5 + 0 &= -5 \\ 99 - 0 &= 99\end{aligned}$$

The inverse operation of addition is subtraction.

$$x + 5 - 5 = x \quad \text{When we subtract what we have added, we get back to where we started!}$$

When you add a number to its **opposite**, you get the identity element for addition.

$$5 + (-5) = 0$$

You can see that the **addition of an opposite is an equivalent operation to subtraction**.

For multiplication and division, the **identity element** is **one**.

$$\begin{aligned}2 \times 1 &= 2 \\ -5 \times 1 &= -5 \\ 99 \div 1 &= 99\end{aligned}$$

In this lesson, we will learn about **multiplying by a multiplicative inverse** as an equivalent operation to division. Just as we can use **opposites** to turn a **subtraction** problem into an **addition** problem, we can use **reciprocals** to turn a **division** problem into a **multiplication** problem.

Find Multiplicative Inverses

The **multiplicative inverse** of a number, x , is the number when multiplied by x yields **one**. In other words, any number times the multiplicative inverse of that number equals one. The multiplicative inverse is commonly the reciprocal, and the multiplicative inverse of x is denoted by $\frac{1}{x}$.

Look at the following multiplication problem:

$$\text{Simplify } \frac{2}{3} \times \frac{3}{2}$$

We know that we can cancel terms that appear on both the numerator and the denominator. Remember we leave a one when we cancel all terms on either the numerator or denominator!

$$\frac{2}{3} \times \frac{3}{2} = \frac{\cancel{2}}{\cancel{3}} \times \frac{\cancel{3}}{\cancel{2}} = 1$$

It is clear that $\frac{2}{3}$ is the multiplicative inverse of $\frac{3}{2}$. Here is the rule.

To find the multiplicative inverse of a rational number, we simply **invert the fraction**.

The multiplicative inverse of $\frac{a}{b}$ is $\frac{b}{a}$, as long as $a \neq 0$

Example 1

Find the multiplicative inverse of each of the following.

a) $\frac{3}{7}$

b) $\frac{4}{7}$

c) $3\frac{1}{2}$

d) $-\frac{x}{y}$

e) $\frac{1}{11}$

a) Solution

The multiplicative inverse of $\frac{3}{7}$ is $\frac{7}{3}$.

b) Solution

The multiplicative inverse of $\frac{4}{9}$ is $\frac{9}{4}$.

c) To find the multiplicative inverse of $3\frac{1}{2}$ we first need to convert $3\frac{1}{2}$ to an **improper fraction**:

$$3\frac{1}{2} = \frac{3}{1} + \frac{1}{2} = \frac{6}{2} + \frac{1}{2} = \frac{7}{2}$$

Solution

The multiplicative inverse of $3\frac{1}{2}$ is $\frac{2}{7}$.

d) Do not let the negative sign confuse you. The multiplicative inverse of a negative number is also negative!

Solution

The multiplicative inverse of $-\frac{x}{y}$ is $-\frac{y}{x}$.

e) The multiplicative inverse of $\frac{1}{11}$ is $\frac{11}{1}$. Remember that when we have a denominator of one, we omit the denominator.

Solution

The multiplicative inverse of $\frac{1}{11}$ is 11.

Look again at the last example. When we took the multiplicative inverse of $\frac{1}{11}$ we got a whole number, 11. This, of course, is expected. We said earlier that the multiplicative inverse of x is $\frac{1}{x}$.

The multiplicative inverse of a whole number is one divided that number.

Remember the idea of the **invisible denominator**. The idea that every integer is actually a rational number whose denominator is one. $5 = \frac{5}{1}$.

Divide Rational Numbers

Division can be thought of as the inverse process of multiplication. If we multiply a number by seven, we can divide the answer by seven to return to the original number. Another way to return to our original number is to multiply the answer by the **multiplicative inverse of seven**.

In this way, we can simplify the process of dividing rational numbers. We can turn a division problem into a multiplication process by replacing the divisor (the number we are dividing by) with its multiplicative inverse, or **reciprocal**.

To divide rational numbers, invert the divisor and multiply $\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c}$.

Also, $\frac{\frac{a}{b}}{\frac{c}{d}} = \frac{a}{b} \cdot \frac{d}{c}$

Example 2

Divide the following rational numbers, giving your answer in the **simplest form**.

a) $\frac{1}{2} \div \frac{1}{4}$

b) $\frac{7}{3} \div \frac{2}{3}$

c) $\frac{x}{2} \div \frac{1}{4y}$

d) $\frac{11}{2x} \div \left(-\frac{x}{y}\right)$

a) Replace $\frac{1}{4}$ with $\frac{4}{1}$ and multiply. $\frac{1}{2} \times \frac{4}{1} = \frac{1}{2} \times \frac{2 \cdot 2}{2} = \frac{1}{2}$.

Solution

$$\frac{1}{2} \div \frac{1}{4} = 2$$

b) Replace $\frac{2}{3}$ with $\frac{3}{2}$ and multiply. $\frac{7}{3} \times \frac{3}{2} = \frac{7}{2}$.

Solution

$$\frac{7}{3} \div \frac{2}{3} = \frac{7}{2}$$

c) Replace $\frac{1}{4y}$ with $\frac{4y}{1}$ and multiply. $\frac{x}{2} \times \frac{4y}{1} = \frac{x}{2} \times \frac{2 \cdot 2 \cdot y}{1} = \frac{x \cdot 2y}{1}$

Solution

$$\frac{x}{2} \div \frac{1}{4y} = 2xy$$

d) Replace $\left(-\frac{x}{y}\right)$ with $\left(-\frac{y}{x}\right)$ and multiply. $\frac{11}{2x} \times \left(-\frac{y}{x}\right) = -\frac{11 \cdot y}{2x \cdot x}$.

Solution

$$\frac{11}{2x} \left(-\frac{x}{y}\right) = -\frac{11y}{2x^2}$$

Solve Real-World Problems Using Division

Speed, Distance and Time

An object moving at a certain **speed** will cover a fixed **distance** in a set **time**. The quantities speed, distance and time are related through the equation:

$$\text{Speed} = \frac{\text{Distance}}{\text{Time}}$$

Example 3

Andrew is driving down the freeway. He passes mile marker 27 at exactly mid-day. At 12:35 he passes mile marker 69. At what speed, in miles per hour, is Andrew traveling?

To determine speed, we need the distance traveled and the time taken. If we want our speed to come out in miles per hour, we will need distance in **miles** and time in **hours**.

$$\begin{aligned}\text{Distance} &= 69 - 27 = 42 \text{ miles} \\ \text{Time taken} &= 35 \text{ minutes} = \frac{35}{60} = \frac{\cancel{7} \cdot 5}{\cancel{2} \cdot 3 \cdot 2} = \frac{7}{12} \text{ hour}\end{aligned}$$

We now *plug in* the values for distance and time into our equation for speed.

$$\begin{aligned}\text{Speed} &= \frac{42}{\left(\frac{7}{12}\right)} = \frac{42}{1} \div \frac{7}{12} && \text{Replace } \frac{7}{12} \text{ with } \frac{12}{7} \text{ and multiply.} \\ \text{Speed} &= \frac{42}{1} \times \frac{12}{7} = \frac{\cancel{7} \cdot 6 \cdot 12}{\cancel{7}} = \frac{6 \cdot 12}{1}\end{aligned}$$

Solution

Andrew is driving at 72 miles per hour .

Example 4

Anne runs a mile and a half in a quarter hour. What is her speed in miles per hour?

We already have the distance and time in the correct units (miles and hours). We simply write each as a rational number and plug them into the equation.

$$\begin{aligned}\text{Speed} &= \frac{\left(\frac{3}{2}\right)}{\left(\frac{1}{4}\right)} = \frac{3}{2} \div \frac{1}{4} && \text{Replace } \frac{1}{4} \text{ with } \frac{4}{1} \text{ and multiply.} \\ \text{Speed} &= \frac{3}{2} \times \frac{4}{1} = \frac{12}{2} = 6\end{aligned}$$

Solution

Anne runs at 6 miles per hour.

Example 5 – Newton’s Second Law

Newton’s second law ($F = ma$) relates the force applied to a body (F), the mass of the body (m) and the acceleration (a). Calculate the resulting acceleration if a Force of $7\frac{1}{3}$ Newtons is applied to a mass of $\frac{1}{5}$ kg.

First, we rearrange our equation to isolate the acceleration, a

$$a = \frac{F}{m}$$

Substitute in the known values.

$$a = \frac{\left(7\frac{1}{3}\right)}{\left(\frac{1}{5}\right)} = \left(\frac{7.3}{3} + \frac{1}{3}\right) \div \left(\frac{1}{5}\right)$$

Determine improper fraction, then invert $\frac{1}{5}$ and multiply.

$$a = \frac{22}{3} \times \frac{5}{1} = \frac{110}{3}$$

Solution

The resultant acceleration is $36\frac{2}{3}$ m/s².

Lesson Summary

- The **multiplicative inverse** of a number is the number which produces one when multiplied by the original number. The multiplicative inverse of x is the reciprocal $\frac{1}{x}$.
- To find the multiplicative inverse of a rational number, we simply **invert the fraction**: $\frac{a}{b}$ inverts to $\frac{b}{a}$.
- To divide rational numbers, invert the divisor and multiply $\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \times \frac{d}{c}$.

Review Questions

- Find the multiplicative inverse of each of the following.
 - 100
 - $\frac{2}{8}$
 - $-\frac{19}{21}$
 - 7
 - $-\frac{x^3}{2xy^2}$
- Divide the following rational numbers, be sure that your answer is in the simplest form.
 - $\frac{5}{2} \div \frac{1}{4}$
 - $\frac{1}{2} \div \frac{7}{9}$
 - $\frac{5}{11} \div \frac{6}{7}$
 - $\frac{1}{2} \div \frac{1}{2}$
 - $-\frac{x}{2} \div \frac{5}{7}$
 - $\frac{1}{2} \div \frac{x}{4y}$
 - $\left(-\frac{1}{3}\right) \div \left(-\frac{3}{5}\right)$
 - $\frac{7}{2} \div \frac{7}{4}$
 - $11 \div \left(-\frac{x}{4}\right)$
- The label on a can of paint states that it will cover 50 square feet per pint. If I buy a $\frac{1}{8}$ pint sample, it will cover a square two feet long by three feet high. Is the coverage I get more, less or the same as that stated on the label?
- The world's largest trench digger, "Bagger 288", moves at $\frac{3}{8}$ mph. How long will it take to dig a trench $\frac{2}{3}$ mile long?
- A $\frac{2}{7}$ Newton force applied to a body of unknown mass produces an acceleration of $\frac{3}{10}$ m/s². Calculate the mass of the body. Note: Newton = kg m/s².

Review Answers

1. (a) $\frac{1}{101}$
(b) $\frac{2}{8}$
(c) $-\frac{21}{19}$
(d) $\frac{1}{7}$
(e) $-\frac{2xy^2}{z^3}$
2. (a) 10
(b) $\frac{9}{14}$
(c) $\frac{35}{66}$
(d) 1
(e) $-\frac{7x}{10}$
(f) $\frac{2y}{x}$
(g) $\frac{5}{9}$
(h) 2
(i) $-\frac{44}{x}$
3. At 48 square feet per pint I get less coverage.
4. Time = $\frac{16}{9}$ hour (1 hr 46 min 40 sec)
5. mass = $\frac{20}{21}$ kg

1.7 Square Roots and Real Numbers

Learning Objectives

- Find square roots.
- Approximate square roots.
- Identify irrational numbers.
- Classify real numbers.
- Graph and order real numbers.


Find Square Roots

The square root of a number is a number which, when multiplied by itself gives the original number. In algebraic terms, the square root of x is a number, b , such that $b^2 = x$.

Note: There are two possibilities for a numerical value for b . The **positive** number that satisfies the equation $b^2 = x$ is called the **principal square root**. Since $(-b) \cdot (-b) = +b^2 = x$, $-b$ is also a valid solution.

The square root of a number, x , is written as \sqrt{x} or sometimes as $\pm\sqrt{x}$. For example, $2^2 = 4$, so the square root of 4, $\sqrt{4} = \pm 2$.

Some numbers, like 4, have integer square roots. Numbers with integer square roots are called **perfect squares**. The first five perfect squares (1, 4, 9, 16, 25) are shown below.



You can determine whether a number is a perfect square by looking at its prime factors. If every number in the factor tree appears an even number of times, the number is a perfect square. Further, to find the square root of that number, simply take one of each pair of factors and multiply them together.

Example 1

Find the principal square root of each of these perfect squares.

- 121
 - 225
 - 324
 - 576
- a) $121 = 11 \times 11$

Solution

$$\sqrt{121} = 11$$

b) $225 = (5 \times 5) \times (3 \times 3)$

Solution

$$\sqrt{225} = 5 \times 3 = 15$$

$$c) 324 = (2 \times 2) \times (3 \times 3) \times (3 \times 3)$$

Solution

$$\sqrt{324} = 2 \times 3 \times 3 = 18$$

$$d) 576 = (2 \times 2) \times (2 \times 2) \times (2 \times 2) \times (3 \times 3)$$

Solution

$$\sqrt{576} = 2 \times 2 \times 2 \times 3 = 24$$

Approximate Square Roots

When we have perfect squares, we can write an exact numerical solution for the principal square root. When we have one or more unpaired primes in the factor tree of a number, however, we do not get integer values for the square root and we have seen that we leave a radical in the answer. Terms like $\sqrt{2}$, $\sqrt{3}$ and $\sqrt{7}$ (square roots of prime numbers) cannot be written as **rational numbers**. That is to say, they cannot be expressed as the ratio of two integers. We call them **irrational numbers**. In decimal form, they have an unending, seemingly random, string of numbers after the decimal point.

To find approximate values for square roots, we use the $\sqrt{\quad}$ or \sqrt{x} button on a calculator. When the number we are finding the square root of is a perfect square, or the square of a rational number, we will get an exact answer. When the number is a non-perfect square, the decimals will appear random and we will have an irrational number as our answer. We call this an **approximate answer**. Even though we may have an answer to eight or nine decimal places, it still represents an **approximation** of the real answer which has an **infinite number of non-repeating decimals**.

Example 4

Use a calculator to find the following square roots. Round your answer to three decimal places.

$$a) \sqrt{99}$$

$$b) \sqrt{5}$$

$$c) \sqrt{0.5}$$

$$d) \sqrt{1.75}$$

a) The calculator returns 9.949874371.

Solution

$$\sqrt{99} \approx 9.950$$

b) The calculator returns 2.236067977.

Solution

$$\sqrt{5} \approx 2.236$$

c) The calculator returns 0.7071067812 .

Solution

$$\sqrt{0.5} \approx 0.707$$

d) The calculator returns 1.322875656.

Solution

$$\sqrt{1.75} \approx 1.323$$

Identify Irrational Numbers

Any square root that cannot be simplified to a form without a square root is **irrational**, but **not all** square roots are irrational. For example, $\sqrt{49} = 7$. This is a terminating decimal, which makes $\sqrt{49}$ **rational**, but $\sqrt{50}$ does not simplify perfectly. $\sqrt{50} \approx 7.071067812$. The fact that it is a non-terminating non-repeating decimal makes $\sqrt{50}$ **irrational**.

Example 5

Identify which of the following are rational numbers and which are irrational numbers.

a) 23.7

b) 2.8956

c) π

d) $\sqrt{6}$

e) $3.\bar{27}$

a) $23.7 = 23\frac{7}{10}$. This is clearly a **rational number**.

b) $2.8956 = 2\frac{8956}{10000}$. Again, this is a **rational number**.


c) $\pi = 3.141592654\dots$ The decimals appear random, and from the definition of π we know they do not repeat. This is an **irrational number**.

d) $\sqrt{6} = 2.44949489743\dots$ Again the decimals appear to be random. We also know that $\sqrt{6} = \sqrt{2} \times \sqrt{3}$. Square roots of **primes** are irrational. $\sqrt{6}$ is an irrational number.

e) $3.\bar{27} = 3.2727272727\dots$ Although these decimals are recurring they are certainly *not* unpredictable. This is a **rational number** (in actual fact, $3.\bar{27} = \frac{36}{11}$)

Classify Real Numbers

We can now see how real numbers fall into one of several categories.



If a real number can be expressed as a rational number, it falls into one of two categories. If the denominator of its **simplest form** is one, then it is an **integer**. If not, it is a fraction (this term is used here to also include decimals, as $3.27 = 3\frac{27}{100}$). **Rational numbers will always be terminating decimals or non-terminating repeating decimals.**

If the number cannot be expressed as the ratio of two integers (i.e. as a fraction), it is **irrational**. **Irrational numbers will always be non-terminating, non-repeating decimals.**

Example 6

Classify the following real numbers.

a) 0

b) -1

c) $\frac{\pi}{3}$

d) $\frac{\sqrt{2}}{3}$

e) $\frac{\sqrt{36}}{9}$

a) **Solution**

Zero is an **integer**.

b) **Solution**

-1 is an **integer**.

c) Although $\frac{\pi}{3}$ is written as a fraction, the numerator (π) is irrational.

Solution

$\frac{\pi}{3}$ is an **irrational number**.

d) $\frac{\sqrt{2}}{3}$ cannot be simplified to remove the square root.

Solution

$\frac{\sqrt{2}}{3}$ is an **irrational number**.

e) $\frac{\sqrt{36}}{9}$ can be simplified to $\frac{\sqrt{36}}{3} = \frac{6}{9} = \frac{2}{3}$ **Solution**

$\frac{\sqrt{36}}{9}$ is a **rational number**.

Graph and Order Real Numbers

We have already talked about plotting integers on the number line. It gives a visual representation of which number is bigger, smaller, etc. It would therefore be helpful to plot non-integer rational numbers (fractions) on the number line also. There are two ways to graph rational numbers on the number line. You can convert them to a mixed number (graphing is one of the few instances in algebra when mixed numbers are preferred to improper fractions), or you can convert them to decimal form.

Example 7

Plot the following rational numbers on the number line.

a) $\frac{2}{3}$


b) $-\frac{3}{7}$

c) $\frac{17}{3}$


d) $\frac{57}{16}$

If we divide the intervals on the number line into the number on the denominator, we can look at the fraction's numerator to determine how many of these **sub-intervals** we need to include.


a) $\frac{2}{3}$ falls between 0 and 1. We divide the interval into three units, and include two sub-intervals.




b) $-\frac{3}{7}$ falls between 0 and -1 . We divide the interval into seven units, and move **left** from zero by three sub-intervals.



c) $\frac{17}{5}$ as a mixed number is $2\frac{2}{5}$ and falls between 3 and 4. We divide the interval into five units, and move over two sub-intervals.



d) $\frac{57}{16}$ as a mixed number is $3\frac{9}{16}$ and falls between 3 and 4. We need to make sixteen sub-divisions.



Example 8


Plot the following numbers, in the correct order, on a number line.

- π
- $\frac{22}{7}$
- 3.14
- $\sqrt{10}$

We will use a calculator to find decimal expansions for each of these, and use a number line divided into 1000 sub-divisions. When we have two extremely close numbers, we will ensure that we place them in the correct order by looking at the expansion to the 3rd decimal place and writing as a fraction of 1000.

- $\pi = 3.14159\dots \approx 3\frac{142}{1000}$
- $\frac{22}{7} = 3.14288\dots \approx 3\frac{143}{1000}$
- $3.14 \approx 3\frac{140}{1000}$
- $\sqrt{10} = 3.16227\dots \approx 3\frac{162}{1000}$

Solution



Lesson Summary

- The **square root** of a number is a number which gives the original number when multiplied by itself. In algebraic terms, the square root of x is a number, b , such that $b^2 = x$, or $b = \sqrt{x}$

- There are two possibilities for a numerical value for b . A positive value called the **principal square root** and a negative value (the opposite of the positive value).
- A **perfect square** is a number with an integer square root.
- Here are some mathematical properties of square roots.

$$\begin{aligned}\sqrt{a} \times \sqrt{b} &= \sqrt{ab} & \sqrt{a} \div \sqrt{b} &= \sqrt{\frac{a}{b}} \\ A\sqrt{a} \times B\sqrt{b} &= AB\sqrt{ab} & A\sqrt{a} \div B\sqrt{b} &= \frac{A}{B}\sqrt{\frac{a}{b}}\end{aligned}$$

- Square roots of prime numbers are **irrational numbers**. They cannot be written as rational numbers (the ratio of two integers). In decimal form, they have an unending, seemingly random, string of numbers after the decimal point.
- Computing a square root on a calculator will produce an **approximate solution** since there are a finite number of digits after the decimal point.

Review Questions

- Find the following square roots **exactly without using a calculator**, giving your answer in the simplest form.
 - $\sqrt{25}$
 - $\sqrt{24}$
 - $\sqrt{20}$
 - $\sqrt{200}$
 - $\sqrt{2000}$
 - $\sqrt{\frac{1}{4}}$
 - $\sqrt{\frac{9}{4}}$
 - $\sqrt{0.16}$
 - $\sqrt{0.1}$
 - $\sqrt{0.01}$
- Use a calculator to find the following square roots. Round to two decimal places.
 - $\sqrt{13}$
 - $\sqrt{99}$
 - $\sqrt{123}$
 - $\sqrt{2}$
 - $\sqrt{2000}$
 - $\sqrt{0.25}$
 - $\sqrt{1.35}$
 - $\sqrt{0.37}$
 - $\sqrt{0.7}$
 - $\sqrt{0.01}$
- Classify the following numbers as an integer, a rational number or an irrational number.
 - $\sqrt{0.25}$

- (b) $\sqrt{1.35}$
- (c) $\sqrt{20}$
- (d) $\sqrt{25}$
- (e) $\sqrt{100}$

4. Place the following numbers in numerical order, from lowest to highest.


$$\frac{\sqrt{6}}{2}$$

$$\frac{61}{50}$$

$$\sqrt{1.5}$$

$$\frac{16}{13}$$

5. Use the marked points on the number line and identify each proper fraction.



Review Answers

1. (a) 5
 (b) $2\sqrt{6}$
 (c) $2\sqrt{5}$
 (d) $10\sqrt{2}$
 (e) $20\sqrt{5}$
 (f) $\frac{1}{2}$
 (g) $\frac{3}{2}$
 (h) 0.4
 (i) $\frac{1}{\sqrt{10}}$ or $\frac{\sqrt{10}}{10}$
 (j) 0.1
2. (a) 3.61
 (b) 9.95
 (c) 11.09
 (d) 1.41
 (e) 44.72
 (f) 0.5
 (g) 1.16
 (h) 0.61
 (i) 0.84
 (j) 0.1
3. (a) rational
 (b) irrational
 (c) irrational
 (d) integer
 (e) integer

4.

$$\frac{61}{50}$$

$$\frac{\sqrt{6}}{2}$$

$$\frac{16}{13}$$

$$\sqrt{1.6}$$

5. (a) $\frac{2}{3}$
 (b) $\frac{3}{5}$
 (c) $\frac{4}{9}$
 (d) $\frac{34}{9}$

CHAPTER **2** Expressions and Equations

Chapter Outline

- 2.1** VARIABLE EXPRESSIONS
 - 2.2** ORDER OF OPERATIONS
 - 2.3** ONE-STEP EQUATIONS
 - 2.4** TWO-STEP EQUATIONS
 - 2.5** MULTI-STEP EQUATIONS
 - 2.6** EQUATIONS WITH VARIABLES ON BOTH SIDES
 - 2.7** RATIOS AND PROPORTIONS
-

2.1 Variable Expressions

Learning Objectives

- Evaluate algebraic expressions.
- Evaluate algebraic expressions with exponents.

Introduction The Language of Algebra

Do you like to do the same problem over and over again? No? Well, you are not alone. **Algebra** was invented by mathematicians so that they could solve a problem once and then use that solution to solve a group of similar problems. The big idea of algebra is that once you have solved one problem you can **generalize** that solution to solve other similar problems.

In this course, we'll assume that you can already do the basic operations of arithmetic. In arithmetic, only numbers and their arithmetical operations (such as $+$, \times , \div) occur. In algebra, numbers (and sometimes processes) are denoted by symbols (such as x, y, a, b, c, \dots). These symbols are called **variables**.


The letter x , for example, will often be used to represent some number. The value of x , however, is not fixed from problem to problem. The letter x will be used to represent a number which may be unknown (and for which we may have to solve) or it may even represent a quantity which varies within that problem.

Using variables offers advantages over solving each problem from scratch:

- It allows the general formulation of arithmetical laws such as $a + b = b + a$ for all real numbers a and b .
- It allows the reference to unknown numbers, for instance: Find a number x such that $3x + 1 = 10$.
- It allows short-hand writing about functional relationships such as, If you sell x tickets, then your profit will be $3x - 10$ dollars, or $f(x) = 3x - 10$, where f is the profit function, and x is the input (i.e. how many tickets you sell).

Example 1

Write an algebraic expression for the perimeter and area of the rectangle as follows.



To find the perimeter, we add the lengths of all 4 sides. We can start at the top-left and work clockwise. The perimeter, P , is therefore:

$$P = l + w + l + w$$

We are adding 2 l 's and 2 w 's. Would say that:

$$P = 2 \times l + 2 \times w$$

You are probably familiar with using \cdot instead of \times for multiplication, so you may prefer to write:

$$P = 2 \cdot l + 2 \cdot w$$

It's customary in algebra to omit multiplication symbols whenever possible. For example, $11x$ means the same thing as $11 \cdot x$ or $11 \times x$. We can therefore write the expression for P as:

$$P = 2l + 2w$$

Area is *length multiplied by width*. In algebraic terms we get the expression:

$$A = 1 \times w \quad \rightarrow \quad A = 1 \cdot w \quad \rightarrow \quad A = 1w$$

Note: An example of a **variable expression** is $2l + 2w$; an example of an **equation** is $P = 2l + 2w$. The main difference between equations and expressions is the presence of an equals sign ($=$).

In the above example, there is no simpler form for these equations for the perimeter and area. They are, however, perfectly general forms for the perimeter and area of a rectangle. They work whatever the numerical values of the length and width of some particular rectangle are. We would simply substitute values for the length and width of a **real** rectangle into our equation for perimeter and area. This is often referred to as substituting (or **plugging in**) values. In this chapter we will be using the process of substitution to evaluate expressions when we have numerical values for the variables involved.

Evaluate Algebraic Expressions

When we are given an algebraic expression, one of the most common things we will have to do with it is **evaluate** it for some given value of the variable. The following example illustrates this process.

Example 2

Let $x = 12$. Find the value of $2x - 7$.

To find the solution, substitute 12 for x in the given expression. Every time we see x we will replace it with 12. **Note:** At this stage we place the value in parentheses:

$$\begin{aligned} 2x - 7 &= 2(12) - 7 \\ &= 24 - 7 \\ &= 17 \end{aligned}$$

The reason we place the substituted value in parentheses is twofold:

1. It will make worked examples easier for you to follow.
2. It avoids any confusion that would arise from dropping a multiplication sign: $2 \cdot 12 = 2(12) \neq 212$.

Example 3

Let $x = -1$. Find the value of $-9x + 2$.

Solution

$$\begin{aligned} -9(-1) + 2 &= 9 + 2 \\ &= 11 \end{aligned}$$


Example 4

Let $y = -2$. Find the value of $\frac{7}{y} - 11y + 2$.

Solution

$$\begin{aligned} \frac{7}{(-2)} - 11(-2) + 2 &= -3\frac{1}{2} + 22 + 2 \\ &= 24 - 3\frac{1}{2} \\ &= 20\frac{1}{2} \end{aligned}$$

Many expressions have more than one variable in them. For example, the formula for the perimeter of a rectangle in the introduction has two variables: length (l) and width (w). In these cases be careful to substitute the appropriate value in the appropriate place.

Example 5

The area of a trapezoid is given by the equation $A = \frac{h}{2}(a + b)$. Find the area of a trapezoid with bases $a = 10$ cm, $b = 15$ cm and height $h = 8$ cm.

To find the solution to this problem we simply take the values given for the variables, a , b and h , and *plug them in* to the expression for A :

$$\begin{aligned} A &= \frac{h}{2}(a + b) && \text{Substitute 10 for a, 15 for b and 8 for h.} \\ A &= \frac{8}{2}(10 + 15) && \text{Evaluate piece by piece. } (10 + 15) = 25; \frac{8}{2} = 4 \\ A &= 4(25) = 100 \end{aligned}$$

Solution: The area of the trapezoid is 100 square centimeters.

Example 6

Find the value of $\frac{1}{9}(5x + 3y + z)$ when $x = 7$, $y = -2$ and $z = 11$.

Let's plug in values for x , y and z and then evaluate the resulting expression.

$$\begin{aligned} \frac{1}{9}(5(7) + 3(-2) + (11)) &&& \text{Evaluate the individual terms inside the parentheses.} \\ \frac{1}{9}(35 + (-6) + 11) &&& \text{Combine terms inside parentheses.} \\ \frac{1}{9}(40) = \frac{40}{9} \approx 4.44 \end{aligned}$$

Solution ≈ 4.44 (rounded to the nearest hundredth) **Example 7**

The total resistance of two electronics components wired in parallel is given by

$$\frac{R_1 R_2}{R_1 + R_2}$$

where R_1 and R_2 are the individual resistances (in ohms) of the two components. Find the combined resistance of two such wired components if their individual resistances are 30 ohms and 15 ohms.

Solution

$$\frac{R_1 R_2}{R_1 + R_2} = \frac{(30)(15)}{30 + 15} = \frac{450}{45} = 10 \text{ ohms}$$

Substitute the values $R_1 = 30$ and $R_2 = 15$.

The combined resistance is 10 ohms.

Evaluate Algebraic Expressions with Exponents

Many formulas and equations in mathematics contain exponents. Exponents are used as a short-hand notation for repeated multiplication. For example:

$$2 \cdot 2 = 2^2$$

$$2 \cdot 2 \cdot 2 = 2^3$$

The exponent stands for how many times the number is used as a factor (multiplied). When we deal with integers, it is usually easiest to simplify the expression. We simplify:

$$2^2 = 4$$


and

$$2^3 = 8$$

However, we need exponents when we work with variables, because it is much easier to write x^8 than $x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot x$.

To evaluate expressions with exponents, substitute the values you are given for each variable and simplify. It is especially important in this case to substitute using parentheses in order to make sure that the simplification is done correctly.

Example 8



The area of a circle is given by the formula $A = \pi r^2$. Find the area of a circle with radius $r = 17$ inches.

Substitute values into the equation.

$$A = \pi r^2$$

Substitute 17 for r .

$$A = \pi(17)^2$$

$\pi \cdot 17 \cdot 17 = 907.9202 \dots$ Round to 2 decimal places.

The area is approximately 907.92 square inches.

Example 9

Find the value of $5x^2 - 4y$ for $x = -4$ and $y = 5$.

Substitute values in the following:

$$\begin{aligned} 5x^2 - 4y &= 5(-4)^2 - 4(5) \\ &= 5(16) - 4(5) \\ &= 80 - 20 \\ &= 60 \end{aligned}$$

Substitute $x = -4$ and $y = 5$.

Evaluate the exponent $(-4)^2 = 16$.

Example 10

Find the value of $2x^3 - 3x^2 + 5$, for $x = -5$.

Substitute the value of x in the expression:

$$\begin{aligned} 2x^3 - 3x^2 + 5 &= 2(-5)^3 - 3(-5)^2 + 5 \quad \text{Substitute } -5 \text{ for } x. \\ &= 2(-125) - 3(25) + 5 \quad \text{Evaluate exponents } (-5)^3 = (-5)(-5)(-5) = -125 \text{ and } (-5)^2 = (-5)(-5) = 25 \\ &= -250 - 75 + 5 \\ &= -320 \end{aligned}$$

Example 11

Find the value of $\frac{x^2y^3}{x^3+y^2}$, for $x = 2$ and $y = -4$.

Substitute the values of x and y in the following.

$$\begin{aligned} \frac{x^2y^3}{x^3+y^2} &= \frac{(2)^2(-4)^3}{(2)^3+(-4)^2} \\ \frac{4(-64)}{8+16} &= \frac{-256}{24} = \frac{-32}{3} \end{aligned}$$

Substitute 2 for x and -4 for y .

Evaluate expressions: $(2)^2 = (2)(2) = 4$ and $(2)^3 = (2)(2)(2) = 8$.

$(-4)^2 = (-4)(-4) = 16$ and $(-4)^3 = (-4)(-4)(-4) = -64$.

Example 12

The height (h) of a ball in flight is given by the formula: $h = -32t^2 + 60t + 20$, where the height is given in feet and the time (t) is given in seconds. Find the height of the ball at time $t = 2$ seconds.

Solution

$$\begin{aligned}
 h &= -32t^2 + 60t + 20 \\
 &= -32(2)^2 + 60(2) + 20 \\
 &= -32(4) + 60(2) + 20 \\
 &= 12 \text{ feet}
 \end{aligned}$$

Substitute 2 for t .

Review Questions

Write the following in a more condensed form by leaving out a multiplication symbol.

- $2 \times 11x$
- $1.35 \cdot y$
- $3 \times \frac{1}{4}$
- $\frac{1}{4} \cdot z$

Evaluate the following expressions for $a = -3$, $b = 2$, $c = 5$ and $d = -4$.

- $2a + 3b$
- $4c + d$
- $5ac - 2b$
- $\frac{2a}{c-d}$
- $\frac{3b}{d}$
- $\frac{a-4b}{3c+2d}$
- $\frac{1}{a+b}$
- $\frac{ab}{cd}$

Evaluate the following expressions for $x = -1$, $y = 2$, $z = -3$, and $w = 4$.

- $8x^3$
- $\frac{5x^2}{6z^3}$
- $3z^2 - 5w^2$
- $x^2 - y^2$
- $\frac{z^3 + w^3}{z^3 - w^3}$
- $2x^2 - 3x^2 + 5x - 4$
- $4w^3 + 3w^2 - w + 2$
- $3 + \frac{1}{z^2}$
- The weekly cost C of manufacturing x remote controls is given by the formula $C = 2000 + 3x$, where the cost is given in dollars.
 - What is the cost of producing 1000 remote controls?
 - What is the cost of producing 2000 remote controls?
- The volume of a box without a lid is given by the formula: $V = 4x(10 - x)^2$ where x is a length in inches and V is the volume in cubic inches.
 - What is the volume when $x = 2$?
 - What is the volume when $x = 3$?

Review Answers

- $22x$

2. $1.35y$
3. $\frac{3}{4}$
4. $\frac{x}{4}$
5. 0
6. 16
7. -79
8. $\frac{-2}{3}$
9. $\frac{-3}{2}$
10. $\frac{-11}{7}$
11. -1
12. $\frac{3}{10}$
13. -8
14. $\frac{-5}{162}$
15. -53
16. -3
17. $\frac{37}{-91}$
18. -14
19. 302
20. $3\frac{1}{9}$
21. (a) \$5000;
(b) \$8000
22. (a) 512 in^3 ;
(b) 588 in^3

2.2 Order of Operations

Learning Objectives

- Evaluate algebraic expressions with grouping symbols.
- Evaluate algebraic expressions with fraction bars.
- Evaluate algebraic expressions with a graphing calculator.

Introduction

Look at and evaluate the following expression:

$$2 + 4 \times 7 - 1 = ?$$

How many different ways can we interpret this problem, and how many different answers could someone possibly find for it?

The *simplest* way to evaluate the expression is simply to start at the left and work your way across, keeping track of the total as you go:

$$2 + 4 = 6$$

$$6 \times 7 = 42$$

$$42 - 1 = 41$$

If you enter the expression into a *non-scientific*, non-graphing calculator you will probably get 41 as the answer. If, on the other hand, you were to enter the expression into a scientific calculator or a graphing calculator you would probably get 29 as an answer.

In mathematics, the order in which we perform the various **operations** (such as adding, multiplying, etc.) is important. In the expression above, the operation of **multiplication** takes precedence over **addition** so we evaluate it first. Lets re-write the expression, but put the multiplication in brackets to indicate that it is to be evaluated first.

$$2 + (4 \times 7) - 1 = ?$$

So we first evaluate the brackets: $4 \times 7 = 28$. Our expression becomes:

$$2 + (28) - 1 = ?$$

When we have only addition and subtraction, we start at the left and keep track of the total as we go:

$$2 + 28 = 30$$

$$30 - 1 = 29$$

Algebra students often use the word **PEMDAS** to help remember the order in which we evaluate the mathematical expressions: **P**arentheses, **E**xponents, **M**ultiplication, **D**ivision, **A**ddition and **S**ubtraction.

Order of Operations

1. Evaluate expressions within **P**arentheses (also all brackets [] and braces { }) first.
2. Evaluate all **E**xponents (squared or cubed terms such as 3^2 or x^3) next.
3. **M**ultiplication *and* **D**ivision is next work from left to right completing **both** multiplication and division in the order that they appear.
4. Finally, evaluate **A**ddition *and* **S**ubtraction work from left to right completing **both** addition and subtraction in the order that they appear.

Evaluate Algebraic Expressions with Grouping Symbols

The first step in the order of operations is called **parentheses**, but we include all **grouping symbols** in this step. While we will mostly use parentheses () in this book, you may also see square brackets [] and curly braces { } and you should include them as part of the first step.

Example 1

Evaluate the following:

a) $4 - 7 - 11 - 2$

b) $4 - (7 - 11) + 2$

c) $4 - [7 - (11 + 2)]$

Each of these expressions has the same numbers and the same mathematical operations, in the same order. The placement of the various grouping symbols means, however, that we must evaluate everything in a different order each time. Let's look at how we evaluate each of these examples.

a) This expression doesn't have parentheses. PEMDAS states that we treat addition and subtraction as they appear, starting at the left and working right (its NOT addition *then* subtraction).

Solution

$$\begin{aligned} 4 - 7 - 11 + 2 &= -3 - 11 + 2 \\ &= -14 + 2 \\ &= -12 \end{aligned}$$

b) This expression has parentheses. We first evaluate $7 - 11 = -4$. Remember that when we subtract a negative it is equivalent to adding a positive:

Solution

$$\begin{aligned} 4 - (7 - 11) + 2 &= 4 - (-4) + 2 \\ &= 8 + 2 \\ &= 10 \end{aligned}$$

c) Brackets are often used to group expressions which already contain parentheses. This expression has both brackets and parentheses. Do the innermost group first, $(11 + 2) = 13$. Then complete the operation in the brackets.

Solution

$$\begin{aligned} 4 - [7 - (11 + 2)] &= 4 - [7 - (13)] \\ &= 4 - [-6] \\ &= 10 \end{aligned}$$

Example 2

Evaluate the following:

a) $3 \times 5 - 7 \div 2$

b) $3 \times (5 - 7) \div 2$

c) $(3 \times 5) - (7 \div 2)$

a) There are no grouping symbols. PEMDAS dictates that we evaluate multiplication and division first, working from left to right: $3 \times 5 = 15$; $7 \div 2 = 3.5$. (NOTE: Its not multiplication *then* addition) Next we perform the subtraction:

Solution

$$\begin{aligned} 3 \times 5 - 7 \div 2 &= 15 - 3.5 \\ &= 11.5 \end{aligned}$$

b) First, we evaluate the expression inside the parentheses: $5 - 7 = -2$. Then work from left to right.

Solution

$$\begin{aligned} 3 \times (5 - 7) \div 2 &= 3 \times (-2) \div 2 \\ &= (-6) \div 2 \\ &= -3 \end{aligned}$$

c) First, we evaluate the expressions inside parentheses: $3 \times 5 = 15$, $7 \div 2 = 3.5$. Then work from left to right.

Solution

$$\begin{aligned} (3 \times 5) - (7 \div 2) &= 15 - 3.5 \\ &= 11.5 \end{aligned}$$

Note that in part (c), the result was unchanged by adding parentheses, but the expression does appear easier to read. Parentheses can be used in two distinct ways:

- To alter the order of operations in a given expression
- To clarify the expression to make it easier to understand

Some expressions contain no parentheses, others contain many sets. Sometimes expressions will have sets of parentheses **inside** other sets of parentheses. When faced with **nested parentheses**, start at the innermost parentheses and work outward.

Example 3

Use the order of operations to evaluate:

$$8 - [19 - (2 + 5) - 7]$$

Follow PEMDAS first parentheses, starting with innermost brackets first:

Solution

$$\begin{aligned} 8 - (19 - (2 + 5) - 7) &= 8 - (19 - 7 - 7) \\ &= 8 - 5 \\ &= 3 \end{aligned}$$

In algebra, we use the order of operations when we are substituting values into expressions for variables. In those situations we will be given an expression involving a variable or variables, and also the values to substitute for any variables in that expression.

Example 4

Use the order of operations to evaluate the following:

a) $2 - (3x + 2)$ when $x = 2$

b) $3y^2 + 2y - 1$ when $y = -3$

c) $2 - (t - 7)^2 \times (u^3 - v)$ when $t = 19, u = 4$ and $v = 2$

a) The first step is to substitute in the value for x into the expression. Let's put it in parentheses to clarify the resulting expression.

Solution

$$2 - (3(2) + 2)$$

$$3(2) \text{ is the same as } 3 \times 2$$

Follow PEMDAS first parentheses. Inside parentheses follow PEMDAS again.

$$\begin{aligned} 2 - (3 \times 2 + 2) &= 2 - (6 + 2) \\ 2 - 8 &= -6 \end{aligned}$$

Inside the parentheses, we evaluate the multiplication first.

Now we evaluate the parentheses.

b) The first step is to substitute in the value for y into the expression.

Solution

$$3 \times (-3)^2 + 2 \times (-3) - 1$$

Follow PEMDAS: we cannot simplify parentheses.

$$\begin{aligned} &= 3 \times (-3)^2 + 2 \times (-3) - 1 \\ &= 3 \times 9 + 2 \times (-3) - 1 \\ &= 27 + (-6) - 1 \\ &= 27 - 6 - 1 \\ &= 20 \end{aligned}$$

Evaluate exponents : $(-3)^2 = 9$

Evaluate multiplication : $3 \times 9 = 27; 2 \times -3 = -6$

Evaluate addition and subtraction in order from left to right.

c) The first step is to substitute the values for t , u , and v into the expression.

Solution:

$$2 - (19 - 7)^2 \times (4^3 - 2)$$

Follow **PEMDAS**:

$= 2 - (19 - 7)^2 \times (4^3 - 2)$ $= 2 - 12^2 \times 62$ $= 2 - 144 \times 62$ $= 2 - 8928$ $= -8926$	<p>Evaluate parentheses : $(19 - 7) = 12$; $(4^3 - 2) = (64 - 2) = 62$</p> <p>Evaluate exponents : $12^2 = 144$</p> <p>Evaluate the multiplication : $144 \times 62 = 8928$</p> <p>Evaluate the subtraction.</p>
---	--

In parts (b) and (c) we left the parentheses around the negative numbers to clarify the problem. They did not affect the order of operations, but they did help avoid confusion when we were multiplying negative numbers.

Part (c) in the last example shows another interesting point. When we have an expression inside the parentheses, we use PEMDAS to determine the order in which we evaluate the contents.

Evaluating Algebraic Expressions with Fraction Bars

Fraction bars count as grouping symbols for PEMDAS, and should therefore be evaluated in the first step of solving an expression. All numerators and all denominators can be treated as if they have invisible parentheses. When **real** parentheses are also present, remember that the innermost grouping symbols should be evaluated first. If, for example, parentheses appear on a numerator, they would take precedence over the fraction bar. If the parentheses appear outside of the fraction, then the fraction bar takes precedence.

Example 5

Use the order of operations to evaluate the following expressions:

a) $\frac{z+3}{4} - 1$ When $z = 2$

b) $\left(\frac{a+2}{b+4} - 1\right) + b$ When $a = 3$ and $b = 1$

c) $2 \times \left(\frac{w+(x-2z)}{(y+2)^2} - 1\right)$ When $w = 11$, $x = 3$, $y = 1$ and $z = -2$

a) We substitute the value for z into the expression.

Solution:

$$\frac{2+3}{4} - 1$$

Although this expression has no parentheses, we will rewrite it to show the effect of the fraction bar.

$$\frac{(2+3)}{4} - 1$$

Using PEMDAS, we first evaluate the expression on the numerator.

$$\frac{5}{4} - 1$$

We can convert $\frac{5}{4}$ to a mixed number:

$$\frac{5}{4} = 1\frac{1}{4}$$

Then evaluate the expression:

$$\frac{5}{4} - 1 = 1\frac{1}{4} - 1 = \frac{1}{4}$$

b) We substitute the values for a and b into the expression:

Solution:

$$\left(\frac{3+2}{1+4} - 1\right) - 1$$

This expression has nested parentheses (remember the effect of the fraction bar on the numerator and denominator). The innermost grouping symbol is provided by the fraction bar. We evaluate the numerator ($3 + 2$) and denominator ($1 + 4$) first.

$$\begin{aligned} &\left(\frac{5}{5} - 1\right) - 1 \\ &(1 - 1) - 1 \\ &0 - 1 = -1 \end{aligned}$$

Now we evaluate the inside of the parentheses, starting with division.

Next the subtraction.

c) We substitute the values for w , x , y and z into the expression:

Solution:

This complicated expression has several layers of nested parentheses. One method for ensuring that we start with the innermost parentheses is to make use of the other types of brackets. We can rewrite this expression, putting brackets in for the fraction bar. The outermost brackets we will leave as parentheses (). Next will be the *invisible brackets* from the fraction bar, these will be written as []. The third level of nested parentheses will be the . We will leave negative numbers in round brackets.

$$2\left(\frac{[11 + \{3 - 2(-2)\}]}{[\{1 + 2\}^2]} - 1\right)$$

We start with the innermost grouping sign { } .

$$\{1 + 2\} = 3; \{3 - 2(-2)\} = 3 + 4 = 7$$

$$2\left(\frac{[11 + 7]}{[3^2]} - 1\right)$$

The next level has two square brackets to evaluate.

$$2\left(\frac{18}{9} - 1\right)$$

We now evaluate the round brackets, starting with division.

$$2(2 - 1)$$

Finally, we complete the addition and subtraction.

$$2(1) = 2$$

Evaluate Algebraic Expressions with a TI-83/84 Family Graphing Calculator

A graphing calculator is a very useful tool in evaluating algebraic expressions. The graphing calculator follows PEMDAS. In this section we will explain two ways of evaluating expressions with the graphing calculator.

Method 1: Substitute for the variable first. Then evaluate the numerical expression with the calculator.

Example 6

Evaluate $[3(x^2 - 1)^2 - x^4 + 12] + 5x^3 - 1$ when $x = -3$

Solution:

Substitute the value $x = -3$ into the expression.

$$[3((-3)^2 - 1)^2 - (-3)^4 + 12] + 5(-3)^3 - 1$$

Input this in the calculator just as it is and press **[ENTER]**. (Note, use ^ for exponents)

The answer is -13 .

Method 2: Input the original expression in the calculator first and then evaluate. Lets look at the same example.

Evaluate $[3(x^2 - 1)^2 - x^4 + 12] + 5x^3 - 1$ when $x = -3$

First, store the value $x = -3$ in the calculator. Type -3 **[STO]** x (The letter x can be entered using the x -**[VAR]** button or **[ALPHA]** + **[STO]**). Then type in the expression in the calculator and press **[ENTER]**.

The answer is -13 .

The second method is better because you can easily evaluate the same expression for any value you want. For example, lets evaluate the same expression using the values $x = 2$ and $x = \frac{2}{3}$.

For $x = 2$, store the value of x in the calculator: 2 **[STO]** x . Press **[2nd]** **[ENTER]** twice to get the previous expression you typed in on the screen without having to enter it again. Press **[ENTER]** to evaluate.

The answer is 62 .

For $x = \frac{2}{3}$, store the value of x in the calculator: $\frac{2}{3}$ **[STO]** x . Press **[2nd]** **[ENTER]** twice to get the expression on the screen without having to enter it again. Press **[ENTER]** to evaluate.

The answer is 13.21 or $\frac{1070}{81}$ in fraction form.


Note: On graphing calculators there is a difference between the minus sign and the negative sign. When we stored the value negative three, we needed to use the negative sign which is to the left of the [ENTER] button on the calculator. On the other hand, to perform the subtraction operation in the expression we used the minus sign. The minus sign is right above the plus sign on the right.

You can also use a graphing calculator to evaluate expressions with more than one variable.

Example 7

Evaluate the expression: $\frac{3x^2 - 4y^2 + x^4}{(x+y)^{1/2}}$ for $x = -2$, $y = 1$.

Solution



Store the values of x and y . -2 [STO] x , 1 [STO] y . The letters x and y can be entered using [ALPHA] + [KEY]. Input the expression in the calculator. When an expression shows the division of two expressions be sure to use parentheses: (numerator) \div (denominator)

Press [ENTER] to obtain the answer $-.8\bar{8}$ or $-\frac{8}{9}$.

Review Questions

1. Use the order of operations to evaluate the following expressions.

- $8 - (19 - (2 + 5) - 7)$
- $2 + 7 \times 11 - 12 \div 3$
- $(3 + 7) \div (7 - 12)$
- $\frac{2 \cdot (3 + (2 - 1))}{4 - (6 + 2)} - (3 - 5)$

2. Evaluate the following expressions involving variables.

- $\frac{jk}{j+k}$ when $j = 6$ and $k = 12$.
- $2y^2$ when $x = 1$ and $y = 5$
- $3x^2 + 2x + 1$ when $x = 5$
- $(y^2 - x)^2$ when $x = 2$ and $y = 1$

3. Evaluate the following expressions involving variables.

- $\frac{4x}{9x^2 - 3x + 1}$ when $x = 2$
- $\frac{z^2}{x+y} + \frac{x^2}{x-y}$ when $x = 1$, $y = -2$, and $z = 4$.
- $\frac{4xyz}{y^2 - x^2}$ when $x = 3$, $y = 2$, and $z = 5$
- $\frac{x^2 - z^2}{xz - 2x(z-x)}$ when $x = -1$ and $z = 3$

4. Insert parentheses in each expression to make a true equation.

- $5 - 2 \cdot 6 - 4 + 2 = 5$
- $12 \div 4 + 10 - 3 \cdot 3 + 7 = 11$
- $22 - 32 - 5 \cdot 3 - 6 = 30$
- $12 - 8 - 4 \cdot 5 = -8$

5. Evaluate each expression using a graphing calculator.

- $x^2 + 2x - xy$ when $x = 250$ and $y = -120$
- $(xy - y^4)^2$ when $x = 0.02$ and $y = -0.025$

- (c) $\frac{x+y-z}{xy+yz+xz}$ when $x = \frac{1}{2}$, $y = \frac{3}{2}$, and $z = -1$
- (d) $\frac{(x+y)^2}{4x^2-y^2}$ when $x = 3$ and $y = -5d$

Review Answers

- 3
 - 75
 - 2
 - 2
- 4
 - 300
 - 86
 - 3
- $\frac{8}{31}$
 - $-\frac{47}{3}$
 - 24
 - $-\frac{8}{5}$
- $(5-2) \cdot (6-5) + 2 = 5$
 - $(12 \div 4) + 10 - (3 \cdot 3) + 7 = 11$
 - $(22 - 32 - 5) \cdot (3 - 6) = 30$
 - $12 - (8 - 4) \cdot 5 = -8$
- 93000
 - 0.00000025
 - $-\frac{12}{5}$
 - $\frac{4}{11}$

2.3 One-Step Equations

Learning Objectives

- Solve an equation using addition.
- Solve an equation using subtraction.
- Solve an equation using multiplication.
- Solve an equation using division.

Introduction

Nadia is buying a new mp3 player. Peter watches her pay for the player with a \$100 bill. She receives \$22.00 in change, and from only this information, Peter works out how much the player cost. How much was the player?

In math, we can solve problems like this using an **equation**. An **equation** is an algebraic expression that involves an **equals** sign. If we use the letter x to represent the cost of the mp3 player we could write the following equation.

$$x + 22 = 100$$

This tells us that the value of the player **plus** the value of the change received is **equal** to the \$100 that Nadia paid.

Peter saw the transaction from a different viewpoint. He saw Nadia receive the player, give the salesperson \$100 then he saw Nadia receive \$22 change. Another way we could write the equation would be:

$$x = 100 - 22$$

This tells us that the value of the player is **equal** to the amount of money Nadia paid ($100 - 22$).

Mathematically, these two equations are equivalent. Though it is easier to determine the cost of the mp3 player from the second equation. In this chapter, we will learn how to solve for the variable in a one variable linear equation. Linear equations are equations in which each term is either a constant or the product of a constant and a single variable (to the first power). The term linear comes from the word line. You will see in later chapters that linear equations define lines when graphed.

We will start with simple problems such as the one in the last example.

Solve an Equation Using Addition

When we write an algebraic equation, equality means that whatever we do to one side of the equation, we have to do to the other side. For example, to get from the second equation in the introduction back to the first equation, we would add a quantity of 22 to both sides:

$$x = 100 - 22$$

$$x + 22 = 100 - 22 + 22 \qquad \text{or} \qquad x + 22 = 100$$

Similarly, we can add numbers to each side of an equation to help solve for our unknown.

Example 1

Solve $x - 3 = 9$

Solution

We need to **isolate** x . Change our equation so that x appears by itself on one side of the equals sign. Right now our x has a 3 subtracted from it. To reverse this, we could add 3, but we must do this to **both sides**.

$$\begin{aligned} x - 3 &= 9 \\ x - 3 + 3 &= 9 + 3 \quad \text{The } +3 \text{ and } -3 \text{ on the left cancel each other. We evaluate } 9 + 3 \\ x &= 12 \end{aligned}$$

Example 2

Solve $x - 3 = 11$

Solution

To isolate x we need to add 3 to both sides of the equation. This time we will add vertically.

$$\begin{aligned} x - 3 &= 11 \\ +3 &= +3 \\ x &= 14 \end{aligned}$$

Notice how this format works. One term will always cancel (in this case the three), but we need to remember to carry the x down and evaluate the sum on the other side of the equals sign.

Example 3

Solve $z - 9.7 = -1.026$

Solution

This time our variable is z , but don't let that worry you. Treat this variable like any other variable.

$$\begin{aligned} z - 9.7 &= -1.026 \\ +9.7 &= +9.7 \\ z &= 8.674 \end{aligned}$$

Make sure that you understand the addition of decimals in this example!

Solve an Equation Using Subtraction

When our variable appears with a number added to it, we follow the same process, only this time to isolate the variable we **subtract** a number from both sides of the equation.

Example 4Solve $x + 6 = 26$ **Solution**To isolate x we need to subtract six from both sides.

$$\begin{aligned}x + 6 &= 26 \\-6 &= -6 \\x &= 20\end{aligned}$$

Example 5Solve $x + 20 = -11$ **Solution**To isolate x we need to subtract 20 from both sides of the equation.

$$\begin{aligned}x + 20 &= -11 \\-20 &= -20 \\x &= -31\end{aligned}$$

Example 6Solve $x + \frac{4}{7} = \frac{9}{5}$ **Solution**To isolate x we need to subtract $\frac{4}{7}$ from both sides.

$$\begin{aligned}x + \frac{4}{7} &= \frac{9}{5} \\-\frac{4}{7} &= -\frac{4}{7} \\x &= \frac{9}{5} - \frac{4}{7}\end{aligned}$$

To solve for x , make sure you know how to subtract fractions. We need to find the lowest common denominator. 5 and 7 are both prime. So we can multiply to find the LCD, $\text{LCD} = 5 \cdot 7 = 35$.

$$\begin{aligned}x &= \frac{9}{5} - \frac{4}{7} \\x &= \frac{7 \cdot 9}{35} - \frac{4 \cdot 5}{35} \\x &= \frac{63 - 20}{35} \\x &= \frac{43}{35}\end{aligned}$$

Make sure you are comfortable with decimals and fractions! To master algebra, you will need to work with them frequently.

Solve an Equation Using Multiplication



Suppose you are selling pizza for \$1.50 a slice and you get eight slices out of a single pizza. How much do you get for a single pizza? It shouldn't take you long to figure out that you get $8 \times \$1.50 = \12.00 . You solve this problem by multiplying. The following examples do the same algebraically, using the unknown variable x as the cost in dollars of the whole pizza.

Example 7

Solve $\frac{1}{8} \cdot x = 1.5$

Our x is being multiplied by one-eighth. We need to cancel this factor, so we multiply by the reciprocal 8. Do not forget to multiply **both sides** of the equation.

$$\begin{aligned} 8 \left(\frac{1}{8} \cdot x \right) &= 8(1.5) \\ x &= 12 \end{aligned}$$

In general, when x is multiplied by a fraction, we multiply by the reciprocal of that fraction.

Example 8

Solve $\frac{9x}{5} = 5$

$\frac{9x}{5}$ is equivalent to $\frac{9}{5} \cdot x$ so x is being multiplied by $\frac{9}{5}$. To cancel, multiply by the reciprocal, $\frac{5}{9}$.

$$\begin{aligned} \frac{5}{9} \left(\frac{9x}{5} \right) &= \frac{5}{9} \cdot 5 \\ x &= \frac{25}{9} \end{aligned}$$

Example 9

Solve $0.25x = 5.25$

0.25 is the decimal equivalent of one fourth, so to cancel the 0.25 factor we would multiply by 4.

$$\begin{aligned} 4(0.25x) &= 4(5.25) \\ x &= 21 \end{aligned}$$

Solve an Equation Using Division

Solving by division is another way to cancel any terms that are being multiplied x . Suppose you buy five identical candy bars, and you are charged \$3.25. How much did each candy bar cost? You might just divide \$3.25 by 5. Or you may convert to cents and divide 325 by 5. Let's see how this problem looks in algebra.

Example 10

Solve $5x = 3.25$ To cancel the 5 we divide both sides by 5.

$$\begin{aligned}\cancel{5}x &= \frac{3.25}{\cancel{5}} \\ x &= 0.65\end{aligned}$$

Example 11

Solve $7x = \frac{5}{11}$ Divide both sides by 7.

$$\begin{aligned}x &= \frac{5}{7 \cdot 11} \\ x &= \frac{5}{77}\end{aligned}$$

Example 12

Solve $1.375x = 1.2$ Divide by 1.375

$$\begin{aligned}x &= \frac{1.2}{1.375} \\ x &= 0.8\overline{72}\end{aligned}$$

Notice the bar above the final two decimals. It means recurring or repeating: the full answer is $0.872727272\dots$

Solve Real-World Problems Using Equations**Example 13**

In the year 2017, Anne will be 45 years old. In what year was Anne born?

The unknown here is the year Anne was born. This is x . Here is our equation.


$$\begin{aligned}x + 45 &= 2017 \\ -45 &= -45 \\ x &= 1972\end{aligned}$$

Solution

Anne was born in 1972.

Example 14

A mail order electronics company stocks a new mini DVD player and is using a balance to determine the shipping weight. Using only one lb weights, the shipping department found that the following arrangement balances.



Knowing that each weight is one lb, calculate the weight of one DVD player.

Solution

We know that the system balances, so the weights on each side must be equal. We can write an algebraic expression based on this equality. The unknown quantity, the weight of the DVD player (in pounds), will be x . The combined weight on the right of the balance is $5 \times 1 \text{ lb} = 5\text{lb}$.

$2x = 5$ Divide both sides by 2.

$$\frac{2x}{2} = \frac{5}{2}$$

$x = 2.5$

Each DVD player weighs x 2.5 lbs.

Example 15



In good weather, tomato seeds can grow into plants and bear ripe fruit in as little as 19 weeks . Lora planted her seeds 11weeks ago. How long must she wait before her tomatoes are ready to eat?

Solution

We know that the total time to bear fruit is 19 weeks , and that the time so far is 19 weeks . Our unknown is the time in weeks remaining, so we call that x . Here is our equation.

$$\begin{aligned} x + 11 &= 19 \\ -11 &= -11 \\ x &= 8 \end{aligned}$$

Lora will have to wait another 8 weeks before her tomatoes are ready. We can show this by designing a table.

Tomato Readiness by Week											Time Now										
Week number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
Tomatoes ready to eat?	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	Y

11 weeks passed
8 weeks remain



Example 16

In 2004, Takeru Kobayashi, of Nagano, Japan, ate $53\frac{1}{2}$ hot dogs in 12 minutes . He broke his previous world record, set in 2002, by three hot dogs. Calculate:

- a) How many minutes it took him to eat one hot dog.
- b) How many hot dogs he ate per minute.
- c) What his old record was.

a) We know that the total time for 53.5 hot dogs is 12 minutes . If the time, in minutes, for each hot dog (the unknown) is x then we can write the following equation.

$$53.5x = 12$$

$$x = \frac{12}{53.5} = 0.224$$

Divide both sides by 53.5

minutes Convert to seconds, by multiplying by 60

Solution

The time taken to eat one hot dog is 0.224 minutes , or about 13.5 seconds .

Note: We round off our answer as there is no need to give our answer to an accuracy better than 0.1 (one tenth) of a second.

b) This time, we look at our data slightly differently. We know that he ate for 12 minutes . His **rate per-minute** is our new unknown (to avoid confusion with x , we will call this y). We know that the total number of hot dogs is 53.5 so we can write the following equation.

$$12y = 53.5$$

$$y = \frac{53.5}{12} = 4.458$$

Divide both sides by 12

Solution

Takeru Kobayashi ate approximately 4.5. hot dogs per minute.

c) We know that his new record is 53.5. and also that his new record is three more than his old record. We have a new unknown. We will call his old record z , and write the following equation.

$$x + 3 = 53.5$$

$$-3 = -3$$

$$x = 50.5$$

Solution

Takeru Kobayashis old record was $50\frac{1}{2}$ hot dogs in 12 minutes .

Lesson Summary

- An equation in which each term is either a constant or a product of a constant and a single variable is a **linear equation**.
- Adding, subtracting, multiplying, or dividing both sides of an equation by the same value results in an equivalent equation.
- To solve an equation, **isolate** the unknown variable on one side of the equation by applying one or more arithmetic operations to both sides.

Review Questions

1. Solve the following equations for x .

(a) $x + 11 = 7$

(b) $x - 1.1 = 3.2$

- (c) $7x = 21$
- (d) $4x = 1$
- (e) $\frac{5x}{12} = \frac{2}{3}$
- (f) $x + \frac{5}{2} = \frac{2}{3}$
- (g) $x - \frac{5}{6} = \frac{3}{8}$
- (h) $0.01x = 11$

2. Solve the following equations for the unknown variable.

- (a) $q - 13 = -13$
- (b) $z + 1.1 = 3.0001$
- (c) $21s = 3$
- (d) $t + \frac{1}{2} = \frac{1}{3}$
- (e) $\frac{7f}{11} = \frac{7}{11}$
- (f) $\frac{3}{4} = -\frac{1}{2} \cdot y$
- (g) $6r = \frac{3}{8}$
- (h) $\frac{9b}{16} = \frac{3}{8}$

3. Peter is collecting tokens on breakfast cereal packets in order to get a model boat. In eight weeks he has collected 10 tokens. He needs 25 tokens for the boat. Write an equation and determine the following information.

- (a) How many more tokens he needs to collect, n .
- (b) How many tokens he collects per week, w .
- (c) How many more weeks remain until he can send off for his boat, r .

4. Juan has baked a cake and wants to sell it in his bakery. He is going to cut it into 12 slices and sell them individually. He wants to sell it for three times the cost of making it. The ingredients cost him \$8.50, and he allowed \$1.25 to cover the cost of electricity to bake it. Write equations that describe the following statements

- (a) The amount of money that he sells the cake for (u).
- (b) The amount of money he charges for each slice (c).
- (c) The total profit he makes on the cake (w).

Review Answers

1.
 - (a) $x = -4$
 - (b) $x = 4.3$
 - (c) $x = 3$
 - (d) $x = 0.25$
 - (e) $x = 1.6$
 - (f) $x = -\frac{11}{6}$
 - (g) $x = \frac{29}{24}$
 - (h) $x = 1100$
2.
 - (a) $q = 0$
 - (b) $z = 1.9001$
 - (c) $s = 1/7$
 - (d) $t = -\frac{1}{6}$
 - (e) $f = 1$
 - (f) $y = -1.5$
 - (g) $r = \frac{1}{16}$
 - (h) $b = \frac{2}{3}$
3.
 - (a) $n + 10 = 25, n = 15$
 - (b) $8w = 10, w = 1.25$

- (c) $r \cdot w = 15$ or $1.25r = 15$, $r = 12$
4. (a) $u = 3(8.5 + 1.25)$
(b) $12v = u$
(c) $w = u - (8.5 + 1.25)$

2.4 Two-Step Equations

Learning Objectives


- Solve a two-step equation using addition, subtraction, multiplication, and division.
- Solve a two-step equation by combining like terms.
- Solve real-world problems using two-step equations.

Solve a Two-Step Equation

We have seen that in order to solve for an unknown variable we can isolate it on one side of the equal sign and evaluate the numbers on the other side. In this chapter we will expand our ability to do that, with problems that require us to combine more than one technique in order to solve for our unknown.

Example 1

Rebecca has three bags containing the same number of marbles, plus two marbles left over. She places them on one side of a balance. Chris, who has more marbles than Rebecca, added marbles to the other side of the balance. He found that with 29 marbles, the scales balanced. How many marbles are in each bag? Assume the bags weigh nothing.



Solution

We know that the system balances, so the weights on each side must be equal. We can write an algebraic expression based on this equality. The unknown quantity, the number of marbles in each bag, will be our x . We can see that on the left hand scale we have three bags (each containing x marbles) and two extra marbles. On the right scale we have 29 marbles. The balancing of the scales is similar to the balancing of the following equation.

$$3x + 2 = 29$$

Three bags plus two marbles **equals** 29 marbles

To solve for x we need to first get all the variables (terms containing an x) alone on one side. Look at the balance. There are no bags on the right. Similarly, there are no x terms on the right of our equation. We will aim to get all the constants on the right, leaving only the x on the left.

$$3x + 2 = 29$$

$$\cancel{-2} = \cancel{-2}$$

Subtract 2 from both sides :

$$3x = 27$$

$$\frac{\cancel{3}x}{\cancel{3}} = \frac{27}{3}$$


Divide both sides by 3

$$x = 9$$

Solution


There are nine marbles in each bag.

We can do the same with the real objects as we have done with the equation. Our first action was to subtract two from both sides of the equals sign. On the balance, we could remove this number of marbles from each scale. Because we remove the same number of marbles from each side, we know the scales will still balance.



Next, we look at the left hand scale. There are three bags of marbles. To make our job easier, we divide the marbles on the right scale into three equal piles. You can see that there are nine marbles in each.

Three bags of marbles **balances** three piles of nine marbles



So each bag of marble balances nine marbles. Again you see we reach our solution:

Solution

Each bag contains nine marbles.

On the web: <http://www.mste.uiuc.edu/pavel/java/balance/> has interactive balance beam activities!

Example 2

Solve $6(x+4) = 12$

Solution

This equation has the x buried in parentheses. In order to extract it we can proceed in one of two ways: we can either distribute the six on the left, or divide both sides by six to remove it from the left. Since the right hand side of the equation is a multiple of six, it makes sense to divide.

$$6(x+4) = 12$$

Divide both sides by 6.

$$\frac{6(x+4)}{6} = \frac{12}{6}$$

$$x+4 = 2$$

Subtract 4 from both sides.

$$-4 - 4$$

$$\hline x = -2$$

Solution

$$x = -2$$

Example 3

Solve $\frac{x-3}{5} = 7$

This equation has a fraction in it. It is always a good idea to get rid of fractions first.

$$\left(x - \frac{3}{5}\right) = 7$$

Solution:

$$\cancel{5} \left(\frac{x-3}{\cancel{5}} \right) = 5 \cdot 7$$

$$x - 3 = 35$$

$$+ 3 = +3$$

$$x = 38$$

Multiply both sides by 5

Add 3 to both sides

Solution

$$x = 38$$

Example 4

Solve $\frac{5}{9}(x+1) = \frac{2}{7}$

First, we will cancel the fraction on the left (making the coefficient equal to one) by multiplying by the reciprocal (the multiplicative inverse).

$$\cancel{\frac{9}{5}} \cdot \frac{5}{\cancel{9}}(x+1) = \frac{9}{5} \cdot \frac{2}{7}$$

$$x+1 = \frac{18}{35}$$

$$x = \frac{18}{35} - \frac{35}{35}$$

$$x = \frac{18-35}{35}$$

Subtract $1 \left(1 = \frac{35}{35}\right)$ from both sides.

Solution

$$x = -\frac{17}{35}$$

These examples are called **two-step equations**, as we need to perform two separate operations to the equation to isolate the variable.

Solve a Two-Step Equation by Combining Like Terms

When we look at linear equations we predominantly see two terms, those that contain the unknown variable as a factor, and those that don't. When we look at an equation that has an x on both sides, we know that in order to solve,

we need to get all the x -terms on one side of the equation. This is called **combining like terms**. They are **like terms** because they contain the same variable (or, as you will see in later chapters, the same combination of variables).

Like Terms

- $17x, 12x, -1.2x$, and $\frac{17x}{9}$
- $3y, 19y$, and $\frac{y}{99}$
- $xy, 6xy$, and $0.0001xy$

Unlike Terms

- $3x$ and $2y$
- $12xy$ and $2x$
- $0.001x$ and 0.001

To add or subtract like terms, we can use the Distributive Property of Multiplication instead of addition and subtraction.

$$\begin{aligned} 3x + 4x &= (3 + 4)x = 7x \\ 0.03xy - 0.01xy &= (0.03 - 0.01)xy = 0.02xy \\ -y + 16y - 5y &= (-1 + 16 - 5)y = 10y \\ 5z + 2z - 7z &= (5 + 2 - 7)z = 0z = 0 \end{aligned}$$

To solve an equation with two or more like terms we need to combine them before we can solve for the variable.

Example 5

Solve $(x + 5) - (2x - 3) = 6$

There are two like terms. The x and the $-2x$ (do not forget that the negative sign multiplies everything in the parentheses).

Collecting like terms means we write all the terms with matching variables together. We will then add, or subtract them individually. We pull out the x from the first bracket and the $-2x$ from the second. We then rewrite the equation collecting the like terms.

$$(x - 2x) + (5 - (-3)) = 6$$

Combine like terms and constants.

$$-x + 8 = 6$$

$$\cancel{-8} = -8$$

Subtract 8 from both sides

$$-x = -2$$

Multiply both sides by -1 to get the variable by itself

Solution

$$x = 2$$

Example 6

Solve $\frac{x}{2} - \frac{x}{3} = 6$

Solution

This problem involves fractions. Combining the variable terms will require dealing with fractions. We need to write all the terms on the left over a common denominator of six.

$$\begin{aligned}\frac{3x}{6} - \frac{2x}{6} &= 6 \\ \frac{x}{6} &= 6 \\ x &= 36\end{aligned}$$

Next we combine the fractions.

Multiply both sides by 6.

Solve Real-World Problems Using Two-Step Equations

When we are faced with real world problems the thing that gives people the most difficulty is going from a problem in words to an equation. First, look to see what the equation is asking. What is the **unknown** for which you have to solve? That will determine the quantity we will use for our **variable**. The text explains what is happening. Break it down into small, manageable chunks. Then, follow what is going on with our variable all the way through the problem.

Example 7

An emergency plumber charges \$65 as a call-out fee plus an additional \$75 per hour per hour. He arrives at a house at 9 : 30 and works to repair a water tank. If the total repair bill is \$196.25, at what time was the repair completed?

In order to solve this problem, we collect the information from the text and convert it to an equation.

Unknown time taken in hours this will be our x

The bill is made up of two parts: a call out fee and a per-hour fee. The call out is a flat fee, and independent of x . The per-hour part depends on x . Lets look at how this works algebraically.

$$\begin{array}{r} \$65 \text{ as a call-out fee} \\ \text{Plus an additional } \$75 \text{ per hour} \end{array} \qquad \begin{array}{r} 65 \\ +75x \end{array}$$

So the bill, made up from the call out fee plus the per hour charge times the hours taken creates the following equation.

$$\text{Total Bill} = 65 + 75x$$

Lastly, we look at the final piece of information. The total on the bill was \$196.25. So our final equation is:

$$196.25 = 65 + 75x$$

We solve for x :

$$\begin{array}{r} 196.25 = \cancel{65} + 75x \\ - 65 = \cancel{-65} \\ \hline 131.25 = 75x \end{array}$$

To isolate x first subtract 65 from both sides :

Divide both sides by 75

$$\frac{131.25}{75} = x = 1.75$$

The time taken was one and three quarter hours.

Solution

The repair job was completed at 11:15AM.

Example 8

When Asia was young her Daddy marked her height on the door frame every month. Asias Daddy noticed that between the ages of one and three, he could predict her height (in inches) by taking her age in months, adding 75inches and multiplying the result by one-third. Use this information to determine the following

a) Write an equation linking her predicted height, h , with her age in months, m .

b) Determine her predicted height on her second birthday.

c) Determine at what age she is predicted to reach three feet tall.

a) To convert the text to an equation, first determine the type of equation we have. We are going to have an equation that links **two variables**. Our unknown will change, depending on the information we are given. For example, we could solve for height given age, or solve for age given height. However, the text gives us a way to determine **height**. Our equation will start with $h =$.

Next we look at the text.

$$(m + 75)$$

Take her age in months, and add 75.

$$\frac{1}{3}(m + 75)$$

Multiply the result by one-third.

Solution

Our full equation is $h = \frac{1}{3}(m + 75)$.

b) To determine the prediction of Asias height on her second birthday, we substitute $m = 24$ (2 years = 24 months) into our equation and solve for h .

$$h = \frac{1}{3}(24 + 75)$$

Combine terms in parentheses.

$$h = \frac{1}{3}(99)$$

Multiply.

$$h = 33$$

Solution

Asias height on her second birthday was predicted to be 33 inches .

c) To determine the predicted age when she reached three feet, substitute $h = 36$ into the equation and solve for m .

$$36 = \frac{1}{3}(m + 75)$$

Multiply both sides by 3.

$$108 = m + 75$$

Subtract 75 from both sides.

$$33 = m$$

Solution

Asia was predicted to be 33 months old when her height was three feet.

Example 9



To convert temperatures in Fahrenheit to temperatures in Celsius follow the following steps: Take the temperature in Fahrenheit and subtract 32. Then divide the result by 1.8 and this gives temperature in degrees Celsius.

a) Write an equation that shows the conversion process.

b) Convert 50 degrees Fahrenheit to degrees Celsius.

c) Convert 25 degrees Celsius to degrees Fahrenheit.

d) Convert -40 degrees Celsius to degrees Fahrenheit.

a) The text gives the process to convert Fahrenheit to Celsius. We can write an equation using two variables. We will use f for temperature in Fahrenheit, and c for temperature in Celsius. Follow the text to see it work.

$$C = \frac{(F - 32)}{1.8}$$

Take the temperature in Fahrenheit and subtract 32.

Then divide the result by 1.8.

This gives temperature in degrees Celsius.

In order to convert from one temperature scale to another, simply substitute in for the **known** temperature and solve for the **unknown**.

b) To convert 50 degrees Fahrenheit to degrees Celsius substitute $F = 50$ into the equation.

$$C = \frac{50 - 32}{1.8}$$

Evaluate numerator.

$$C = \frac{18}{1.8}$$

Perform division operation.

Solution

$C = 10$, so 50 degrees Fahrenheit is equal to 10 degrees Celsius.

ci) To convert 25 degrees Celsius to degrees Fahrenheit substitute $C = 25$ into the equation:

$$25 = \frac{F - 32}{1.8}$$

Multiply both sides by 1.8

$$45 = F - 32$$

$$+32 = +32$$

Add 32 to both sides.

$$77 = F$$

Solution

25 degrees Celsius is equal to 77 degrees Fahrenheit.

d) To convert -40 degrees Celsius to degrees Fahrenheit substitute $C = -40$ into the equation.

$$-40 = \frac{F - 32}{1.8}$$

Multiply both sides by 1.8.

$$-72 = F - 32$$

$$+32 = +32$$

Add 32 to both sides.

$$-40 = F$$

Solution

-40 degrees Celsius is equal to -40 degrees Fahrenheit.

Lesson Summary

- Some equations require more than one operation to solve. Generally it, is good to go from the outside in. If there are parentheses around an expression with a variable in it, cancel what is outside the parentheses first.
- Terms with the same variable in them (or no variable in them) are **like terms**. **Combine like terms** (adding or subtracting them from each other) to simplify the expression and solve for the unknown.

Review Questions

1. Solve the following equations for the unknown variable.

- $1.3x - 0.7x = 12$
- $6x - 1.3 = 3.2$
- $5x - (3x + 2) = 1$
- $4(x + 3) = 1$
- $5q - 7 = \frac{2}{3}$
- $\frac{3}{5}x + \frac{5}{2} = \frac{2}{3}$
- $s - \frac{3s}{8} = \frac{5}{6}$
- $0.1y + 11 = 0$
- $\frac{5q-7}{12} = \frac{2}{3}$
- $\frac{5(q-7)}{12} = \frac{2}{3}$
- $33t - 99 = 0$
- $5p - 2 = 32$

2. Jade is stranded downtown with only \$10 to get home. Taxis cost \$0.75 per mile, but there is an additional \$2.35 hire charge. Write a formula and use it to calculate how many miles she can travel with her money. Determine how many miles she can ride.

3. Jasmin's Dad is planning a surprise birthday party for her. He will hire a bouncy castle, and will provide party food for all the guests. The bouncy castle costs \$150 dollars for the afternoon, and the food will cost \$3.00 per person. Andrew, Jasmin's Dad, has a budget of \$300. Write an equation to help him determine the maximum number of guests he can invite.

Review Answers

- $x = 20$
 - $x = 0.75$
 - $x = 1.5$
 - $x = -2.75$
 - $q = \frac{23}{15}$
 - $= -\frac{55}{18}$
 - $s = \frac{4}{3}$
 - $y = -110$
 - $q = 3$
 - $q = \frac{43}{5}$
 - $t = 3$
 - $p = \frac{34}{5}$
- $0.75x + 2.35 = 10 ; x = 10.2$ miles
- $3x + 150 = 300 ; x = 50$ guests

2.5 Multi-Step Equations

Learning Objectives

- Solve a multi-step equation by combining like terms.
- Solve a multi-step equation using the distributive property.
- Solve real-world problems using multi-step equations.

Solving Multi-Step Equations by Combining Like Terms

We have seen that when we solve for an unknown variable, it can be a simple matter of moving terms around in one or two steps. We can now look at solving equations that take several steps to isolate the unknown variable. Such equations are referred to as **multi-step equations**.

In this section, we will simply be combining the steps we already know how to do. Our goal is to end up with all the constants on one side of the equation and all of the variables on the other side. We will do this by collecting like terms. Don't forget, like terms have the same combination of variables in them.

Example 1

Solve $\frac{3x+4}{3} - 5x = 6$

This problem involves a fraction. Before we can combine the variable terms we need to deal with it. Let's put all the terms on the left over a common denominator of three.

$$\frac{3x+4}{3} - \frac{15x}{3} = 6$$

$$\frac{3x+4-15x}{3} = 6$$

$$\frac{4-12x}{3} = 6$$

$$4-12x = 18$$

$$-12x = 14$$

$$\frac{-12}{-12}x = -\frac{14}{12}$$

Next we combine the fractions.

Combine like terms.

Multiply both sides by 3.

Subtract 4 from both sides.

Divide both sides by -12

Solution

$$x = -\frac{7}{6}$$

Solving Multi-Step Equations Using the Distributive Property

You have seen in some of the examples that we can choose to divide out a constant or distribute it. The choice comes down to whether or not we would get a fraction as a result. We are trying to simplify the expression. If we can divide out large numbers without getting a fraction, then we avoid large coefficients. Most of the time, however, we will have to distribute and then collect like terms.

Example 2

Solve $17(3x+4) = 7$

This equation has the x buried in parentheses. In order to extract it we can proceed in one of two ways. We can either distribute the seventeen on the left, or divide both sides by seventeen to remove it from the left. If we divide by seventeen, however, we will end up with a fraction. We wish to avoid fractions if possible!

$17(3x+4) = 7$	Distribute the 17.
$51x + 68 = 7$	
$-68 = -68$	Subtract 68 from both sides.
$51x = -61$	Divide by 51.

Solution

$$x = -\frac{61}{51}$$

Example 3

Solve $4(3x-4) - 7(2x+3) = 3$

This time we will need to collect like terms, but they are hidden inside the brackets. We start by expanding the parentheses.

$12x - 16 - 14x - 21 = 3$	Collect the like terms ($12x$ and $-14x$).
$(12x - 14x) + (-16 - 21) = 3$	Evaluate each set of like terms.
$-2x - 37 = 3$	
$+37 + 37$	Add 37 to both sides.
$-2x = 40$	
$\frac{-2x}{-2} = \frac{40}{-2}$	Divide both sides by -2 .

Solution

$$x = -20$$

Example 4

Solve the following equation for x .

$$0.1(3.2 + 2x) + \frac{1}{2}\left(3 - \frac{x}{5}\right) = 0$$

This function contains both fractions and decimals. We should convert all terms to one or the other. It is often easier to convert decimals to fractions, but the fractions in this equation are easily moved to decimal form. Decimals do not require a common denominator!

Rewrite in decimal form.

$$\begin{aligned}
 0.1(3.2 + 2x) + 0.5(3 - 0.2x) &= 0 \\
 0.32 + 0.2x + 1.5 - 0.1x &= 0 \\
 (0.2x - 0.1x) + (0.32 + 1.5) &= 0 \\
 0.1x + 1.82 &= 0 \\
 -1.82 &- 1.82 \\
 0.1x &= -1.82 \\
 \frac{0.1x}{0.1} &= \frac{-1.82}{0.1}
 \end{aligned}$$

Multiply out decimals:

Collect like terms:

Evaluate each collection:

Subtract 1.82 from both sides

Divide by -0.1

Solution

$$x = 18.2$$

Solve Real-World Problems Using Multi-Step Equations

Real-world problems require you to translate from a problem in words to an equation. First, look to see what the equation is asking. What is the **unknown** you have to solve for? That will determine the quantity we will use for our **variable**. The text explains what is happening. Break it down into small, manageable chunks, and follow what is going on with our variable all the way through the problem.

Example 5

A growers cooperative has a farmers market in the town center every Saturday. They sell what they have grown and split the money into several categories. 8.5% of all the money taken is removed for sales tax. \$150 is removed to pay the rent on the space they occupy. What remains is split evenly between the seven growers. How much money is taken in total if each grower receives a \$175 share?

Let us translate the text above into an equation. The unknown is going to be the total money taken in dollars. We will call this x .

"8.5% of all the money taken is removed for sales tax". This means that 91.5% of the money remains. This is $0.915x$.

$$(0.915x - 150)$$

\$150 is removed to pay the rent on the space they occupy

$$\frac{0.915x - 150}{7}$$

What remains is split evenly between the 7 growers

If each growers share is \$175, then we can write the following equation.

$$\frac{0.915x - 150}{7} = 175$$

$$0.915x - 150 = 1225$$

$$0.915x = 1375$$

$$\frac{0.915x}{0.915} = \frac{1375}{0.915}$$

$$= 1502.7322 \dots$$

Multiply by both sides 7.

Add 150 to both sides.

Divide by 0.915.

Round to two decimal places.

Solution

If the growers are each to receive a \$175 share then they must take at least \$1,502.73.



Example 6

A factory manager is packing engine components into wooden crates to be shipped on a small truck. The truck is designed to hold sixteen crates, and will safely carry a 1200lb cargo. Each crate weighs twelve lbs empty. How much weight should the manager instruct the workers to put in each crate in order to get the shipment weight as close as possible to 1200lbs?

The unknown quantity is the weight to put in each box. This is x . Each crate, when full will weigh:

$$(x + 12)$$

$$16(x + 12)$$

$$16(x + 12) = 1200$$

$$x + 12 = 75$$

$$x = 63$$

16 crates must weigh.

And this must equal 16 lbs.

Isolate x first, divide both sides by 16.

Next subtract 12 from both sides.

Solution

The manager should tell the workers to put 63 lbs of components in each crate.

Ohms Law

The electrical current, I (amps), passing through an electronic component varies directly with the applied voltage, V (volts), according to the relationship:

$V = I \cdot R$ where R is the resistance (measured in Ohms - Ω)



Example 7

A scientist is trying to deduce the resistance of an unknown component. He labels the resistance of the unknown component $x\Omega$. The resistance of a circuit containing a number of these components is $(5x + 20)\Omega$. If a 120 volt potential difference across the circuit produces a current of 2.5 amps, calculate the resistance of the unknown component.

Substitute $V = 120$, $I = 2.5$ and $R = 5x + 20$ into $V = I \cdot R$:

$$\begin{aligned} 120 &= 2.5(5x + 20) && \text{Distribute the 2.5.} \\ 120 &= 12.5x + 50 && \text{Subtract 50 from both sides.} \\ -50 &= -50 \\ 70 &= 12.5x && \text{Divide both sides by 12.5.} \\ \frac{70}{12.5} &= \frac{12.5x}{12.5} \\ 5.6\Omega &= x \end{aligned}$$

Solution

The unknown components have a resistance of 5.6Ω .

Distance, Speed and Time

The speed of a body is the distance it travels per unit of time. We can determine how far an object moves in a certain amount of time by multiplying the speed by the time. Here is our equation.

$$\text{distance} = \text{speed} \times \text{time}$$

Example 8

Shanices car is traveling 10 miles per hour slower than twice the speed of Brandons car. She covers 93 miles in 1 hour 30 minutes. How fast is Brandon driving?

Here we have two unknowns in this problem. Shanices speed and Brandons speed. We do know that Shanices speed is ten less than twice Brandons speed. Since the question is asking for Brandons speed, it is his speed in miles per hour that will be x .

Substituting into the distance time equation yields:

$$\begin{aligned} 93 &= 2x - 10 \times 1.5 && \text{Divide by 1.5.} \\ 62 &= 2x - 10 \\ +10 &= +10 && \text{Add 10 to both sides.} \\ 72 &= 2x \\ \frac{72}{2} &= \frac{2x}{2} && \text{Divide both sides by 2.} \\ 36 &= x \end{aligned}$$

Solution

Peter is driving at 36 miles per hour.

This example may be checked by considering the situation another way: We can use the fact that Shanice's covers 93 miles in 1 hour 30 minutes to determine her speed (we will call this y as x has already been defined as Brandons speed):

$$93 = y \cdot 1.5$$

$$\frac{93}{1.5} = \frac{1.5y}{1.5}$$

$$y = 62\text{mph}$$

Divide both sides by 1.5.

We can then use this information to determine Shanices speed by converting the text to an equation.
Shanices car is traveling at 10miles per hour slower than twice the speed of Peters car
 Translates to

$$y = 2x - 10$$

It is then a simple matter to substitute in our value in fory and then solve for x:

$$62 = (2x - 10)$$

$$+ 10 + 10$$

$$72 = 2x$$

$$72 = 2x$$

$$\frac{72}{2} = \frac{2x}{2}$$

$$x = 36 \text{ miles per hour.}$$

Add 10 to both sides.

Divide both sides by 2.

Solution

Brandon is driving at 36 milesperhour.

You can see that we arrive at exactly the same answer whichever way we solve the problem. In algebra, there is almost always more than one method of solving a problem. If time allows, it is an excellent idea to try to solve the problem using two different methods and thus confirm that you have calculated the answer correctly.

Speed of Sound

The speed of sound in dry air, v, is given by the following equation.

$$v = 331 + 0.6T \text{ where } T \text{ is the temperature in Celsius and } v \text{ is the speed of sound in meters per second.}$$

Example 9

Tashi hits a drainpipe with a hammer and 250 metersaway Minh hears the sound and hits his own drainpipe. Unfortunately, there is a one second delay between him hearing the sound and hitting his own pipe. Tashi accurately measures the time from her hitting the pipe and hearing Mihns pipe at 2.46 seconds . What is the temperature of the air?

This complex problem must be carefully translated into equations:

$$\text{Distance traveled} = (331 + 0.6T) \times \text{time}$$

$$\text{time} = (2.46 - 1)$$

$$\text{Distance} = 2 \times 250$$

Do not forget, for one second the sound is not traveling

Our equation is:

$$\begin{aligned}
 2(250) &= (331 + 0.6T) \cdot (2.46 - 1) && \text{Simplify terms.} \\
 \frac{500}{1.46} &= \frac{1.46(331 + 0.6T)}{1.46} && \text{Divide by 1.46.} \\
 342.47 - 331 &= 331 + 0.6T - 331 && \text{Subtract 331 from both sides.} \\
 \frac{11.47}{0.6} &= \frac{0.6T}{0.6} && \text{Divide by 0.6.} \\
 19.1 &= T
 \end{aligned}$$

Solution

The temperature is 19.1 degrees Celsius.

Lesson Summary

- If dividing a number outside of parentheses will produce fractions, it is often better to use the **Distributive Property** (for example, $3(x + 2) = 3x + 6$) to expand the terms and then combine like terms to solve the equation.

Review Questions

- Solve the following equations for the unknown variable.
 - $3(x - 1) - 2(x + 3) = 0$
 - $7(w + 20) - w = 5$
 - $9(x - 2) = 3x + 3$
 - $2\left(5a - \frac{1}{3}\right) = \frac{2}{7}$
 - $\frac{2}{9}\left(i + \frac{2}{3}\right) = \frac{2}{5}$
 - $4\left(v + \frac{1}{4}\right) = \frac{35}{2}$
 - $\frac{s-4}{11} = \frac{2}{5}$
 - $\frac{p}{16} - \frac{2p}{3} = \frac{1}{9}$
- An engineer is building a suspended platform to raise bags of cement. The platform has a mass of 200 kg, and each bag of cement is 40 kg. He is using two steel cables, each capable of holding 250 kg. Write an equation for the number of bags he can put on the platform at once, and solve it.
- A scientist is testing a number of identical components of unknown resistance which he labels $x\Omega$. He connects a circuit with resistance $(3x + 4)\Omega$ to a steady 12 Volt supply and finds that this produces a current of 1.2 Amps. What is the value of the unknown resistance?
- Lydia inherited a sum of money. She split it into five equal chunks. She invested three parts of the money in a high interest bank account which added 10% to the value. She placed the rest of her inheritance plus \$500 in the stock market but lost 20% on that money. If the two accounts end up with exactly the same amount of money in them, how much did she inherit?
- Pang drove to his mothers house to drop off her new TV. He drove at 50 miles per hour there and back, and spent 10 minutes dropping off the TV. The entire journey took him 94 minutes. How far away does his mother live?

Review Answers

- (a) $x = 9$

(b) $w = -22.5$

(c) $x = 3.5$

(d) $a = \frac{2}{21}$

(e) $i = \frac{17}{15}$

(f) $v = \frac{33}{8}$

(g) $s = \frac{42}{5}$

(h) $p = -\frac{16}{87}$

2. $2(250) = 200 + 40x; x = 7.5 \rightarrow 7$ bags

3. 2Ω

4. \$1,176.50

5. 35 miles

2.6 Equations with Variables on Both Sides

Learning Objectives

- Solve an equation with variables on both sides.
- Solve an equation with grouping symbols.
- Solve real-world problems using equations with variables on both sides.

Solve an Equation with Variables on Both Sides

When the variable appears on both sides of the equation, we need to manipulate our equation such that all variables appear on one side, and only constants remain on the other.

Example 1

Dwayne was told by his chemistry teacher to measure the weight of an empty beaker using a balance. Dwayne found only one lb weights, and so devised the following way of balancing the scales.



Knowing that each weight is one lb, calculate the weight of one beaker.

Solution

We know that the system balances, so the weights on each side must be equal. We can write an algebraic expression based on this fact. The unknown quantity, the weight of the beaker, will be our x . We can see that on the left hand scale we have one beaker and four weights. On the right scale, we have four beakers and three weights. The balancing of the scales is analogous to the balancing of the following equation.

$$x + 4 = 4x + 3$$

One beaker plus 4 lbs **equals** 4 beakers plus 3 lbs

To solve for the weight of the beaker, we want all the constants (numbers) on one side and all the variables (terms with x in) on the other side. Look at the balance. There are more beakers on the right and more weights on the left. We will aim to end up with only x terms (beakers) on the right, and only constants (weights) on the left.

$$\begin{aligned}x + 4 &= 4x + 3 \\ - 3 &= -3\end{aligned}$$

Subtract 3 from both sides.

$$\begin{aligned}x + 1 &= 4x \\ - x &= -x\end{aligned}$$

Subtract x from both sides.

$$\begin{aligned}1 &= 3x \\ \frac{1}{3} &= \frac{3x}{3} \\ x &= \frac{1}{3}\end{aligned}$$

Divide both sides by 3.


Answer The weight of the beaker is one-third of a pound.

We can do the same with the real objects as we have done with the equation. Our first action was to subtract three from both sides of the equals sign. On the balance, we could remove a certain number of weights or beakers from each scale. Because we remove the same number of weights from each side, we know the scales will still balance.

On the balance, we could remove three weights from each scale. This would leave one beaker and one weight on the left and four beakers on the right (in other words $x + 1 = 4x$):




The next step we could do is remove one beaker from each scale leaving only one weight on the left and three beakers on the right and you will see our final equation: $1 = 3x$.



Looking at the balance, it is clear that the weight of the beaker is one-third of a pound.

Example 2

Sven was told to find the weight of an empty box with a balance. Sven found one lb weights and five lb weights. He placed two one lb weights in three of the boxes and with a fourth empty box found the following way of balancing the scales.



Knowing that small weights are one lb and big weights are five lbs, calculate the weight of one box.

We know that the system balances, so the weights on each side must be equal. We can write an algebraic expression based on this equality. The unknown quantity, the weight of each empty box, in pounds, will be our x . A box with two 1 lb weights in it weighs $(x + 2)$. Here is the equation.

$$\begin{array}{r}
 3(x+2) = x+3(5) \\
 3x+6 = x+5 \\
 -x = -x \\
 \hline
 2x+6 = 15 \\
 -6 = -6 \\
 \hline
 2x = 9 \\
 x = 4.5
 \end{array}$$

Distribute the 3.

Subtract x from both sides.

Subtract 6 from both sides.

Divide both sides by 2.

Solution

Each box weighs 4.5 lbs.

Solve an Equation with Grouping Symbols

When we have a number of like terms on one side of the equal sign we **collect like terms** then add them in order to solve for our variable. When we move variables from one side of the equation to the other we sometimes call it grouping symbols. Essentially we are doing exactly what we would do with the constants. We can add and subtract variable terms just as we would with numbers. In fractions, occasionally we will have to multiply and divide by variables in order to get them all on the numerator.

Example 3

Solve $3x + 4 = 5x$

Solution

This equation has x on both sides. However, there is only a number term on the left. We will therefore move all the x terms to the right of the equal sign leaving the constant on the left.

$$\begin{array}{r}
 3x+4 = 5x \\
 -3x \quad -3x \\
 \hline
 4 = 2x \\
 \hline
 \frac{4}{2} = \frac{2x}{2}
 \end{array}$$

Subtract 3x from both sides.

Divide by 2

Solution

$$x = 2$$

Example 4

Solve $9x = 4 - 5x$

This time we will collect like terms (x terms) on the left of the equal sign.

$$\begin{array}{r} 9x = 4 - 5x \\ + 5x \quad + 5x \\ \hline \end{array}$$

Add $5x$ to both sides.

$$14x = 4$$

$$14x = 4$$

$$\frac{14x}{14} = \frac{4}{14}$$

$$x = \frac{2}{7}$$

Divide by 14.

Solution

$$x = \frac{2}{7}$$

Example 5

Solve $3x + 2 = \frac{5x}{3}$

This equation has x on both sides and a fraction. It is always easier to deal with equations that do not have fractions. The first thing we will do is get rid of the fraction.

$$\begin{array}{r} 3x + 2 = \frac{5x}{3} \\ 3(3x + 2) = 5x \end{array}$$

Multiply both sides by 3.

Distribute the 3.

$$\begin{array}{r} 9x + 6 = 5x \\ - 9x \quad - 9x \\ \hline \end{array}$$

Subtract $9x$ from both sides :

Divide by -4 .

$$\begin{array}{r} \frac{6}{-4} = \frac{-4x}{-4} \\ \frac{6}{-4} = x \\ -\frac{3}{2} = x \end{array}$$

Solution

$$x = -1.5$$

Example 6

Solve $7x + 2 = \frac{5x-3}{6}$

Again we start by eliminating the fraction.

$$7x + 2 = \frac{5x - 3}{6}$$

$$6(7x + 2) = \frac{5x - 3}{6} \cdot 6$$

Multiply both sides by 6.

$$6(7x + 2) = 5x - 3$$

Distribute the 6.

$$42x + 12 = \cancel{5x} - 3$$

Subtract 5x from both sides :

$$-5x \quad -\cancel{5x}$$

$$37x + \cancel{12} = -3$$

Subtract 12 from both sides.

$$-\cancel{12} \quad -12$$

$$37x = -15$$

Divide by -37 .

$$\frac{37x}{x} = \frac{-15}{37}$$

Solution

$$x = -\frac{15}{37}$$

Example 7

Solve the following equation for x .

$$\frac{14x}{(x+3)} = 7$$

The form of the left hand side of this equation is known as a **rational function** because it is the ratio of two other functions ($14x$) and $(x+3)$. However, we wish simply to solve for x so we start by eliminating the fraction. We do this as we have always done, by multiplying by the denominator.

$$\frac{14x}{(x+3)}(x+3) = 7(x+3)$$

Multiply by $(x+3)$.

$$14x = 7(x+3)$$

Distribute the 7.

$$14x = \cancel{7x} + 21$$

Subtract $7x$ from both sides.

$$-7x = -\cancel{7x}$$

$$7x = 21$$

Divide both sides by 7

$$\frac{7x}{7} = \frac{21}{7}$$

$$x = 3$$

Solve Real-World Problems Using Equations with Variables on Both Sides

Build your skills in translating problems from words to equations. What is the equation asking? What is the **unknown** variable? What quantity will we use for our variable?

The text explains what is happening. Break it down into small, manageable chunks, and follow what is going on with our variable all the way through the problem.

More on Ohms Law

The electrical current, I (amps), passing through an electronic component varies directly with the applied voltage, V (volts), according to the relationship:

$$V = I \cdot R \quad \text{where } R \text{ is the resistance (measured in Ohms)}$$

The resistance R of a number of components wired in a **series** (one after the other) is given by: $R = r_1 + r_2 + r_3 + r_4 + \dots$



Example 8

In an attempt to find the resistance of a new component, a scientist tests it in series with standard resistors. A fixed voltage causes a 4.8amp current in a circuit made up from the new component plus a 15Ω resistor in series. When the component is placed in a series circuit with a 50Ω resistor the same voltage causes a 2.0amp current to flow. Calculate the resistance of the new component.

This is a complex problem to translate, but once we convert the information into equations it is relatively straight forward to solve. Firstly we are trying to find the resistance of the new component (in Ohms, Ω). This is our x . We do not know the voltage that is being used, but we can leave that as simple V . Our first situation has the unknown resistance plus 15Ω . The current is 4.8 amps. Substitute into the formula $V = I \cdot R$.

$$V = 4.8(x + 15)$$

Our second situation has the unknown resistance plus 50Ω . The current is 2.0 amps.

$$V = 2(x + 50)$$

We know the voltage is fixed, so the V in the first equation must equal the V in the second. This means that:

$$4.8(x + 15) = 2(x + 50)$$

$$4.8x + 72 = 2x + 100$$

$$- 2x \quad - 2x$$

$$2.8x + 72 = 100$$

$$- 72 \quad - 72$$

$$2.8x = 28$$

$$\frac{2.8x}{2.8} = \frac{28}{2.8}$$

$$x = 10$$

Distribute the constants.

Subtract $2x$ from both sides.

Subtract 72 from both sides.

Divide both sides by 2.8 .

Solution

The resistance of the component is 10Ω .

Lesson Summary

- If an unknown variable appears on both sides of an equation, distribute as necessary. Then subtract (or add) one term to both sides to simplify the equation to have the unknown on only one side.

Review Questions

1. Solve the following equations for the unknown variable.

(a) $3(x - 1) = 2(x + 3)$

(b) $7(x + 20) = x + 5$

(c) $9(x - 2) = 3x + 3$

(d) $2\left(a - \frac{1}{3}\right) = \frac{2}{5}\left(a + \frac{2}{3}\right)$

(e) $\frac{2}{7}\left(t + \frac{2}{3}\right) = \frac{1}{5}\left(t - \frac{2}{3}\right)$

(f) $\frac{1}{7}\left(v + \frac{1}{4}\right) = 2\left(\frac{3v}{2} - \frac{5}{2}\right)$

(g) $\frac{y-4}{11} = \frac{2}{5} \cdot \frac{2y+1}{3}$

(h) $\frac{z}{16} = \frac{2(3z+1)}{9}$

(i) $\frac{q}{16} + \frac{q}{6} = \frac{(3q+1)}{9} + \frac{3}{2}$

2. Manoj and Tamar are arguing about how a number trick they heard goes. Tamar tells Andrew to think of a number, multiply it by five and subtract three from the result. Then Manoj tells Andrew to think of a number add five and multiply the result by three. Andrew says that whichever way he does the trick he gets the same answer. What was Andrew's number?
3. I have enough money to buy five regular priced CDs and have \$6 left over. However all CDs are on sale today, for \$4 less than usual. If I borrow \$2, I can afford nine of them. How much are CDs on sale for today?
4. Five identical electronics components were connected in series. A fixed but unknown voltage placed across them caused a 2.3 amp current to flow. When two of the components were replaced with standard 10Ω resistors, the current dropped to 1.9 amps. What is the resistance of each component?
5. Solve the following resistance problems. Assume the same voltage is applied to all circuits.
- Three unknown resistors plus 20Ω give the same current as one unknown resistor plus 70Ω .
 - One unknown resistor gives a current of 1.5 amps and a 15Ω resistor gives a current of 3.0 amps.
 - Seven unknown resistors plus 18Ω gives twice the current of two unknown resistors plus 150Ω .

- (d) Three unknown resistors plus 1.5Ω gives a current of 3.6 amps and seven unknown resistors plus 712Ω resistors gives a current of 0.2 amps.

Review Answers

1. (a) $x = 9$
(b) $x = -22.5$
(c) $x = 3.5$
(d) $a = \frac{7}{12}$
(e) $t = -\frac{34}{9}$
(f) $v = \frac{141}{80}$
(g) $y = -\frac{82}{29}$
(h) $z = -\frac{32}{87}$
(i) $q = -\frac{232}{15}$
2. 9
3. \$7
4. 6.55Ω
5. (a) unknown = 25Ω
(b) unknown = 30Ω
(c) unknown = 94Ω
(d) unknown = 1.213Ω

2.7 Ratios and Proportions

Learning Objectives

- Write and understand a ratio.
- Write and solve a proportion.
- Solve proportions using cross products.

Introduction

Nadia is counting out money with her little brother. She gives her brother all the nickels and pennies. She keeps the quarters and dimes for herself. Nadia has four quarters (worth 25 cents each) and six dimes (worth 10 cents each). Her brother has fifteen nickels (worth 5 cents each) and five pennies (worth one cent each) and is happy because he has more coins than his big sister. How would you explain to him that he is actually getting a bad deal?

Write a ratio

A **ratio** is a way to compare two numbers, measurements or quantities. When we write a ratio, we divide one number by another and express the answer as a fraction. There are two distinct ratios in the problem above. For example, the ratio of the **number** of Nadias coins to her brothers is:

$$\frac{4 + 6}{15 + 5} = \frac{10}{20}$$

When we write a ratio, the correct way is to simplify the fraction.

$$\frac{10}{20} = \frac{\cancel{2} \cdot 5}{\cancel{2} \cdot 20} = \frac{1}{2}$$

In other words, Nadia has half the number of coins as her brother.

Another ratio we could look at in the problem is the **value** of the coins. The value of Nadias coins is $(4 \times 25) + (6 \times 10) = 160$ cents . The value of her brothers coins is $(15 \times 5) + (5 \times 1) = 80$ cents . The ratio of the **value** of Nadias coins to her brothers is:

$$\frac{160}{80} = \frac{2}{1}$$

So the value of Nadias money is twice the value of her brothers.

Notice that even though the denominator is one, it is still written. A ratio with a denominator of one is called a **unit rate**. In this case, it means Nadia is gaining money at twice the rate of her brother.



Example 1

The price of a Harry Potter Book on Amazon.com is \$10.00. The same book is also available used for \$6.50. Find two ways to compare these prices.

Clearly, the cost of a new book is greater than the used book price. We can compare the two numbers using a difference equation:

$$\text{Difference in price} = 10.00 - \$6.50 = \$3.50$$

We can also use a ratio to compare the prices:

$$\frac{\text{new price}}{\text{used price}} = \frac{\$10.00}{\$6.50}$$

$$\frac{10}{6.50} = \frac{1000}{650} = \frac{20}{13}$$

We can cancel the units of \$ as they are the same.

We remove the decimals and simplify the fraction.

Solution

The new book is \$3.50 more than the used book.

The new book costs $\frac{20}{13}$ times the cost of the used book.

**Example 2**

The State Dining Room in the White House measures approximately 48feet long by 36feet wide. Compare the length of room to the width, and express your answer as a ratio.

Solution

$$\frac{48 \text{ feet}}{36 \text{ feet}} = \frac{48}{36} = \frac{4}{3}$$

Example 3

A tournament size shuffleboard table measures 30inches wide by 14feet long. Compare the length of the table to its width and express the answer as a ratio.

We could write the ratio immediately as:

$$\frac{14 \text{ feet}}{30 \text{ inches}}$$

Notice that we cannot cancel the units.

Sometimes it is OK to leave the units in, but as we are comparing two lengths, it makes sense to convert all the measurements to the same units.

Solution

$$\frac{14 \text{ feet}}{30 \text{ inches}} = \frac{14 \times 12 \text{ inches}}{30 \text{ inches}} = \frac{168}{30} = \frac{28}{5}$$



Example 4

A family car is being tested for fuel efficiency. It drives non-stop for 100 miles, and uses 3.2 gallons of gasoline. Write the ratio of distance traveled to fuel used as a **unit rate**.

$$\text{Ratio} = \frac{100 \text{ miles}}{3.2 \text{ gallons}}$$

A unit rate has a denominator of one, so we need to divide both numerator and denominator by 3.2.

$$\text{Unit Rate} = \frac{\left(\frac{100}{3.2}\right) \text{ miles}}{\left(\frac{3.2}{3.2}\right) \text{ gallons}} = \frac{31.25 \text{ miles}}{1 \text{ gallon}}$$

Solution

The ratio of distance to fuel used is $\frac{31.25 \text{ miles}}{1 \text{ gallon}}$ or 31.25 miles per gallon.

Write and Solve a Proportion

When two ratios are equal to each other, we call it a proportion.

$$\frac{10}{15} = \frac{6}{9}$$

This statement is a proportion. We know the statement is true because we can reduce both fractions to $\frac{2}{3}$.

Check this yourself to make sure!

We often use proportions in science and business. For example, when scaling up the size of something. We use them to solve for an unknown, so we will use algebra and label our unknown variable x . We assume that a certain ratio holds true whatever the size of the thing we are enlarging (or reducing). The next few examples demonstrate this.

**Example 5**

A small fast food chain operates 60 stores and makes \$1.2 million profit every year. How much profit would the chain make if it operated 250 stores?

First, we need to write a **ratio**. This will be the ratio of profit to number of stores.

$$\text{Ratio} = \frac{\$1,200,000}{60 \text{ stores}}$$

We now need to determine our unknown, x which will be in dollars. It is the profit of 250 stores. Here is the ratio that compares unknown dollars to 250 stores.

$$\text{Ratio} = \frac{\$x}{250 \text{ stores}}$$

We now write equal ratios and solve the resulting **proportion**.

$$\frac{\$1,200,000}{60 \text{ stores}} = \frac{\$x}{250 \text{ stores}} \text{ or } \frac{1,200,000}{60} = \frac{x}{250}$$

Note that we can drop the units not because they are the same on the numerator and denominator, but because they are the same on both sides of the equation.

$$\frac{1,200,000}{60} = \frac{x}{250}$$

Simplify fractions.

$$20,000 = \frac{x}{250}$$

Multiply both sides by 250.

$$5,000,000 = x$$

Solution

If the chain operated 250 stores the annual profit would be 5 million dollars .



Example 6

A chemical company makes up batches of copper sulfate solution by adding 250 kg of copper sulfate powder to 1000 liters of water. A laboratory chemist wants to make a solution of identical concentration, but only needs 350 ml (0.35 liters) of solution. How much copper sulfate powder should the chemist add to the water?

First we write our ratio. The mass of powder divided by the volume of water used by the chemical company.

$$\text{Ratio} = \frac{250 \text{ kg}}{1000 \text{ liters}}$$

$$\text{We can reduce this to : } \frac{1 \text{ kg}}{4 \text{ liters}}$$

Our unknown is the mass in kilograms of powder to add. This will be x . The volume of water will be 0.35 liters .

$$\text{Ratio} = \frac{x \text{ kg}}{0.35 \text{ liters}}$$

Our proportion comes from setting the two ratios equal to each other:

$$\frac{1 \text{ kg}}{4 \text{ liters}} = \frac{x \text{ kg}}{0.35 \text{ liters}} \text{ which becomes } \frac{1}{4} = \frac{x}{0.35}$$

We now solve for x .

$$\begin{aligned} \frac{1}{4} &= \frac{x}{0.35} \\ 0.35 \cdot \frac{1}{4} &= \frac{x}{0.35} \cdot 0.35 \\ x &= 0.0875 \end{aligned}$$

Multiply both sides by 0.35.

Solution

The mass of copper sulfate that the chemist should add is 0.0875 kg or 87.5 grams .

Solve Proportions Using Cross Products

One neat way to simplify proportions is to cross multiply. Consider the following proportion.

$$\frac{16}{4} = \frac{20}{5}$$

If we want to eliminate the fractions, we could multiply both sides by 4 and then multiply both sides by 5. In fact we *could* do both at once:

$$\begin{aligned} 4 \cdot 5 \cdot \frac{16}{4} &= 4 \cdot 5 \cdot \frac{20}{5} \\ 5 \cdot 16 &= 4 \cdot 20 \end{aligned}$$

Now comparing this to the proportion we started with, we see that the denominator from the left hand side ends up multiplying with the numerator on the right hand side.

You can also see that the denominator from the *right* hand side ends up multiplying the numerator on the *left* hand side.

In effect the two denominators have *multiplied* across the equal sign:

$$\frac{16}{4} \times \frac{20}{5}$$

=

$$5 \cdot 16 = 4 \cdot 20$$

This movement of denominators is known as **cross multiplying**. It is extremely useful in solving proportions, especially when the unknown variable is on the denominator.

Example 7

Solve the proportion for x .

$$\frac{4}{3} = \frac{9}{x}$$

Cross multiply:

$$x \cdot 4 = 9 \cdot 3$$

$$\frac{4x}{4} = \frac{27}{4}$$

Divide both sides by 4.

Solution

$$x = 6.75$$

Example 8

Solve the following proportion for x .

$$\frac{0.5}{3} = \frac{56}{x}$$

Cross multiply:

$$\begin{aligned} x \cdot 0.5 &= 56 \cdot 3 \\ \frac{0.5x}{0.5} &= \frac{168}{0.5} \end{aligned}$$

Divide both sides by 0.5.

Solution:

$$x = 336$$

Solve Real-World Problems Using Proportions

When we are faced with a word problem that requires us to write a proportion, we need to identify both the unknown (which will be the quantity we represent as x) and the ratio which will stay fixed.

**Example 9**

A cross-country train travels at a steady speed. It covers 15 miles in 20 minutes. How far will it travel in 7 hours assuming it continues at the same speed?

This example is a Distance = speed \times time problem. We came across a similar problem in Lesson 3.3. Recall that the speed of a body is the quantity distance/time. This will be our ratio. We simply plug in the known quantities. We will, however convert to hours from minutes.

$$\text{Ratio} = \frac{15 \text{ miles}}{20 \text{ minutes}} = \frac{15 \text{ miles}}{\frac{1}{3} \text{ hour}}$$

This is a very awkward looking ratio, but since we will be cross multiplying we will leave it as it is. Next, we set up our proportion.

$$\frac{15 \text{ miles}}{\frac{1}{3} \text{ hour}} = \frac{x \text{ miles}}{7 \text{ hours}}$$

Cancel the units and cross-multiply.

$$\begin{aligned} 7 \cdot 15 &= \frac{1}{3} \cdot x \\ 3 \cdot 7 \cdot 15 &= 3 \cdot \frac{1}{3} \cdot x \\ 315 &= x \end{aligned}$$

Multiply both sides by 3.

Solution

The train will travel 315 miles in 7 hours .

Example 10

Rain is falling at 1 inch every 1.5 hours. How high will the water level be if it rains at the same rate for 3 hours?

Although it may not look it, this again uses the Distance = speed \times time relationship. The distance the water rises in inches will be our x . The ratio will again be $\frac{\text{distance}}{\text{time}}$.

$$\begin{aligned}\frac{1 \text{ inch}}{1.5 \text{ hours}} &= \frac{x \text{ inch}}{3 \text{ hours}} \\ \frac{3(1)}{1.5} &= \frac{1.5x}{1.5} \\ 2 &= x\end{aligned}$$

Cancel units and cross multiply.

Divide by 1.5

Solution

The water will be 2 inches high if it rains for 3 hours .

Example 11

In the United Kingdom, Alzheimers disease is said to affect one in fifty people over 65 years of age. If approximately 250000 people over 65 are affected in the UK, how many people over 65 are there in total?

The fixed ratio in this case will be the 1 person in 50. The unknown (x) is the number of persons over 65. Note that in this case, the ratio does not have units, as they will cancel between the numerator and denominator.

We can go straight to the proportion.

$$\begin{aligned}\frac{1}{50} &= \frac{250000}{x} \\ 1 \cdot x &= 250000 \cdot 50 \\ x &= 12,500,000\end{aligned}$$

Cross multiply :

Solution

There are approximately 12.5 million people over the age of 65.

Lesson Summary

- A **ratio** is a way to compare two numbers, measurements or quantities by dividing one number by the other and expressing the answer as a fraction. $\frac{2}{3}$, $\frac{32 \text{ miles}}{1.4 \text{ gallons}}$, and $\frac{x}{13}$ are all ratios.
- A **proportion** is formed when two ratios are set equal to each other.
- **Cross multiplication** is useful for solving equations in the form of proportions. To cross multiply, multiply the bottom of each ratio by the top of the other ratio and set them equal. For instance, cross multiplying

$$\frac{11}{5} \times \frac{x}{3}$$

results in $11 \cdot 3 = 5 \cdot x$.

Review Questions

- Write the following comparisons as ratios. Simplify fractions where possible.
 - \$150 to \$3
 - 150 boys to 175 girls
 - 200 minutes to 1 hour
 - 10 days to 2 weeks
- Write the following ratios as a unit rate.
 - 54 hotdogs to 12 minutes
 - 5000 lbs to $250in^2$
 - 20 computers to 80 students
 - 180 students to 6 teachers
 - 12 meters to 4 floors
 - 18 minutes to 15 appointments
- Solve the following proportions.
 - $\frac{13}{6} = \frac{5}{x}$
 - $\frac{1.25}{7} = \frac{3.6}{x}$
 - $\frac{6}{19} = \frac{x}{11}$
 - $\frac{1}{1} = \frac{0.01}{5}$
 - $\frac{300}{4} = \frac{x}{99}$
 - $\frac{2.75}{9} = \frac{x}{(\frac{2}{5})}$
 - $\frac{1.3}{4} = \frac{x}{1.3}$
 - $\frac{0.1}{1.01} = \frac{1.9}{x}$
- A restaurant serves 100 people per day and takes \$908. If the restaurant were to serve 250 people per day, what might the taking be?
- The highest mountain in Canada is Mount Yukon. It is $\frac{298}{67}$ the size of Ben Nevis, the highest peak in Scotland. Mount Elbert in Colorado is the highest peak in the Rocky Mountains. Mount Elbert is $\frac{220}{67}$ the height of Ben Nevis and $\frac{44}{48}$ the size of Mont Blanc in France. Mont Blanc is 4800 meters high. How high is Mount Yukon?
- At a large high school it is estimated that two out of every three students have a cell phone, and one in five of all students have a cell phone that is one year old or less. Out of the students who own a cell phone, what proportion owns a phone that is more than one year old?

Review Answers

- $\frac{50}{1}$
 - $\frac{6}{7}$
 - $\frac{10}{3}$
 - $\frac{5}{7}$
- 4.5 hot-dogs per minute
 - 20 lbs per in^2
 - 0.25 computers per student
 - 30 students per teacher
 - 3 meters per floor
 - 1.2 minutes per appointment
- $x = \frac{30}{13}$
 - $x = 20.16$
 - $x = \frac{66}{19}$
 - $x = 500$

(e) $x = 7425$

(f) $x = \frac{11}{162}$

(g) $x = 0.4225$

(h) $x = \frac{100}{1919}$

4. \$2270

5. 5960 meters .

6. $\frac{3}{10}$ or 30%

CHAPTER **3** Graphing Linear Equations

Chapter Outline


- 3.1 THE COORDINATE PLANE**
 - 3.2 GRAPHS OF LINEAR EQUATIONS**
 - 3.3 GRAPHING USING INTERCEPTS**
 - 3.4 SLOPE AND RATE OF CHANGE**
 - 3.5 GRAPHS USING SLOPE-INTERCEPT FORM**
 - 3.6 DIRECT VARIATION MODELS**
 - 3.7 PROBLEM-SOLVING STRATEGIES - GRAPHS**
-

3.1 The Coordinate Plane

Learning Objectives

- Identify coordinates of points.
- Plot points in a coordinate plane.
- Graph a function given a table.
- Graph a function given a rule.

Introduction



We now make our transition from a number line that stretches in only one dimension (left to right) to something that exists in two dimensions. The **coordinate plane** can be thought of as two number lines that meet at right angles. The horizontal line is called the **x-axis** and the vertical line is the **y-axis**. Together the lines are called the **axes**, and the point at which they cross is called the **origin**. The axes split the coordinate plane into four **quadrants**. The first quadrant (I) contains all the positive numbers from both axes. It is the top right quadrant. The other quadrants are numbered sequentially (II, III, IV) moving counterclockwise from the first.


Identify Coordinates of Points

When given a point on a coordinate plane, it is a relatively easy task to determine its **coordinates**. The coordinates of a point are two numbers written together they are called an **ordered pair**. The numbers describe how far along the x-axis and y-axis the point is. The ordered pair is written in parenthesis, with the **x-coordinate** (also called the **ordinate**) first and the **y-coordinate** (or the **ordinate**) second.

$(1, 7)$	An ordered pair with an x-value of one and a y-value of seven
$(0, 5)$	an ordered pair with an x-value of zero and a y-value of five
$(-2.5, 4)$	An ordered pair with an x-value of -2.5 and a y-value of four
$(-107.2, -0.005)$	An ordered pair with an x-value of -107.2 and a y-value of -0.005 .

The first thing to do is realize that identifying coordinates is just like reading points on a number line, except that now the points do not actually lie **on** the number line! Look at the following example.

Example 1



Find the coordinates of the point labeled P in the diagram to the right.

Imagine you are standing at the origin (the points where the x -axis meets the y -axis). In order to move to a position where P was directly above you, you would move 3 units to the **right** (we say this is in the **positive** x direction).

The x -coordinate of P is $+3$.


Now if you were standing at the three marker on the x -axis, point P would be 7 units above you (above the axis means it is in the **positive** y direction).

The y -coordinate of P is $+7$.

Solution

The coordinates of point P are $(3, 7)$.

Example 2



Find the coordinates of the points labeled Q and R in the diagram to the right.

In order to get to Q we move three units to the right, in the positive x direction, then two units **down**. This time we are moving in the **negative** y direction. The x coordinate of Q is $+3$, the y coordinate of Q is -2 .

The coordinates of R are found in a similar way. The x coordinate is $+5$ (five units in positive x) and the y -coordinate is again -2 .


Solution

$Q(3, -2)$

$R(5, -2)$

Example 3

Triangle ABC is shown in the diagram to the right. Find the coordinates of the vertices A , B and C .



Point A :

x -coordinate = -2

y -coordinate = $+5$

Point B :

x -coordinate = $+3$

y -coordinate = -3

Point C :

x -coordinate = -4

y -coordinate = -1

Solution

$$A(-2, 5)$$

$$B(3, -3)$$

$$C(-4, -1)$$

Plot Points in a Coordinate Plane

Plotting points is a simple matter once you understand how to read coordinates and read the scale on a graph. As a note on scale, in the next two examples pay close attention to the labels on the axes.

Example 4

Plot the following points on the coordinate plane.

$A(2, 7)$


$B(-5, 6)$

$C(-6, 0)$

$D(-3, -3)$

$E(0, 2)$

$F(7, -5)$



Point $A(2, 7)$ is 2 units right, 7 units up. It is in Quadrant I.

Point $B(-5, 6)$ is 5 units left, 6 units up. It is in Quadrant II.

Point $C(-6, 0)$ is 6 units left, 0 units up. It is **on the x axis**.

Point $D(-3, -3)$ is 3 units left, 3 units down. It is in Quadrant III.

Point $E(0, 2)$ is 2 units up from the origin. It is **on the y axis**.

Point $F(7, -5)$ is 7 units right, 5 units down. It is in Quadrant IV.

Example 5

Plot the following points on the coordinate plane.


$A(2.5, 0.5)$

$B(\pi, 1.2)$

$C(2, 1.75)$

$D(0.1, 1.2)$

$E(0, 0)$




Choice of axes is always important. In Example Four, it was important to have all four quadrants visible. In this case, all the coordinates are positive. There is no need to show the negative values of x or y . Also, there are no x values bigger than about 3.14, and 1.75 is the largest value of y . We will therefore show these points on the following scale $0 \leq x \leq 3.5$ and $0 \leq y \leq 2$. The points are plotted to the right.

Here are some important points to note about this graph.


- The tick marks on the axes do not correspond to unit increments (i.e. the numbers do not go up by one).
- The scale on the x -axis is different than the scale on the y -axis.
- The scale is **chosen** to maximize the clarity of the plotted points.

Lesson Summary



- The **coordinate plane** is a two-dimensional space defined by a horizontal number line (the x -**axis**) and a vertical number line (the y -**axis**). The **origin** is the point where these two lines meet. Four areas, or **quadrants**, are formed as shown in the diagram at right.
- Each point on the coordinate plane has a set of **coordinates**, two numbers written as an **ordered pair** which describe how far along the x -axis and y -axis the point is. The x -**coordinate** is always written first, then the y -**coordinate**. Here is an example (x, y) .

Review Questions




1. Identify the coordinates of each point, $A - F$, on the graph to the right.
2. Plot the following points on a graph and identify which quadrant each point lies in:
 - (a) $(4, 2)$
 - (b) $(-3, 5.5)$
 - (c) $(4, -4)$
 - (d) $(-2, -3)$
3. The following three points are three vertices of square $ABCD$. Plot them on a graph then determine what the coordinates of the fourth point, D , would be. Plot that point and label it.


$A (-4, -4)$
 $B (3, -4)$
 $C (3, 3)$
4. Becky has a large bag of M&Ms that she knows she should share with Jaeyun. Jaeyun has a packet of Starburst. Becky tells Jaeyun that for every Starburst he gives her, she will give him three M&Ms in return. If x is the number of Starburst that Jaeyun gives Becky, and y is the number of M&Ms he gets in return then complete each of the following.
 - (a) Write an algebraic rule for y in terms of x
 - (b) Make a table of values for y with x values of $0, 1, 2, 3, 4, 5$.
 - (c) Plot the function linking x and y on the following scale $0 \leq x \leq 10, 0 \leq y \leq 10$.

Review Answers

1. $A(5, 6)B(-5, 5)C(-2, 3)D(-2, -2)E(3, -4)F(2, -6)$




- (a) Quadrant I
- (b) Quadrant II
- (c) Quadrant IV
- (d) Quadrant III



4. (a) $y = 3x$
 (b)

x	y
0	0
1	3
2	6
3	9
4	12
5	15



3.2 Graphs of Linear Equations

Learning Objectives

- Graph a linear function using an equation.
- Write equations and graph horizontal and vertical lines.
- Analyze graphs of linear functions and read conversion graphs.

Graph a Linear Equation

At the end of Lesson 4.1 we looked at graphing a function from a rule. A rule is a way of writing the relationship between the two quantities we are graphing. In mathematics, we tend to use the words **formula** and **equation** to describe what we get when we express relationships algebraically. Interpreting and graphing these equations is an important skill that you will use frequently in math.


Example 1

A taxi fare costs more the further you travel. Taxis usually charge a fee on top of the per-mile charge to cover hire of the vehicle. In this case, the taxi charges \$3 as a set fee and \$0.80 per mile traveled. Here is the equation linking the cost in dollars (y) to hire a taxi and the distance traveled in miles (x).

$$y = 0.8x + 3$$

Graph the equation and use your graph to estimate the cost of a seven mile taxi ride.

We will start by making a table of values. We will take a few values for x 0, 1, 2, 3, 4, find the corresponding y values and then plot them. Since the question asks us to find the cost for a seven mile journey, we will choose a scale that will accommodate this.



x	y
0	3
1	3.8
2	4.6
3	5.4
4	6.2

The graph is shown to the right. To find the cost of a seven mile journey we first locate $x = 7$ on the horizontal axis and draw a line up to our graph. Next we draw a horizontal line across to the y axis and read where it hits. It appears to hit around half way between $y = 8$ and $y = 9$. Let's say it is 8.5.

Solution

A seven mile taxi ride would cost approximately \$8.50 (\$8.60 exactly).

There are a few interesting points that you should notice about this graph and the formula that generated it.

- The graph is a straight line (this means that the equation is **linear**), although the function is **discrete** and will graph as a series of points.
- The graph crosses the y -axis at $y = 3$ (look at the equation you will see a $+3$ in there!). This is the base cost of the taxi.
- Every time we move **over** by one square we move **up** by 0.8 squares (look at the coefficient of x in the equation). This is the rate of charge of the taxi (cost per mile).
- If we move over by three squares, we move up by 3×0.8 squares.

Example 2


A small business has a debt of \$500000 incurred from start-up costs. It predicts that it can pay off the debt at a rate of \$85000 per year according to the following equation governing years in business (x) and debt measured in thousands of dollars(y).

$$y = -85x + 500$$

Graph the above equation and use your graph to predict when the debt will be fully paid.

First, we start with our table of values. We plug in x -values and calculate our corresponding y -values.

x	y
0	500
1	415
2	330
3	245
4	160



Then we plot our points and draw the line that goes through them.

Take note of the scale that has been chosen. There is no need to have any points above $y = 500$, but it is still wise to allow a little extra.

We need to determine how many years (the x value) that it takes the debt (y value) to reach zero. We know that it is greater than four (since at $x = 4$ the y value is still positive), so we need an x scale that goes well past $x = 4$. In this case the x value runs from 0 to 12, though there are plenty of other choices that would work well.

To read the time that the debt is paid off, we simply read the point where the line hits $y = 0$ (the x axis). It looks as if the line hits pretty close to $x = 6$. So the debt will definitely be paid off in six years.

Solution

The debt will be paid off in six years.

Graphs and Equations of Horizontal and Vertical Lines


Example 3

"Mad-cabs" have an unusual offer going on. They are charging \$7.50 for a taxi ride of any length within the city limits. Graph the function that relates the cost of hiring the taxi (y) to the length of the journey in miles (x).

To proceed, the first thing we need is an **equation**. You can see from the problem that the cost of a journey does not depend on the length of the journey. It should come as no surprise that the equation then, does not have x in it. In fact, any value of x results in the same value of y (7.5). Here is the equation.

$$y = 7.5$$

The graph of this function is shown to the right. You can see that the graph $y = 7.5$ is simply a horizontal line.




Any time you see an equation of the form $y = \text{constant}$ then the graph is a horizontal line that intercepts the y -axis at the value of the constant.

Similarly, when you see an equation of the form $x = \text{constant}$ then the graph is a vertical line that intercepts the x -axis at the value of the constant. Notice that this a relation, and not a function because each x value (theres only one in this case) corresponds to many (actually an infinite number) y values.

Example 4

Plot the following graphs.

- (a) $y = 4$
- (b) $y = -4$
- (c) $x = 4$
- (d) $x = -4$



- (a) $y = 4$ is a horizontal line that crosses the y -axis at 4
 (b) $y = -4$ is a horizontal line that crosses the y -axis at -4
 (c) $x = 4$ is a vertical line that crosses the x -axis at 4
 (d) $x = -4$ is a vertical line that crosses the x -axis at -4

Example 5

Find an equation for the x -axis and the y -axis.

Look at the axes on any of the graphs from previous examples. We have already said that they intersect at the origin (the point where $x = 0$ and $y = 0$). The following definition could easily work for each axis.


x -axis: A horizontal line crossing the y -axis at zero.

y -axis: A vertical line crossing the x -axis at zero.


So using example 3 as our guide, we could define the x -axis as the line $y = 0$ and the y -axis as the line $x = 0$.

Analyze Graphs of Linear Functions

We often use line graphs to represent relationships between two linked quantities. It is a useful skill to be able to interpret the information that graphs convey. For example, the chart below shows a fluctuating stock price over ten weeks. You can read that the index closed the first week at about \$68, and at the end of the third week it was at about \$62. You may also see that in the first five weeks it lost about 20% of its value and that it made about 20% gain between weeks seven and ten. Notice that this relationship is discrete, although the dots are connected for ease of interpretation.



Analyzing line graphs is a part of life whether you are trying to decide to buy stock, figure out if your blog readership is increasing, or predict the temperature from a weather report. Many of these graphs are very complicated, so for now we will start off with some simple linear conversion graphs. Algebra starts with basic relationships and builds to the complicated tasks, like reading the graph above. In this section, we will look at reading information from simple linear conversion graphs.

**Example 6**


The graph shown at the right shows a chart for converting marked prices in a downtown store into prices that include sales tax. Use the graph to determine the cost inclusive of sales tax for a \$6.00 pen in the store.

To find the relevant price with tax we find the correct pre-tax price on the x -axis. This is the point $x = 6$.

Draw the line $x = 6$ up until it meets the function, then draw a horizontal line to the y -axis. This line hits at $y \approx 6.75$ (about three fourths of the way from $y = 6$ to $y = 7$).

Solution

The approximate cost including tax is \$6.75



Example 7

The chart for converting temperature from Fahrenheit to Celsius is shown to the right. Use the graph to convert the following:

1. 70° Fahrenheit to Celsius
2. 0° Fahrenheit to Celsius
3. 30° Celsius to Fahrenheit
4. 0° Celsius to Fahrenheit

1. To find 70° Fahrenheit we look along the Fahrenheit-axis (in other words the x -axis) and draw the line $x = 70$ up to the function. We then draw a horizontal line to the Celsius-axis (y -axis). The horizontal line hits the axis at a little over 20 (21 or 22).

Solution

70° Fahrenheit is approximately equivalent to 21° Celsius

2. To find 0° Fahrenheit, we are actually looking at the y -axis. Don't forget that this axis is simply the line $x = 0$. We just look to see where the line hits the y -axis. It hits just below the half way point between -15 and -20 .

Solution: 0° Fahrenheit is approximately equivalent to -18° Celsius .

3. To find 30° Celsius, we look up the Celsius-axis and draw the line $y = 30$ along to the function. When this horizontal line hits the function, draw a line straight down to the Fahrenheit-axis. The line hits the axis at approximately 85.

Solution

30° Celsius is approximately equivalent to 85° Fahrenheit.

4. To find 0° Celsius we are looking at the Fahrenheit-axis (the line $y = 0$). We just look to see where the function hits the x -axis. It hits just right of 30.

Solution


0° Celsius is equivalent to 32° Fahrenheit.

Lesson Summary


- Equations with the variables y and x can be graphed by making a chart of values that fit the equation and then plotting the values on a coordinate plane. This graph is simply another representation of the equation and can be analyzed to solve problems.
- Horizontal lines are defined by the equation $y = \text{constant}$ and vertical lines are defined by the equation $x = \text{constant}$.
- Be aware that although we graph the function as a line to make it easier to interpret, the function may actually be discrete.

Review Questions

- Make a table of values for the following equations and then graph them.
 - $y = 2x + 7$
 - $y = 0.7x - 4$
 - $y = 6 - 1.25x$
- "Think of a number. Triple it, and then subtract seven from your answer"**. Make a table of values and plot the function that represents this sentence.




- Write the equations for the five (A through E) lines plotted in the graph to the right.
- At the Airport, you can change your money from dollars into Euros. The service costs \$5, and for every additional dollar you get 0.7 Euros. Make a table for this and plot the function on a graph. Use your graph to determine how many Euros you would get if you give the office \$50.
- The graph to below shows a conversion chart for converting between weight in kilograms to weight in pounds. Use it to convert the following measurements.




- 4 kilograms into weight in pounds
- 9 kilograms into weight in pounds
- 12 pounds into weight in kilograms
- 17 pounds into weight in kilograms

Review Answers




1.

2. $y = 3x - 7$



- $Ay = 5By = -2Cy = -7Dx = -4Ex = 6$
- $y = 0.7(x - 5)$




5.
 - (a) 9 lb
 - (b) 20 lb
 - (c) 5.5 kg
 - (d) 7.75 kg

3.3 Graphing Using Intercepts


Learning Objectives

- Find intercepts of the graph of an equation.
- Use intercepts to graph an equation.
- Solve real-world problems using intercepts of a graph.


Introduction



Only two distinct points are needed to uniquely define a graph of a line. After all, there are an infinite number of lines that pass through a single point (a few are shown in the graph at right). But if you supplied just one more point, there can only be one line that passes through both points. To plot the line, just plot the two points and use a ruler, edge placed on both points, to trace the graph of the line.



There are a lot of options for choosing which two points on the line you use to plot it. In this lesson, we will focus on two points that are rather convenient for graphing: the points where our line crosses the x and y axes, or **intercepts**. We will be finding intercepts algebraically and using them to quickly plot graphs. Similarly, the x -intercept occurs at the point where the graph crosses the x -axis. The x -value in the graph at the right is 6.



Look at the graph to the right. The **y-intercept** occurs at the point where the graph crosses the y-axis. The y-value at this point is 8.

Similarly the **x-intercept** occurs at the point where the graph crosses the x-axis. The x-value at this point is 6.

Now we know that the x value of all the points on the y-axis is zero, and the y value of all the points on the x-axis is also zero. So if we were given the coordinates of the two intercepts (0, 8) and (6, 0) we could quickly plot these points and join them with a line to recreate our graph.

Note: Not all lines will have both intercepts but most do. Specifically, horizontal lines never cross the x-axis and vertical lines never cross the y-axis. For examples of these special case lines, see the graph at right.

Finding Intercepts by Substitution


Example 1

Find the intercepts of the line $y = 13 - x$ and use them to graph the function.

The first intercept is easy to find. The y-intercept occurs when $x = 0$ Substituting gives:

$$y = 13 - 0 = 13$$

(0, 13) is the y-intercept.



We know that the x-intercept has, by definition, a y-value of zero. Finding the corresponding x-value is a simple case of substitution:

$$0 = 13 - x$$

$$-13 = -x$$

To isolate x subtract 13 from both sides.
Divide by -1 .

Solution

$(13, 0)$ is the x -intercept.

To draw the graph simply plot these points and join them with a line.

Example 2


Graph the following functions by finding intercepts.

a. $y = 2x + 3$

b. $y = 7 - 2x$

c. $4x - 2y = 8$

d. $2x + 3y = -6$



a. Find the y -intercept by plugging in $x = 0$.

$$y = 2 \cdot 0 + 3 = 3$$

The y -intercept is $(0, 3)$

Find the x -intercept by plugging in $y = 0$.

$$0 = 2x + 3$$


$$-3 = 2x$$

$$-\frac{3}{2} = x$$

Subtract 3 from both sides.

Divide by 2.

The x -intercept is $(-1.5, 0)$.



b. Find the y -intercept by plugging in $x = 0$.

$$y = 7 - 2 \cdot 0 = 7$$

The y -intercept is $(0, 7)$.

Find the x -intercept by plugging in $y = 0$.

$$0 = 7 - 2x$$


$$-7 = -2x$$

$$\frac{7}{2} = x$$

Subtract 7 from both sides.

Divide by -2 .

The x -intercept is $(3.5, 0)$.



c. Find the y -intercept by plugging in $x = 0$.

$$4 \cdot 0 - 2y = 8$$

$$-2y = 8$$

$$y = -4$$

Divide by -2 .

The y -intercept is $(0, -4)$.

Find the x -intercept by plugging in $y = 0$.


$$4x - 2 \cdot 0 = 8$$

$$4x = 8$$

$$x = 2$$

Divide by 4.

The x -intercept is $(2, 0)$.



d. Find the y -intercept by plugging in $x = 0$.

$$2 \cdot 0 + 3y = -6$$

$$3y = -6$$

$$y = -2$$

Divide by 3.

The y -intercept is $(0, -2)$.

Find the x -intercept by plugging in $y = 0$.

$$2x + 3 \cdot 0 = -6$$

$$2x = -6$$

$$x = -3$$

Divide by 2.

The x -intercept is $(-3, 0)$

Finding Intercepts for Standard Form Equations Using the Cover-Up Method

Look at the last two equations in Example 2. These equations are written in **standard form**. Standard form equations are always written "**positive coefficient** times x plus (or minus) **positive coefficient** times y equals **value**". Note that the x term *always* has a positive value in front of it while the y value may have a negative term. The equation looks like this:

$$ax + by + c \text{ or } ax - by = c$$

(a and b are positive numbers)

There is a neat method for finding intercepts in standard form, often referred to as the cover-up method.

Example 3

Find the intercepts of the following equations.

a. $7x - 3y = 21$

b. $12x - 10y = -15$

c. $x + 3y = 6$

To solve for each intercept, we realize that on the intercepts the value of **either** x or y is zero, and so any terms that contain the zero variable effectively disappear. To make a term disappear, simply cover it (a finger is an excellent way to cover up terms) and solve the resulting equation.

a. To solve for the y -intercept we set $x = 0$ and cover up the x term:

$$\text{Cover } x \text{ term} \quad -3y = 21$$

$$\begin{aligned} -3y &= 21 \\ y &= -7 \end{aligned}$$

$(0, -7)$ is the y – intercept

Now we solve for the x –intercept:

$$\text{Cover } y \text{ term} \quad 7x = 21$$

$$\begin{aligned} 7x &= 21 \\ x &= 3 \end{aligned}$$

$(3, 0)$ is the x – intercept.

b. Solve for the y –intercept ($x = 0$) by covering up the x term.

$$\text{Cover } x \text{ term} \quad -10y = -15$$

$$\begin{aligned} -10y &= -15 \\ y &= -1.5 \end{aligned}$$

$(0, -1.5)$ is the y – intercept.

Solve for the x –intercept ($y = 0$):

$$\text{Cover } y \text{ term} \quad 12x = -15$$

$$\begin{aligned} 12x &= -15 \\ x &= -\frac{5}{4} \end{aligned}$$


$(-1.25, 0)$ is the x – intercept.

c. Solve for the y –intercept ($x = 0$) by covering up the x term:

$$\text{Cover } x \text{ term} \quad 3y = 6$$

$$\begin{aligned} 3y &= 6 \\ y &= 2 \end{aligned}$$

$(0, 2)$ is the y – intercept.



Solve for the y -intercept:

$$x \cdot (-1) = 6$$

$$x = 6$$

$(6, 0)$ is the x -intercept.

The graph of these functions and the intercepts is shown in the graph on the right.

Solving Real-World Problems Using Intercepts of a Graph

Example 4


The monthly membership cost of a gym is \$25 per month. To attract members, the gym is offering a \$100 cash rebate if members sign up for a full year. Plot the cost of gym membership over a 12 month period. Use the graph to determine the final cost for a 12 month membership.

Let us examine the problem. Clearly the cost is a function of the number of months (not the other way around). Our independent variable is the number of months (the domain will be whole numbers) and this will be our x value. The cost in dollars is the dependent variable and will be our y value. Every month that passes the money paid to the gym goes up by \$25. However, we start with a \$100 cash gift, so our **initial cost** (y -intercept) is \$100. This pays for four months ($4 \times \$25 = 100$) so after four months the cost of membership (y -value) is zero.

The y -intercept is $(0, -100)$. The x -intercept is $(4, 0)$.

We plot our points, join them with a straight line and extend that line out all the way to the $x = 12$ line. The graph is shown below.

Cost of Gym Membership by Number of Months



To find the cost of a 12 month membership we simply read off the value of the function at the 12 month point. A line drawn up from $x = 12$ on the x axis meets the function at a y value of \$200.

Solution

The cost of joining the gym for one year is \$200.

Example 5

Jesus has \$30 to spend on food for a class barbecue. Hot dogs cost \$0.75 each (including the bun) and burgers cost \$1.25 (including bun and salad). Plot a graph that shows all the combinations of hot dogs and burgers he could buy for the barbecue, without spending more than \$30.

This time we will find an equation first, and then we can think logically about finding the intercepts.

If the number of burgers that John buys is x , then the money spent on burgers is $1.25x$.

If the number of hot dogs he buys is y then the money spent on hot dogs is $0.75y$.

$$1.25x + 0.75y$$

The total cost of the food.

The total amount of money he has to spend is \$30. If he is to spend it ALL, then we can use the following equation.

$$1.25x + 0.75y = 30$$

We solve for the intercepts using the cover-up method.

First the y -intercept ($x = 0$).

$$0.75y = 30$$

$$0.75y = 30$$

$$y = 40$$

y - intercept(0,40)

Then the x -intercept ($y = 0$)


$$1.25x + 0 = 30$$

$$1.25x = 30$$

$$x = 24$$

x - intercept(24,0)

Possible Numbers of Hot Dogs and Hamburgers Purchased for \$30



We can now plot the points and join them to create our graph, shown right.

Here is an alternative to the equation method.

If Jesus were to spend ALL the money on hot dogs, he could buy $\frac{30}{0.75} = 40$ hot dogs. If on the other hand, he were to buy only burgers, he could buy $\frac{30}{1.25} = 24$ burgers. So you can see that we get two intercepts: (0 burgers, 40 hot dogs) and (24 burgers, 0 hot dogs). We would plot these in an identical manner and design our graph that way.

As a final note, we should realize that Jesus problem is really an example of an **inequality**. He can, in fact, spend any amount up to \$30. The only thing he cannot do is spend more than \$30. So our graph reflects this. The shaded region shows where Jesus' solutions all lie. We will see inequalities again in Chapter 6.

Lesson Summary

- A **y-intercept** occurs at the point where a graph crosses the y -axis ($x = 0$) and an **x-intercept** occurs at the point where a graph crosses the x -axis ($y = 0$).
- The y -intercept can be found by **substituting** $x = 0$ into the equation and solving for y . Likewise, the x -intercept can be found by **substituting** $y = 0$ into the equation and solving for x .
- A linear equation is in **standard form** if it is written as positive coefficient times x plus (or minus) positive coefficient times y equals value. Equations in standard form can be solved for the intercepts by covering up the x (or y) term and solving the equation that remains.




Review Questions

1. Find the intercepts for the following equations using substitution.
 - (a) $y = 3x - 6$
 - (b) $y = -2x + 4$
 - (c) $y = 14x - 21$
 - (d) $y = 7 - 3x$
2. Find the intercepts of the following equations using the cover-up method.
 - (a) $5x - 6y = 15$
 - (b) $3x - 4y = -5$
 - (c) $2x + 7y = -11$
 - (d) $5x + 10y = 25$
3. Use any method to find the intercepts and then graph the following equations.
 - (a) $y = 2x + 3$
 - (b) $6(x - 1) = 2(y + 3)$
 - (c) $x - y = 5$
 - (d) $x + y = 8$
4. At the local grocery store strawberries cost \$3.00 per pound and bananas cost \$1.00 per pound. If I have \$10 to spend between strawberries and bananas, draw a graph to show what combinations of each I can buy and spend exactly \$10.
5. A movie theater charges \$7.50 for adult tickets and \$4.50 for children. If the theater takes \$900 in ticket sales for a particular screening, draw a graph which depicts the possibilities for the number of adult tickets and the number of child tickets sold.
6. Why can't we use the intercept method to graph the following equation?
 $3(x + 2) = 2(y + 3)$

Review Answers

1. (a) $(0, -6), (2, 0)$
(b) $(0, 4), (2, 0)$

- (c) $(0, -21), (1.5, 0)$
- (d) $(0, 7), (\frac{7}{3}, 0)$
- 2. (a) $(0, -2.5), (3, 0)$
- (b) $(0, 1.25), (-\frac{5}{3}, 0)$
- (c) $(0, -\frac{11}{7}), (-\frac{11}{2}, 0)$
- (d) $(0, 2.5), (5, 0)$



6. This equation reduces to $3x = 2y$, which passes through $(0, 0)$ and therefore only has **one intercept**. Two intercepts are needed for this method to work.


3.4 Slope and Rate of Change

Learning Objectives

- Find positive and negative slopes.
- Recognize and find slopes for horizontal and vertical lines.
- Understand rates of change.
- Interpret graphs and compare rates of change.

Introduction

We come across many examples of slope in everyday life. For example, a slope is in the pitch of a roof, the grade or incline of a road, and the slant of a ladder leaning on a wall. In math, we use the word **slope** to define steepness in a particular way.



$$\text{Slope} = \frac{\text{distance moved vertically}}{\text{distance moved horizontally}}$$

This is often reworded to be easier to remember:


$$\text{Slope} = \frac{\text{rise}}{\text{run}}$$

Essentially, slope is the change in y if x increases by 1.

In the picture to the right, the slope would be the ratio of the **height** of the hill (the **rise**) to the horizontal **length** of the hill (the **run**).

$$\text{Slope} = \frac{3}{4} = 0.75$$

If the car were driving to the **right** it would **climb** the hill. We say this is a positive slope. Anytime you see the graph of a line that goes up as you move to the right, the slope is **positive**.



If the car were to keep driving after it reached the top of the hill, it may come down again. If the car is driving to the **right** and **descending**, then we would say that the slope is **negative**. The picture at right has a **negative slope** of -0.75 .

Do not get confused! If the car turns around and drives back down the hill shown, we would still classify the slope as positive. This is because the rise would be -3 , but the run would be -4 (think of the x -axis if you move from right to left you are moving in the negative x -direction). Our ratio for moving **left** is:

$$\text{Slope} = \frac{-3}{-4} = 0.75 \qquad \text{A negative divided by a negative is a positive.}$$

So as we move from left to right, positive slopes increase while negative slopes decrease.

Find a Positive Slope

We have seen that a function with a positive slope increases in y as we increase x . A simple way to find a value for the slope is to draw a right angled triangle whose hypotenuse runs along the line. It is then a simple matter of measuring the distances on the triangle that correspond to the rise (the vertical dimension) and the run (the horizontal dimension).

Example 1

Find the slopes for the three graphs shown right.

There are already right-triangles drawn for each of the lines. In practice, you would have to do this yourself. Note that it is easiest to make triangles whose vertices are **lattice points** (i.e. the coordinates are all integers).

a. The rise shown in this triangle is 4 units, the run is 2 units.

$$\text{Slope} = \frac{4}{2} = 2$$

b. The rise shown in this triangle is 4 units, the run is also 4 units.


$$\text{Slope} = \frac{4}{4} = 1$$

c. The rise shown in this triangle is 2 units, the run is 4 units.

$$\text{Slope} = \frac{2}{4} = \frac{1}{2}$$

Example 2

Find the slope of the line that passes through the points (1, 2) and (4, 7).



We already know how to graph a line if we are given two points. We simply plot the points and connect them with a line. Look at the graph shown at right.

Since we already have coordinates for our right triangle, we can quickly work out that the rise would be 5 and the run would be 3 (see diagram). Here is our slope.

$$\text{Slope} = \frac{7 - 2}{4 - 1} = \frac{5}{3}$$

If you look closely at the calculations for the slope you will notice that the 7 and 2 are the y -coordinates of the two points and the 4 and 1 are the x -coordinates. This suggests a pattern we can follow to get a general formula for the slope between two points (x_1, y_1) and (x_2, y_2) .

Slope between (x_1, y_1) and $(x_2, y_2) = \frac{y_2 - y_1}{x_2 - x_1}$ or $m = \frac{\Delta y}{\Delta x}$


In the second equation, the letter m denotes the slope (you will see this a lot in this chapter) and the Greek letter delta (Δ) means **change**. So another way to express slope is *change in y* divided by *change in x* . In the next section, you will see that it does not matter which point you choose as point 1 and which you choose as point 2.

Find a Negative Slope

Any function with a negative slope is simply a function that decreases as we increase x . If you think of the function as the incline of a road a negative slope is a road that goes **downhill** as you drive to the **right**.

Example 3

Find the slopes of the lines on the graph to the right.



Look at the lines. Both functions fall (or decrease) as we move from left to right. Both of these lines have a **negative slope**.

Neither line passes through a great number of lattice points, but by looking carefully you can see a few points that look to have integer coordinates. These points have been identified (with rings) and we will use these to determine the slope. We will also do our calculations twice, to show that we get the same slope whichever way we choose point 1 and point 2.

For line A:

$$(x_1, y_1) = (-6, 3) \quad (x_2, y_2) = (5, -1)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{(-1) - (3)}{(5) - (-6)} = \frac{-4}{11} \approx -0.364$$

$$(x_1, y_1) = (5, -1) \quad (x_2, y_2) = (-6, 3)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{(3) - (-1)}{(-6) - (5)} = \frac{-4}{-11} \approx -0.364$$

For line B:

$$(x_1, y_1) = (-4, 6) \quad (x_2, y_2) = (4, -5)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{(-5) - (6)}{(4) - (-4)} = \frac{-11}{8} = -1.375$$

$$(x_1, y_1) = (4, -5) \quad (x_2, y_2) = (-4, 6)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{(6) - (-5)}{(-4) - (4)} = \frac{11}{-8} = -1.375$$


You can see that whichever way you select the points, the answers are the same!

Solution

Line A has slope -0.364 . Line B has slope -1.375 .

Find the Slopes of Horizontal and Vertical lines

Example 4



Determine the slopes of the two lines on the graph at the right.

There are two lines on the graph. A ($y = 3$) and B ($x = 5$).

Let's pick two points on line A . say, $(x_1, y_1) = (-4, 3)$ and $(x_2, y_2) = (5, 3)$ and use our equation for slope.

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{(3) - (3)}{(5) - (-4)} = \frac{0}{9} = 0$$

If you think about it, this makes sense. If there is no change in y as we increase x then there is no slope, or to be correct, a slope of zero. You can see that this must be true for all horizontal lines.

Horizontal lines ($y = \text{constant}$) all have a slope of 0.

Now consider line B . Pick two distinct points on this line and plug them in to the slope equation.

$(x_1, y_1) = (5, -3)$ and $(x_2, y_2) = (5, 4)$.

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{(4) - (-3)}{(5) - (5)} = \frac{7}{0} \quad \text{A division by zero!}$$

Divisions by zero lead to infinities. In math we often use the term **undefined** for any division by zero.

Vertical lines ($x = \text{constant}$) all have an infinite (or undefined) slope.

Find a Rate of Change

The slope of a function that describes real, measurable quantities is often called a **rate of change**. In that case, the slope refers to a change in one quantity (y) **per** unit change in another quantity (x).

Example 5

*Andrea has a part time job at the local grocery store. She saves for her vacation at a rate of \$15 every week. Express this rate as money saved **per day** and money saved **per year**.*

Converting rates of change is fairly straight forward so long as you remember the equations for rate (i.e. the equations for slope) and know the conversions. In this case 1 week = 7 days and 52 weeks = 1 year.

$$\begin{aligned} \text{rate} &= \frac{\$15}{1 \text{ week}} \cdot \frac{1 \text{ week}}{7 \text{ days}} = \frac{\$15}{7 \text{ days}} = \frac{15}{7} \text{ dollars per day} \approx \$2.14 \text{ per day} \\ \text{rate} &= \frac{\$15}{1 \text{ week}} \cdot \frac{52 \text{ week}}{1 \text{ year}} = \$15 \cdot \frac{52}{\text{year}} = \$780 \text{ per year} \end{aligned}$$


Example 6

A candle has a starting length of 10 inches. Thirty minutes after lighting it, the length is 7 inches. Determine the rate of change in length of the candle as it burns. Determine how long the candle takes to completely burn to nothing.

In this case, we will graph the function to visualize what is happening.

We have two points to start with. We know that at the moment the candle is lit (time = 0) the length of the candle is 10 inches. After thirty minutes (time = 30) the length is 7 inches. Since the candle length is a function of time we will plot time on the horizontal axis, and candle length on the vertical axis. Here is a graph showing this information.

Candle Length by Burning Time



The rate of change of the candle is simply the slope. Since we have our two points $(x_1, y_1) = (0, 10)$ and $(x_2, y_2) = (30, 7)$ we can move straight to the formula.

$$\text{Rate of change} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{(7 \text{ inches}) - (10 \text{ inches})}{(30 \text{ minutes}) - (0 \text{ minutes})} = \frac{-3 \text{ inches}}{30 \text{ minutes}} = -0.1 \text{ inches per minute}$$

The slope is negative. A negative rate of change means that the quantity is decreasing with time.

We can also convert our rate to inches per hour.

$$\text{rate} = \frac{-0.1 \text{ inches}}{1 \text{ minute}} \cdot \frac{60 \text{ minutes}}{1 \text{ hour}} = \frac{-6 \text{ inches}}{1 \text{ hour}} = -6 \text{ inches per hour}$$

To find the point when the candle reaches zero length we can simply read off the graph (100 minutes). We can use the rate equation to verify this algebraically.

$$\text{Length burned} = \text{rate} \times \text{time}$$

$$0.1 \times 100 = 10$$


Since the candle length was originally 10 inches this confirms that 100 minutes is the correct amount of time.

Interpret a Graph to Compare Rates of Change

Example 7

Examine the graph below. It represents a journey made by a large delivery truck on a particular day. During the day, the truck made two deliveries, each one taking one hour. The driver also took a one hour break for lunch. Identify what is happening at each stage of the journey (stages A through E)

Trucks Distance from Home by Time



Here is the driver’s journey.

- A. The truck sets off and travels 80 miles in 2 hours .
- B. The truck covers no distance for 1 hours .
- C. The truck covers $(120 - 80) = 40$ miles in 1 hours

D. the truck covers no distance for 2 hours .

E. The truck covers 120 miles in 2 hours .

Lets look at the rates of change for each section.

A. Rate of change = $\frac{\Delta y}{\Delta x} = \frac{80 \text{ miles}}{2 \text{ hours}} = 40 \text{ miles per hour}$

- The rate of change is a **velocity!** This is a very important concept and one that deserves a special note!

The slope (or rate of change) of a distance-time graph is a velocity.

You may be more familiar with calling **miles per hour** a **speed**. **Speed** is the **magnitude** of a **velocity**, or, put another way, velocity has a direction, speed does not. This is best illustrated by working through the example.

On the first part of the journey sees the truck travel at a constant velocity of 40 mph for 2 hours covering a distance of 80 miles .

B. **Slope** = 0 so **rate of change** = 0 mph. The truck is stationary for one hour. This could be a lunch break, but as it is only 2 hours since the truck set off it is likely to be the first delivery stop.

C. Rate of change = $\frac{\Delta y}{\Delta x} = \frac{(120-80) \text{ miles}}{(4-3) \text{ hours}} = 40 \text{ miles per hour}$. The truck is traveling at 40 mph.

D. **Slope** = 0 so **rate of change** = 0 mph . The truck is stationary for two hours. It is likely that the driver used these 2 hours for a lunch break plus the second delivery stop. At this point the truck is 120 miles from the start position.

E. Rate of change = $\frac{\Delta y}{\Delta x} = \frac{(0-120) \text{ miles}}{(8-6) \text{ hours}} = \frac{-120 \text{ miles}}{2 \text{ hours}} = -60 \text{ miles per hour}$. The truck is traveling at **negative** 60 mph .

Wait, a negative velocity? Does this mean that the truck is reversing? Well, probably not. What it means is that the distance (and dont forget that is the distance measured from the starting position) is decreasing with time. The truck is simply driving in the opposite direction. In this case, back to where it started from. So, the speed of the truck would be 60 mph, but the velocity (which includes direction) is negative because the truck is getting closer to where it started from. The fact that it no longer has two heavy loads means that it travels faster (60 mph as opposed to 40 mph) covering the 120 mile return trip in 2 hours .




Lesson Summary

- **Slope** is a measure of change in the vertical direction for each step in the horizontal direction. Slope is often represented as "m".
- $Slope = \frac{\text{rise}}{\text{run}}$ or $m = \frac{\Delta y}{\Delta x}$
- The slope between two points (x_1, y_1) and $(x_2, y_2) = \frac{y_2 - y_1}{x_2 - x_1}$
- **Horizontal lines** ($y = \text{constant}$) all have a slope of 0.
- **Vertical lines** ($x = \text{constant}$) all have an infinite (or undefined) slope.
- The slope (or **rate of change**) of a distance-time graph is a **velocity**.

Review Questions


1. Use the slope formula to find the slope of the line that passes through each pair of points.
 - (a) $(-5, 7)$ and $(0, 0)$
 - (b) $(-3, -5)$ and $(3, 11)$
 - (c) $(3, -5)$ and $(-2, 9)$
 - (d) $(-5, 7)$ and $(-5, 11)$
 - (e) $(9, 9)$ and $(-9, -9)$
 - (f) $(3, 5)$ and $(-2, 7)$

2. Use the points indicated on each line of the graphs to determine the slopes of the following lines.



3. The graph below is a distance-time graph for Marks three and a half mile cycle ride to school. During this ride, he rode on cycle paths but the terrain was hilly. He rode slower up hills and faster down them. He stopped once at a traffic light and at one point he stopped to mend a tire puncture. Identify each section of the graph accordingly.

Andrews Distance from Home by Time



Review Answers

1. (a) -1.4
- (b) 2.67
- (c) -2.8
- (d) undefined

- (e) 1
 - (f) -0.4
- 2.
- (a) 3
 - (b) 0.5
 - (c) -2
 - (d) 1
 - (e) undefined
 - (f) $\frac{1}{3}$
- 3.
- (a) A. uphill
 - (b) B. stopped (traffic light)
 - (c) C. uphill
 - (d) D. downhill
 - (e) E. stopped (puncture)
 - (f) F. uphill

3.5 Graphs Using Slope-Intercept Form

Learning Objectives

- Identify the slope and y -intercept of equations and graphs.
- Graph an equation in slope-intercept form.
- Understand what happens when you change the slope or intercept of a line.
- Identify parallel lines from their equations.

Identify Slope and y -intercept

One of the most common ways of writing linear equations prior to graphing them is called **slope-intercept form**. We have actually seen several slope-intercept equations so far. They take the following form:

$y = mx + b$ where m is the slope and the point $(0, b)$ is the y -intercept.

We know that the y -intercept is the point at which the line passes through the y -axis. The slope is a measure of the steepness of the line. Hopefully, you can see that if we know **one point** on a line and the slope of that line, we know what the line is. Being able to quickly identify the y -intercept and slope will aid us in graphing linear functions.

Example 1

Identify the slope and y -intercept of the following equations.

a) $y = 3x + 2$

b) $y = 0.5x - 3$

c) $y = -7x$

d) $y = -4$

Solution

a)

$y = 3x + 2$ with $y = mx + b$

Comparing, we see that $m = 3$ and $b = 2$.

$y = 3x + 2$ has a **slope of 3** and a **y -intercept of $(0, 2)$**

b)

$y = 0.5x - 3$

has a **slope of 0.5** and a **y -intercept of $(0, -3)$** .

Note that the y -intercept is **negative**. The b term includes the sign of the operator in front of the number. Just remember that $y = 0.5x - 3$ is identical to $y = 0.5x + (-3)$ and is in the form $y = mx + b$.

c) **At first glance, this does not appear to fit the slope-intercept form. To illustrate how we deal with this, let us rewrite the equation.**

A small diagram showing a line on a coordinate plane. The equation $y = -7x + 0$ is written next to the line. An arrow points to the coefficient -7 with the label "slope". Another arrow points to the constant term 0 with the label "intercept".

. We now see that we get a slope of -7 and a y -intercept of $(0,0)$.

Note that the slope is negative. The $(0,0)$ intercept means that the line passes through origin.

d) Rewrite as $y = 0x - 4$, giving us a slope of 0 and an intercept of $(0, -4)$.

Remember:

- When $m < 0$ the slope is negative.

For example, $y = -3x + 2$ has a slope of -3 .

- When $b < 0$ the intercept is below the x axis.

For example, $y = 4x - 2$ has a y -intercept of $(0, -2)$.

- When $m = 0$ the slope is zero and we have a horizontal line.


For example, $y = 3$ can be written as $y = 0x + 3$.

- When $b = 0$ the graph passes through the origin.

For example, $y = 4x$ can be written as $y = 4x + 0$.

Example 2

Identify the slope and y -intercept of the lines on the graph shown to the right.



The intercepts have been marked, as have a number of lattice points that lines pass through.

a. The y -intercept is $(0,5)$. The line also passes through $(2,3)$.

$$\text{slope } m = \frac{\Delta y}{\Delta x} = \frac{-2}{2} = -1$$

b. The y -intercept is $(0,2)$. The line also passes through $(1,5)$.

$$\text{slope } m = \frac{\Delta y}{\Delta x} = \frac{3}{1} = 3$$

c. The y -intercept is $(0, -1)$. The line also passes through $(2, 3)$.

$$\text{slope } m = \frac{\Delta y}{\Delta x} = \frac{4}{2} = 2$$

d. The y -intercept is $(0, -3)$. The line also passes through $(4, -4)$.

$$\text{slope } m = \frac{\Delta y}{\Delta x} = \frac{-1}{4} = \frac{-1}{4} \text{ or } -0.25$$

Graph an Equation in Slope-Intercept Form

Once we know the slope and intercept of a line it is easy to graph it. Just remember what slope means. Let's look back at this example from Lesson 4.1.


Example 3

Ahiga is trying to work out a trick that his friend showed him. His friend started by asking him to think of a number. Then double it. Then add five to what he got. Ahiga has written down a rule to describe the first part of the trick. He is using the letter x to stand for the number he thought of and the letter y to represent the result of applying the rule. His rule is:

$$y = 2x + 5$$

Help him visualize what is going on by graphing the function that this rule describes.

In that example, we constructed the following table of values.



x	y
0	$2 \cdot 0 + 5 = 0 + 5 = 5$
1	$2 \cdot 1 + 5 = 2 + 5 = 7$
2	$2 \cdot 2 + 5 = 4 + 5 = 9$
3	$2 \cdot 3 + 5 = 6 + 5 = 11$


The first entry gave us our y intercept $(0, 5)$. The other points helped us graph the line.

We can now use our equation for slope, and two of the given points.

Slope between $(x_1, y_1) = (0, 5)$ and $(x_2, y_2) = (3, 11)$.

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{11 - 5}{3 - 0} = \frac{6}{3} = 2$$

Thus confirming that the slope, $m = 2$.



An easier way to graph this function is the slope-intercept method. We can now do this quickly, by identifying the intercept and the slope.

$$y = 2x + 5$$


slope = 2 y-intercept = 5

Look at the graph we drew, the line intersects the y -axis at 5, and every time we move to the right by one unit, we move up by two units.

So what about plotting a function with a negative slope? Just remember that a negative slope means the function decreases as we increase x .

Example 4


Graph the following function. $y = -3x + 5$



- Identify y -intercept $b = 5$
- Plot intercept $(0, 5)$
- Identify slope $m = -3$
- Draw a line through the intercept that has a slope of -3 .

To do this last part remember that slope = $\frac{\text{rise}}{\text{run}}$ so for every unit we move to the right the function increases by -3 (in other words, for every square we move right, the function comes **down** by 3).

Changing the Slope of a Line



Look at the graph on the right. It shows a number of lines with different slopes, but all with the same y -intercept $(0, 3)$.

You can see all the positive slopes increase as we move from left to right while all functions with negative slopes fall as we move from left to right.

Notice that the higher the value of the slope, the steeper the graph.

The graph of $y = 2x + 3$ appears as the mirror image of $y = -2x + 3$. The two slopes are equal but opposite.

Fractional Slopes and Rise Over Run

Look at the graph of $y = 0.5x + 3$. As we increase the x value by 1, the y value increases by 0.5. If we increase the x value by 2, then the y value increases by 1. In fact, if you express any slope as a fraction, you can determine how to plot the graph by looking at the numerator for the *rise* (keep any negative sign included in this term) and the denominator for the *run*.


Example 5

Find integer values for the *rise* and *run* of following slopes then graph lines with corresponding slopes.


- a. $m = 3$
- b. $m = -2$
- c. $m = 0.75$
- d. $m = -0.375$


Solution:

a.




b.






c.



d.



Changing the Intercept of a Line



When we take an equation (such as $y = 2x$) and change the y intercept (leaving the slope intact) we see the following pattern in the graph on the right.

Notice that changing the intercept simply translates the graph up or down. Take a point on the graph of $y = 2x$, such as $(1, 2)$. The corresponding point on $y = 2x + 3$ would be $(1, 4)$. Similarly the corresponding point on the $y = 2x - 3$ line would be $(1, -1)$.

Will These Lines Ever Cross?

To answer that question, let us take two of the equations $y = 2x$ and $y = 2x + 3$ and solve for values of x and y that satisfy both equations. This will give us the (x, y) coordinates of the point of intersection.

$$2x = 2x + 3$$

$$0 = 0 + 3$$

or

Subtract $2x$ from both sides.

$$0 = 3 \text{ This statement is FALSE!}$$

When we get a false statement like this, it means that there are **no** (x, y) values that satisfy both equations simultaneously. The lines will **never** cross, and so they **must be parallel**.

Identify Parallel Lines

In the previous section, when we changed the intercept but left the slope the same, the new line was parallel to the original line. This would be true whatever the slope of the original line, as changing the intercept on a $y = mx + b$ graph does nothing to the slope. This idea can be summed up as follows.

Any two lines with identical slopes are parallel.

Lesson Summary

- A common form of a line (linear equation) is **slope-intercept form**:

$y = mx + b$ where m is the slope and the point $(0, b)$ is the y -intercept


- Graphing a line in slope-intercept form is a matter of first plotting the y -intercept $(0, b)$, then plotting more points by moving a step to the right (adding 1 to x) and moving the value of the slope vertically (adding m to y) before plotting each subsequent point.
- Any two lines with identical slopes are **parallel**.

Review Questions


- Identify the slope and y -intercept for the following equations.


- $y = 2x + 5$
- $y = -0.2x + 7$
- $y = x$
- $y = 3.75$

- Identify the slope of the following lines.



- Identify the slope and y -intercept for the following functions.





4. Plot the following functions on a graph.


- (a) $y = 2x + 5$
- (b) $y = -0.2x + 7$
- (c) $y = -x$
- (d) $y = 3.75$

5. Which two of the following lines are parallel?

- (a) $y = 2x + 5$
- (b) $y = -0.2x + 7$
- (c) $y = -x$
- (d) $y = 3.75$
- (e) $y = -\frac{1}{5}x - 11$
- (f) $y = -5x + 5$
- (g) $y = -3x + 11$
- (h) $y = 3x + 3.5$

Review Answers

1. (a) $m = 2, (0, 5)$
 (b) $m = -0.2, (0, 7)$
 (c) $m = 1, (0, 0)$
 (d) $m = 0, (0, 3.75)$
2. (a) A. $m = -2$
 (b) B. $m = -\frac{4}{3}$
 (c) C. $m = 0$
 (d) D. $m = \frac{2}{5}$
 (e) E. $m = -0.25$
 (f) F. $m = -0.5$
 (g) G. $m = 4$
3. (a) A. $y = -\frac{2}{3}x + 1.5$
 (b) B: $y = 3x + 1$
 (c) C: $y = 0.5x - 2$
 (d) D: $y = -x$
 (e) E: $y = 3$
 (f) F: $y = -0.2x - 1$



5. b and e

3.6 Direct Variation Models

Learning Objectives

- Identify direct variation.
- Graph direct variation equations.
- Solve real-world problems using direct variation models.



Introduction

Suppose you see someone buy five pounds of strawberries at the grocery store. The clerk weighs the strawberries and charges \$12.50 for them. Now suppose you wanted two pounds of strawberries for yourself. How much would you expect to pay for them?

Identify Direct Variation

The preceding problem is an example of a **direct variation**. We would expect that the strawberries are priced on a "per pound" basis, and that if you buy two-fifths of the amount of strawberries, you would pay two-fifths of \$12.50 for your strawberries.

$$\frac{2}{5} \times \$12.50 = \$5.00$$

Similarly, if you bought 10 pounds of strawberries (twice the amount) you would pay $2 \times \$12.50$ and if you did not buy any strawberries you would pay nothing.

If variable y varies directly with variable x , then we write the relationship as:

$$y = k \cdot x$$

k is called the **constant of proportionality**.


If we were to graph this function you can see that it passes through the origin, because $y = 0$, when $x = 0$ whatever the value of k . So we know that a direct variation, when graphed, has a single intercept at $(0, 0)$.

Example 1

If y varies directly with x according to the relationship $y = k \cdot x$, and $y = 7.5$ when $x = 2.5$, determine the constant of proportionality, k .

We can solve for the constant of proportionality using substitution.

Substitute $x = 2.5$ and $y = 7.5$ into the equation $y = k \cdot x$



$$7.5 = k(2.5)$$

Divide both sides by 2.5.

$$\frac{7.5}{2.5} = k = 3$$

Solution

The constant of proportionality, $k = 3$.

We can graph the relationship quickly, using the intercept $(0,0)$ and the point $(2.5, 7.5)$. The graph is shown right. It is a straight line with a slope $= 3$.

The graph of a direct variation has a slope that is equal to the constant of proportionality, k .

Example 2



The volume of water in a fish-tank, V , varies directly with depth, d . If there are 15 gallons in the tank when the depth is eight inches, calculate how much water is in the tank when the depth is 20 inches.

This is a good example of a direct variation, but for this problem we will need to determine the equation of the variation ourselves. Since the volume, V , depends on depth, d , we will use the previous equation to create new one that is better suited to the content of the new problem.

$$y = k \cdot x$$

In place of y we will use V and in place of x we will use d .

$$V = k \cdot d$$

We know that when the depth is 8 inches, the volume is 15 gallons. Now we can substitute those values into our equation.

Substitute $V = 15$ and $x = 8$:

$$V = k \cdot d$$

$$15 = k(8)$$

Divide both sides by 8.

$$\frac{15}{8} = k = 1.875$$

Now to find the volume of water at the final depth we use $V = k \cdot d$ and substitute for our new d .


$$V = k \cdot d$$

$$V = 1.875 \times 20$$

$$V = 37.5$$

Solution

At a depth of 20 inches, the volume of water in the tank is 37.5 gallons.

Example 3

The graph shown to the right shows a conversion chart used to convert U.S. dollars (US\$) to British pounds (GB) in a bank on a particular day. Use the chart to determine the following.

- (i) The number of pounds you could buy for \$600.
- (ii) The number of dollars it would cost to buy 200.
- (iii) The exchange rate in pounds per dollar.
- (iv) Is the function continuous or discrete?

Solution

In order to solve (i) and (ii) we could simply read off the graph: it looks as if at $x = 600$ the graph is about one fifth of the way between 350 and 400. So \$600 would buy 360. Similarly, the line $y = 200$ would appear to intersect the graph about a third of the way between \$300 and \$400. We would probably round this to \$330. So it would cost approximately \$330 to buy 200.

To solve for the exchange rate we should note that as this is a direct variation, because the graph is a straight line passing through the origin. The slope of the line gives the constant of proportionality (in this case the **exchange rate**) and it is equal to the ratio of the y -value to x -value. Looking closely at the graph, it is clear that there is one lattice point that the line passes through (500, 300). This will give us the most accurate estimate for the slope (exchange rate).

$$y = k \cdot x \Rightarrow k = \frac{y}{x}$$

$$\text{rate} = \frac{300 \text{ pounds}}{500 \text{ dollars}} = 0.60 \text{ pounds per dollar}$$


Graph Direct Variation Equations

We know that all direct variation graphs pass through the origin, and also that the slope of the line is equal to the constant of proportionality, k . Graphing is a simple matter of using the point-slope or point-point methods discussed earlier in this chapter.

Example 4

Plot the following direct relations on the same graph.

- a. $y = 3x$
- b. $y = -2x$
- c. $y = -0.2x$
- d. $y = \frac{2}{9}x$



Solution

- a. The line passes through (0,0). All these functions will pass through this point. It is plotted in red. This function has a slope of 3. When we move across by one unit, the function increases by three units.
- b. The line has a slope of -2 . When we move across the graph by one unit the function **falls** by two units.
- c. The line has a slope of -0.2 . As a fraction this is equal to $-\frac{1}{5}$. When we move across by five units, the function **falls** by one unit.
- d. The line passes through (0,0) and has a slope of $\frac{2}{9}$. When we move across the graph by 9 units, the function increases by two units.

Solve Real-World Problems Using Direct Variation Models

Direct variations are seen everywhere in everyday life. Any time that we have one quantity that doubles when another related quantity doubles, we say that they follow a direct variation.

Newton’s Second Law

In 1687, Sir Isaac Newton published the famous *Principia Mathematica*. It contained, among other things, his Second Law of Motion. This law is often written as:

$$F = m \cdot a$$

A force of F (Newtons) applied to a mass of m (kilograms) results in acceleration of a (meterspersecond²).

Example 5

If a 175 Newton force causes a heavily loaded shopping cart to accelerate down the aisle with an acceleration of 2.5 m/s^2 , calculate

- (i) *The mass of the shopping cart.*
- (ii) *The force needed to accelerate the same cart at 6 m/s^2 .*

Solution

(i) This question is basically asking us to solve for the constant of proportionality. Let us compare the two formulas.

$y = k \cdot x$	The direct variation equation
$F = m \cdot a$	Newton’s Second law

We see that the two equations have the same form; y is analogous to force and x analogous to acceleration.

We can solve for m (the mass) by substituting our given values for force and acceleration:

Substitute $F = 175, a = 2.5$

$175 = m(2.5)$	Divide both sides by 2.5.
$70 = m$	

The mass of the shopping cart is 70 kg.

(ii) Once we have solved for the mass we simply substitute that value, plus our required acceleration back into the formula $F = m \cdot a$ and solve for F :

Substitute $m = 70$, $a = 6$

$$F = 70 \times 6 = 420$$

The force needed to accelerate the cart at 6 m/s^2 is 420 Newtons.

Ohm's Law

The electrical current, I (amps), passing through an electronic component varies directly with the applied voltage, V (volts), according to the relationship:

$$V = I \cdot R \quad \text{where } R \text{ is the resistance (measured in Ohms)}$$

The resistance is considered to be a constant for all values of V and I .


Example 6

A certain electronics component was found to pass a current of 1.3 amps at a voltage of 2.6 volts. When the voltage was increased to 12.0 volts the current was found to be 6.0 amps.

- Does the component obey Ohms law?*
- What would the current be at 6 volts?*

Solution

a) Ohm's law is a simple direct proportionality law. Since the resistance R is constant, it acts as our constant of proportionality. In order to know if the component obeys Ohm's law we need to know if it follows a direct proportionality rule. In other words **is V directly proportional to I ?**



Method One Graph It

If we plot our two points on a graph and join them with a line, does the line pass through $(0,0)$?

Point 1 = 2.6, $I = 1.3$ our point is $(1.3, 2.6)^*$

Point 2 $V = 12.0$, $I = 6.0$ our point is $(6, 12)$

Plotting the points and joining them gives the following graph.

The graph does appear to pass through the origin, so

Yes, the component obeys Ohms law.

Method Two Solve for

We can quickly determine the value of R in each case. It is the ratio of the voltage to the resistance.

$$\text{Case 1 } R = \frac{V}{I} = \frac{2.6}{1.3} = 2 \text{ Ohms}$$

$$\text{Case 2 } R = \frac{V}{I} = \frac{12}{6} = 2 \text{ Ohms}$$

The values for R agree! This means that the line that joins point 1 to the origin is the same as the line that joins point 2 to the origin. **The component obeys Ohms law.**

b) To find the current at 6 volts, simply substitute the values for V and R into $V = I \cdot R$

Substitute $V = 6$, $R = 2$

- In physics, it is customary to plot voltage on the horizontal axis as this is most often the independent variable. In that situation, the slope gives the **conductance**, σ . However, by plotting the current on the horizontal axis, the **slope** is equal to the **resistance**, R .

$$6 = I(2)$$

Divide both sides by 2.

$$3 = I$$

Solution

The current through the component at a voltage of 6 volts is 3 amps.

Lesson Summary

- If a variable y varies *directly* with variable x , then we write the relationship as


$$y = k \cdot x$$

Where k is a constant called the **constant of proportionality**.

- Direct variation** is very common in many areas of science.

Review Questions

- Plot the following direct variations on the same graph.
 - $y = \frac{4}{3}x$
 - $y = -\frac{2}{3}x$
 - $y = -\frac{1}{6}x$
 - $y = 1.75x$
- Dasans mom takes him to the video arcade for his birthday. In the first 10 minutes, he spends \$3.50 playing games. If his allowance for the day is \$20.00, how long can he keep playing games before his money is gone?
- The current standard for low-flow showerheads heads is 2.5 gallons per minute. Calculate how long it would take to fill a 30 gallon bathtub using such a showerhead to supply the water.
- Amen is using a hose to fill his new swimming pool for the first time. He starts the hose at 10 P.M. and leaves it running all night. At 6 AM he measures the depth and calculates that the pool is four sevenths full. At what time will his new pool be full?
- Land in Wisconsin is for sale to property investors. A 232 acre lot is listed for sale for \$200500. Assuming the same price per acre, how much would a 60 acre lot sell for?
- The force (F) needed to stretch a spring by a distance x is given by the equation $F = k \cdot x$, where k is the spring constant (measured in Newtons per centimeter, N/cm). If a 12 Newton force stretches a certain spring by 10 cm, calculate:
 - The spring constant, k
 - The force needed to stretch the spring by 7 cm .
 - The distance the spring would stretch with a 23 Newton force.



2. 57 minutes 8 seconds
3. 12 minutes
4. 12 : 00 Midday
5. \$51, 853
6. (a) $k = 1.2 \text{ N/cm}$
(b) 8.4 Newtons
(c) 19.17 cm

3.7 Problem-Solving Strategies - Graphs

Learning Objectives

- Read and understand given problem situations.
- Use the strategy: read a graph.
- Develop and apply the strategy: make a graph.
- Solve real-world problems using selected strategies as part of a plan.

Introduction

In this chapter, we have been solving problems where quantities are linearly related to each other. In this section, we will look at a few examples of linear relationships that occur in real-world problems. Remember back to our Problem Solving Plan.

Step 1:

Understand the problem

Read the problem carefully. Once the problem is read, list all the components and data that are involved. This is where you will be assigning your variables.

Step 2:

Devise a plan Translate

Come up with a way to solve the problem. Set up an equation, draw a diagram, make a chart or construct a table as a start to solving your problem.

Step 3:

Carry out the plan Solve

This is where you solve the equation you came up with in Step 2.

Step 4:

Look Check and Interpret

Check to see if your answer makes sense.

Lets look at an example that investigates a geometrical relationship.



Example 1

A cell phone company is offering its costumers the following deal. You can buy a new cell phone for \$60 and pay a monthly flat rate of \$40 per month for unlimited calls. How much money will this deal cost you after 9 months?

Solution

Lets follow the problem solving plan.

Step 1:

cell phone = \$60, calling plan = \$40per month

Let x = number of months


Let y => cost in dollars

Step 2: Lets solve this problem by making a graph that shows the number of months on the horizontal axis and the cost on the vertical axis.


Since you pay \$60 for the phone when you get the phone, then the y -intercept is $(0, 60)$.

You pay \$40 for each month, so the cost rises by \$40 for one month, so the slope = 40.

We can graph this line using the slope-intercept method.



Step 3: The question was: *How much will this deal cost after 9 months?*



We can now read the answer from the graph. We draw a vertical line from 9 months until it meets the graph, and then draw a horizontal line until it meets the vertical axis.

We see that after 9 months **you pay approximately \$420**.


Step 4: To check if this is correct, let's think of the deal again. Originally, you pay \$60 and then \$40 for 9 months.

$$\begin{aligned}\text{Phone} &= \$60 \\ \text{Calling plan} &= \$40 \times 9 = \$360 \\ \text{Total cost} &= \$420.\end{aligned}$$

The answer checks out.

Example 2

A stretched spring has a length of 12 inches when a weight of 2 lbs is attached to the spring. The same spring has a length of 18 inches when a weight of 5 lbs is attached to the spring. It is known from physics that within certain weight limits, the function that describes how much a spring stretches with different weights is a linear function. What is the length of the spring when no weights are attached?



Solution

Let's apply problem solving techniques to our problem.

Step 1:

We know: the length of the spring = 12 inches when weight = 2 lbs.

the length of the spring = 18 inches when weight = 5 lbs.

We want: the length of the spring when the weight = 0 lbs.

Let x = the weight attached to the spring.

Let y = the length of the spring


Step 2

Let's solve this problem by making a graph that shows the weight on the horizontal axis and the length of the spring on the vertical axis.

We have two points we can graph.

When the weight is 2 lbs, the length of the spring is 12 inches. This gives point (2, 12).

When the weight is 5 lbs, the length of the spring is 18 inches. This gives point (5, 18).



If we join these two points by a line and extend it in both directions we get the relationship between weight and length of the spring.

Step 3

The question was: *What is the length of the spring when no weights are attached?*

We can answer this question by reading the graph we just made. When there is no weight on the spring, the x value equals to zero, so we are just looking for the y -intercept of the graph. Looking at the graph we see that the y -intercept is **approximately** 8 inches .

Step 4

To check if this correct, lets think of the problem again.

You can see that the length of the spring goes up by 6 inches when the weight is increased by 3 lbs, so the slope of the line is $\frac{6 \text{ inches}}{3 \text{ lbs}} = 2 \text{ inches/lb}$.

To find the length of the spring when there is no weight attached, we look at the spring when there are 2 lbs attached. For each pound we take off, the spring will shorten by 2 inches. Since we take off 2 lbs, the spring will be shorter by 4 inches. So, the length of the spring with no weights is 12 inches $- 4 \text{ inches} = 8 \text{ inches}$.

The answer checks out.



Example 3

Christine took one hour to read 22 pages of Harry Potter and the Order of the Phoenix. She has 100 pages left to read in order to finish the book. Assuming that she reads at a constant rate of pages per hour, how much time should she expect to spend reading in order to finish the book?

Solution: Lets apply the problem solving techniques:

Step 1

We know that it takes Christine takes 1 hour to read 22 pages.

We want to know how much time it takes her to read 100 pages.

Let x = the time expressed in hours.

Let y = the number of pages.

Step 2

Lets solve this problem by making a graph that shows the number of hours spent reading on the horizontal axis and the number of pages on the vertical axis.

We have two points we can graph.

Christine takes one hour to read 22 pages. This gives point $(1, 22)$.

A second point is not given but we know that Christine takes 0 hours to read 0 pages. This gives the point $(0, 0)$.

If we join these two points by a line and extend it in both directions we get the relationship between the amount of time spent reading and the number of pages read.



Step 3

The question was: *How much time should Christine expect to spend reading 100 pages?*

We find the answer from reading the graph we draw a horizontal line from 100 pages until it meets the graph and then we draw the vertical until it meets the horizontal axis. We see that it takes **approximately** 4.5 hours to read the remaining 100 pages.

Step 4

To check if this correct, lets think of the problem again.

We know that Christine reads 22 pages per hour. This is the slope of the line or the rate at which she is reading. To find how many hours it takes her to read 100 pages, we divide the number of pages by the rate. In this case, $\frac{100 \text{ pages}}{22 \text{ pages per hour}} = 4.54 \text{ hours}$. This is very close to what we gathered from reading the graph.

The answer checks out.

Example 4

Aatif wants to buy a surfboard that costs \$249. He was given a birthday present of \$50 and he has a summer job that pays him \$6.50 per hour. To be able to buy the surfboard, how many hours does he need to work?



Solution

Lets apply the problem solving techniques.

Step 1

We know Surfboard costs \$249.

He has \$50.

His job pays \$6.50 per hour.

We want How many hours does Aatif need to work to buy the surfboard?

Let x = the time expressed in hours

Let y = Aatifs earnings

Step 2

Lets solve this problem by making a graph that shows the number of hours spent working on the horizontal axis and Aatifs earnings on the vertical axis.

Peter has \$50 at the beginning. This is the y -intercept of $(0, 50)$.

He earns \$6.50 per hour. This is the slope of the line.

We can graph this line using the slope-intercept method. We graph the y intercept of $(0, 50)$ and we know that for each unit in the horizontal direction the line rises by 6.5 units in the vertical direction. Here is the line that describes this situation.

**Step 3**

The question was *How many hours does Aatif need to work in order to buy the surfboard?*

We find the answer from reading the graph. Since the surfboard costs \$249, we draw a horizontal line from \$249 on the vertical axis until it meets the graph and then we draw a vertical line downwards until it meets the horizontal axis. We see that it takes **approximately 31 hours** to earn the money.

Step 4

To check if this correct, lets think of the problem again.

We know that Aatif has \$50 and needs \$249 to buy the surfboard. So, he needs to earn $\$249 - \$50 = \$199$ from his job.

His job pays \$6.50 per hour. To find how many hours he need to work we divide $\frac{\$199}{\$6.50 \text{ per hour}} = 30.6$ hours. This is very close to the result we obtained from reading the graph.

The answer checks out.

Lesson Summary

The four steps of the **problem solving plan** are:

1. **Understand the problem**
2. **Devise a plan Translate.** Build a graph.
3. **Carry out the plan Solve.** Use the graph to answer the question asked.
4. **Look Check and Interpret**

Review Questions

Solve the following problems by making a graph and reading a graph.

1. A gym is offering a deal to new members. Customers can sign up by paying a registration fee of \$200 and a monthly fee of \$39. How much will this membership cost a member by the end of the year?
2. A candle is burning at a linear rate. The candle measures five inches two minutes after it was lit. It measures three inches eight minutes after it was lit. What was the original length of the candle?
3. Tali is trying to find the width of a page of his telephone book. In order to do this, he takes a measurement and finds out that 550 pages measures 1.25 inches . What is the width of one page of the phone book?
4. Bobby and Petra are running a lemonade stand and they charge 45 cents for each glass of lemonade. In order to break even they must make \$25. How many glasses of lemonade must they sell to break even?

Review Answers

1. \$668
2. 5.67 inches
3. 0.0023 inches
4. 56 glasses

CHAPTER 4**Determining Linear Equations****Chapter Outline**

- 4.1 LINEAR EQUATIONS IN SLOPE-INTERCEPT FORM**
 - 4.2 LINEAR EQUATIONS IN POINT-SLOPE FORM**
 - 4.3 LINEAR EQUATIONS IN STANDARD FORM**
 - 4.4 EQUATIONS OF PARALLEL AND PERPENDICULAR LINES**
-

4.1 Linear Equations in Slope-Intercept Form

Learning Objectives

- Write an equation given slope and y -intercept.
- Write an equation given the slope and a point.
- Write an equation given two points.
- Write a linear function in slope-intercept form.
- Solve real-world problems using linear models in slope-intercept form.

Introduction

We saw in the last chapter that linear graphs and equations are used to describe a variety of real-life situations. In mathematics, we want to find equations that explain a situation as presented in a problem. In this way, we can determine the rule that describes the relationship between the variables in the problem. Knowing the equation or rule is very important since it allows us to find the values for the variables. There are different ways to find an equation that describes the problem. The methods are based on the information you can gather from the problem. In graphing these equations, we will assume that the domain is all real numbers.

Write an Equation Given Slope and

Lets start by learning how to write an equation in slopeintercept form $y = mx + b$.

b is the y -intercept (*the value of y when $x = 0$. This is the point where the line crosses the y -axis*).

m is the slope *how the quantity y changes with each one unit of x .*

If you are given the slope and y -intercept of a line:

1. Start with the slopeintercept form of the line $y = mx + b$.
2. Substitute the given values of m and b into the equation.

Example 1

- a) Write an equation with a slope = 4 and a y -intercept = -3 .
- b) Write an equation with a slope = -2 and a y -intercept = 7 .
- c) Write an equation with a slope = $\frac{2}{3}$ and a y -intercept = $\frac{4}{5}$.

a) Solution

We are given $m = 4$ and $b = -3$. Plug these values into the slopeintercept form $y = mx + b$.

$$y = 4x - 3$$

b) Solution

We are given $m = -2$ and $b = 7$. Plug these values into the slopeintercept form $y = mx + b$.

$$y = -2x + 7$$

c) Solution

We are given $m = \frac{2}{3}$ and $b = \frac{4}{5}$. Plug these values into the slope-intercept form $y = mx + b$.


$$y = \frac{2}{3}x + \frac{4}{5}$$

You can also write an equation in slope-intercept form if you are given the graph of the line.


Example 2

Write the equation of each line in slope-intercept form.


a)




b)



c)



d)



a) The y -intercept = -4 and the slope = $-\frac{5}{2}$. Plug these values into the slope-intercept form $y = mx + b$.

Solution

$$-\frac{5}{2}x - 4$$

b) The y -intercept = 2 and the slope = $\frac{3}{1}$. Plug these values into the slope-intercept form $y = mx + b$.

Solution

$$y = 3x + 2$$

c) The y -intercept = 4 and the slope = $-\frac{1}{1}$. Plug these values into the slope-intercept form $y = mx + b$.

Solution

$$y = -x + 4$$

d) The y -intercept = -2 and the slope = $\frac{1}{2}$. Plug these values into the slope-intercept form $y = mx + b$.

Solution

$$y = \frac{1}{2}x - 2.$$

Write an Equation Given the Slope and a Point

Often, we don't know the value of the y -intercept, but we know the value of y for a non-zero value of x . In this case we can still use the slope-intercept form to find the equation of the line.

For example, we are told that the slope of a line is two and that the line passes through the point $(1, 4)$. To find the equation of the line, we start with the slope-intercept form of a line.

$$y = mx + b$$

Plug in the value of the slope.

We don't know the value of b but we know that the slope is two, and that point $(1, 4)$ is on this line. Where the x value is one, and the y value is four. We plug this point in the equation and solve for b .

$$\begin{aligned} 4 &= 2(1) + b \\ 4 &= 2 + b \\ -2 &= -2 \\ 2 &= b \end{aligned}$$

Therefore the equation of this line is $y = 2x + 2$.

If you are given the slope and a point on the line:

1. Start with the slope-intercept form of the line $y = mx + b$.
2. Plug in the given value of m into the equation.
3. Plug the x and y values of the given point and solve for b .
4. Plug the value of b into the equation.

Example 3

Write the equation of the line in slope-intercept form.

- a) The slope of the line is 4 and the line contains point $(-1, 5)$.
 b) The slope of the line is $-\frac{2}{3}$ and the line contains point $(2, -2)$.
 c) The slope of the line is 3 and the line contains point $(3, -5)$.

Solution

a)

Start with the slope-intercept form of the line	$y = mx + b$
Plug in the slope.	$y = 4x + b$
Plug point $(-1, 5)$ into the equation.	$5 = 4(-1) + b \Rightarrow b = 9$
Plug the value of b into the equation.	$y = 4x + 9$

b)

Start with the slope-intercept form of the line	$y = mx + b$
Plug in the slope.	$y = -\frac{2}{3}x + b$
Plug point $(2, -2)$ into the equation.	$-2 = -\frac{2}{3}(2) + b \Rightarrow b = -2 + \frac{4}{3} = -\frac{2}{3}$
Plug the value of b into the equation.	$y = -\frac{2}{3}x - \frac{2}{3}$

c)

Start with the slope-intercept form of the line	$y = mx + b$
Plug in the slope.	$y = -3x + b$
Plug point $(3, -5)$ into the equation.	$-5 = -3(3) + b \Rightarrow b = 4$
Plug the value of b into the equation.	$y = -3x + 4$

Write an Equation Given Two Points

One last case is when we are just given two points on the line and we are asked to write the line of the equation in slope-intercept form.

For example, we are told that the line passes through the points $(-2, 3)$ and $(5, 2)$. To find the equation of the line we start with the slope-intercept form of a line

$$y = mx + b$$

Since we don't know the slope, we find it using the slope formula $m = \frac{y_2 - y_1}{x_2 - x_1}$.

Now substitute the x_1 and x_2 and the y_1 and y_2 values into the slope formula to solve for the slope.

$$m = \frac{2-3}{5-(-2)} = -\frac{1}{7}$$

We plug the value of the slope into the slopeintercept form $y = -\frac{1}{7}x + b$

We dont know the value of b but we know two points on the line. We can plug either point into the equation and solve for b . Lets use point $(-2, 3)$.

Therefore, the equation of this line is $y = -\frac{1}{7}x + \frac{19}{7}$.

If you are given two points on the line:

1. Start with the slopeintercept form of the line $y = mx + b$
2. Use the two points to find the slope using the slope formula $m = \frac{y_2 - y_1}{x_2 - x_1}$.
3. Plug the given value of m into the equation.
4. Plug the x and y values of one of the given points into the equation and solve for b .
5. Plug the value of b into the equation.
6. Plug the other point into the equation to check the values of m and b .

Example 4

Write the equations of each line in slopeintercept form.

- a) The line contains the points $(3, 2)$ and $(-2, 4)$.
- b) The line contains the points $(-4, 1)$ and $(-2, 3)$.

Solution:

a)

1. Start with the slopeintercept form of the line $y = mx + b$.
2. Find the slope of the line. $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4-2}{-2-3} = -\frac{2}{5}$
3. Plug in the value of the slope. $y = -\frac{2}{5}x + b$
4. Plug point $(3, 2)$ into the equation. $2 = -\frac{2}{5}(3) + b \Rightarrow b = 2 + \frac{6}{5} = \frac{16}{5}$
5. Plug the value of b into the equation. $y = -\frac{2}{5}x + \frac{16}{5}$
6. Plug point $(-2, 4)$ into the equation to check. $4 = -\frac{2}{5}(-2) + \frac{16}{5} = \frac{4}{5} + \frac{16}{5} = \frac{20}{5} = 4$

b)

1. Start with the slopeintercept form of the line $y = mx + b$.
2. Find the slope of the line. $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{3-1}{-2-(-4)} = \frac{2}{2} = 1$
3. Plug in the value of the slope. $y = x + b$
4. Plug point $(-2, 3)$ into the equation. $3 = -2 + b \Rightarrow b = 5$
5. Plug the value of b into the equation. $y = x + 5$
6. Plug point $(-4, 1)$ into the equation to check. $1 = -4 + 5 = 1$

Solve Real-World Problems Using Linear Models in Slope-Intercept Form

Lets apply the methods we just learned to a few application problems that can be modeled using a linear relationship.



Example 6

Nadia has \$200 in her savings account. She gets a job that pays \$7.50 per hour and she deposits all her earnings in her savings account. Write the equation describing this problem in slope-intercept form. How many hours would Nadia need to work to have \$500 in her account?

Let's define our variables

y = amount of money in Nadia's savings account

x = number of hours

You can see that the problem gives us the y -intercept and the slope of the equation.

We are told that Nadia has \$200 in her savings account, so $b = 200$.

We are told that Nadia has a job that pays \$7.50 per hour, so $m = 7.50$.

If we plug these values in the slope-intercept form $y = mx + b$ we obtain $y = 7.5x + 200$.

To answer the question, we plug in $y = 500$ and solve for x . $500 = 7.5x + 200 \Rightarrow 7.5x = 300 \Rightarrow x = 40$ hours.

Solution

Nadia must work 40 hours if she is to have \$500 in her account.

**Example 7**

A stalk of bamboo of the family *Phyllostachys nigra* grows at a steady rate of 12 inches per day and achieves its full height of 720 inches in 60 days. Write the equation describing this problem in slope-intercept form.

How tall is the bamboo 12 days after it started growing?

Let's define our variables

y = the height of the bamboo plant in inches

x = number of days

You can see that the problem gives us the slope of the equation and a point on the line.

We are told that the bamboo grows at a rate of 12 inches per day, so $m = 12$.

We are told that the plant grows to 720 inches in 60 days, so we have the point $(60, 720)$.

Start with the slope-intercept form of the line

$$y = mx + b$$

Plug in the slope.

$$y = 12x + b$$

Plug in point $(60, 720)$.

$$720 = 12(60) + b \Rightarrow b = 0$$

Plug the value of b back into the equation.

$$y = 12x$$

To answer the question, plug in $x = 12$ to obtain $y = 12(12) = 144$ inches.

Solution

The bamboo is 144 inches (12 feet!) tall 12 days after it started growing.

Example 8

Petra is testing a bungee cord. She ties one end of the bungee cord to the top of a bridge and to the other end she ties different weights and measures how far the bungee stretches. She finds that for a weight of 100 lb, the bungee stretches to 265 feet and for a weight of 120 lb, the bungee stretches to 275 feet. Physics tells us that in a certain range of values, including the ones given here, the amount of stretch is a linear function of the weight. Write the equation describing this problem in slope-intercept form. What should we expect the stretched length of the cord to be for a weight of 150 lbs?

Let's define our variables

y = the stretched length of the bungee cord in feet

x = the weight attached to the bungee cord in pounds

You can see that the problem gives us two points on the line.

We are told that for a weight of 100 lbs the cord stretches to 265 feet, so we have point (100, 265).

We are told that for a weight of 200 lbs the cord stretches to 275 feet, so we have point (120, 270).

Start with the slope-intercept form of the line

$$y = mx + b$$

Find the slope of the line.

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{270 - 265}{120 - 100} = \frac{5}{20} = \frac{1}{4}$$

Plug in the value of the slope.

$$y = \frac{1}{4}x + b$$

Plug point (100, 265) into the equation.

$$265 = \frac{1}{4}(100) + b \Rightarrow b = 265 - 25 = 240$$

Plug the value of b into the equation.

$$y = \frac{1}{4}x + 240$$

To answer the question, we plug in $x = 150$. $y = \frac{1}{4}(150) + 240 \Rightarrow y = 37.5 + 240 = 277.5$ feet

Solution

For a weight of 150 lbs we expect the stretched length of the cord to be 277.5 feet.

Lesson Summary

- The equation of a line in **slope-intercept** form is $y = mx + b$.

Where m is the slope and $(0, b)$ is the y -intercept).

- If you are **given the slope and y -intercept** of a line:

1. Simply plug m and b into the equation.

- If you are **given the slope and a point** on the line:

- Plug in the given value of m into the equation.
- Plug the x and y values of the given point and solve for b .
- Plug the value of b into the equation.

- If you are given two points on the line:

- Use the two points to find the slope using the slope formula $m = \frac{y_2 - y_1}{x_2 - x_1}$.



- Plug the value of m into the equation.
- Plug the x and y values of one of the given points and solve for b .
- Plug the value of b into the equation.
- Plug the other point into the equation to check the values of m and b .

Review Questions

Find the equation of the line in slope-intercept form.

- The line has slope of 7 and y -intercept of -2 .
- The line has slope of -5 and y -intercept of 6.
- The line has slope of $-\frac{1}{4}$ and contains point $(4, -1)$.
- The line has slope of $\frac{2}{3}$ and contains point $(\frac{1}{2}, 1)$.
- The line has slope of -1 and contains point $(\frac{4}{5}, 0)$.
- The line contains points $(2, 6)$ and $(5, 0)$.
- The line contains points $(5, -2)$ and $(8, 4)$.
- The line contains points $(3, 5)$ and $(-3, 0)$.
- The line contains points $(10, 15)$ and $(12, 20)$.

Write the equation of each line in slope-intercept form.



Find the equation of the linear equation in slope-intercept form.

- $m = 5, f(0) = -3$
- $m = -7, f(2) = -1$
- $m = \frac{1}{3}, f(-1) = \frac{2}{3}$
- $m = 4.2, f(-3) = 7.1$
- To buy a car, Andrew puts a down payment of \$1500 and pays \$350 per month in installments. Write an equation describing this problem in slope-intercept form. How much money has Andrew paid at the end of one year?
- Anne transplants a rose seedling in her garden. She wants to track the growth of the rose so she measures its height every week. On the third week, she finds that the rose is 10 inches tall and on the eleventh week she finds that the rose is 14 inches tall. Assuming the rose grows linearly with time, write an equation describing this problem in slope-intercept form. What was the height of the rose when Anne planted it?
- Ravi hangs from a giant spring whose length is 5 m. When his child Nimi hangs from the spring its length is 2 m. Ravi weighs 160 lbs. and Nimi weighs 40 lbs. Write the equation for this problem in slope-intercept form. What should we expect the length of the spring to be when his wife Amardeep, who weighs 140 lbs., hangs from it?

Review Answers

1. $y = 7x - 2$
2. $y = -5x + 6$
3. $y = -\frac{1}{4}x$
4. $y = \frac{2}{3}x + \frac{2}{3}$
5. $y = -1x + \frac{4}{5}$
6. $y = -2x + 10$
7. $y = 2x - 12$
8. $y = \frac{5}{6}x + \frac{5}{2}$
9. $y = \frac{5}{2}x - 10$
10. $y = -x + 3$
11. $y = 4x - 6$
12. $y = 5x - 3$
13. $y = -7x + 13$
14. $y = \frac{1}{3}x + 1$
15. $y = 4.2x + 19.7$
16. $y = 350x + 1500$; $y = \$5700$
17. $y = 0.5x + 8.5$; $y = 8.5$ inches
18. $y = .025x + 1$ or $y = \frac{1}{40}x + 1$; $y = 4.5$ m

4.2 Linear Equations in Point-Slope Form

Learning Objectives

- Write an equation in point-slope form.
- Graph an equation in point-slope form.
- Write a linear function in point-slope form.
- Solve real-world problems using linear models in point-slope form.

Introduction

In the last lesson, we saw how to write the equation of a straight line in slope-intercept form. We can rewrite this equation in another way that sometimes makes solving the problem easier. The equation of a straight line that we are going to talk about is called **point-slope form**.

$$y - y_0 = m(x - x_0)$$

Here m is the slope and (x_0, y_0) is a point on the line. Lets see how we can use this form of the equation in the three cases that we talked about in the last section.

Case 1: You know the slope of the line and the y -intercept. Case 2: You know the slope of the line and a point on the line. Case 3: You know two points on the line.

Write an Equation in Point-Slope Form

Case 1 You know the slope and the y -intercept.

1. Start with the equation in point-slope form $y - y_0 = m(x - x_0)$.
2. Plug in the value of the slope.
3. Plug in 0 for x_0 and b for y_0 .

Example 1

Write the equation of the line in point-slope form, given that the slope = -5 and the y -intercept = 4 .

Solution:

1. Start with the equation in point-slope form. $y - y_0 = m(x - x_0)$
2. Plug in the value of the slope. $y - y_0 = -5(x - x_0)$
3. Plug in 0 for x_0 and 4 for y_0 . $y - (-4) = -5(x - (0))$

Therefore, the equation is $y + 4 = -5x$

Case 2 You know the slope and a point on the line.

1. Start with the equation in point-slope form $y - y_0 = m(x - x_0)$.
2. Plug in the value of the slope.

3. Plug in the x and y values in place of x_0 and y_0 .

Example 2

Write the equation of the line in point-slope form, given that the slope = $\frac{3}{5}$ and the point $(2, 6)$ is on the line.

Solution:

1. Start with the equation in point-slope form. $y - y_0 = m(x - x_0)$
2. Plug in the value of the slope. $y - y_0 = \frac{3}{5}(x - x_0)$
3. Plug in 2 for x_0 and 6 for y_0 . $y - (6) = \frac{3}{5}(x - (2))$

The equation is $y - 6 = \frac{3}{5}(x - 2)$

Notice that the equation in point-slope form is not solved for y .

Case 3 You know two points on the line.

1. Start with the equation in point-slope form $y - y_0 = m(x - x_0)$.
2. Find the slope using the slope formula. $m = \frac{y_2 - y_1}{x_2 - x_1}$
3. Plug in the value of the slope.
4. Plug in the x and y values of one of the given points in place of x_0 and y_0

Example 3

Write the equation of the line in point-slope form, given that the line contains points $(-4, -2)$ and $(8, 12)$.

Solution

1. Start with the equation in point-slope form. $y - y_0 = m(x - x_0)$
2. Find the slope using the slope formula. $m = \frac{12 - (-2)}{8 - (-4)} = \frac{14}{12} = \frac{7}{6}$
3. Plug in the value of the slope. $y - y_0 = \frac{7}{6}(x - x_0)$
4. Plug in -4 for x_0 and -2 for y_0 . $y - (-2) = \frac{7}{6}(x - (-4))$

Therefore, the equation is $y + 2 = \frac{7}{6}(x + 4)$ **Answer 1**

In the last example, you were told that for the last step you could choose either of the points you were given to plug in for the point (x_0, y_0) but it might not seem like you would get the same answer if you plug the second point in instead of the first. Lets redo Step 4.

4. Plug in 8 for x_0 and 12 for y_0 . $y - 12 = \frac{7}{6}(x - 8)$ **Answer 2**

This certainly does not seem like the same answer as we got by plugging in the first point. What is going on?

Notice that the equation in point-slope form is not solved for y . Lets change both answers into slope-intercept form by solving for y .

$$\begin{aligned} &\text{Answer 1} \\ y + 2 &= \frac{7}{6}(x + 4) \\ y + 2 &= \frac{7}{6}x + \frac{28}{6} \\ y &= \frac{7}{6}x + \frac{14}{3} - 2 \\ y &= \frac{7}{6}x + \frac{8}{3} \end{aligned}$$

$$\begin{aligned} &\text{Answer 2} \\ y - 12 &= \frac{7}{6}(x - 8) \\ y - 12 &= \frac{7}{6}x - \frac{56}{6} \\ y &= \frac{7}{6}x - \frac{28}{3} + 12 \\ y &= \frac{7}{6}x + \frac{8}{3} \end{aligned}$$

Now that the two answers are solved for y , you can see that they simplify to the same thing. In point-slope form you can get an infinite number of right answers, because there are an infinite number of points on a line. The slope of the line will always be the same but the answer will look different because you can substitute any point on the line for (x_0, y_0) . However, regardless of the point you pick, the point-slope form should always simplify to the same slope-intercept equation for points that are on the same line.

In the last example you saw that sometimes we need to change between different forms of the equation. To change from point-slope form to slope-intercept form, we just solve for y .

Example 4

Re-write the following equations in slope-intercept form.

a) $y - 5 = 3(x - 2)$

b) $y + 7 = -(x + 4)$

Solution

a) To re-write in slope-intercept form, solve for y .

$$\begin{aligned} y - 5 &= 3(x - 2) \\ -5 &= 3x - 6 \\ y &= 3x - 1 \end{aligned}$$

b) To re-write in slope-intercept form, solve for y .

$$\begin{aligned} y + 7 &= -(x + 4) \\ y + 7 &= -x - 4 \\ y &= -x - 11 \end{aligned}$$

Graph an Equation in Point-Slope Form

If you are given an equation in point-slope form, it is not necessary to re-write it in slope-intercept form in order to graph it. The point-slope form of the equation gives you enough information so you can graph the line $y - y_0 = m(x - x_0)$. From this equation, we know a point on the line (x_0, y_0) and the slope of the line.

To graph the line, you first plot the point (x_0, y_0) . Then the slope tells you how many units you should go up or down and how many units you should go to the right to get to the next point on the line. Lets demonstrate this method with an example.

Example 5


Make a graph of the line given by the equation $y - 2 = \frac{2}{3}(x + 2)$

Solution


Lets rewrite the equation $y - (2) = \frac{2}{3}(x + 2)$.

Now we see that point $(-2, 2)$ is on the line and that the slope $= \frac{2}{3}$.


First plot point $(-2, 2)$ on the graph.



A slope of $\frac{2}{3}$ tells you that from your point you should move 2 units up and 3 units to the right and draw another point.



Now draw a line through the two points and extend the line in both directions.



Solve Real-World Problems Using Linear Models in Point-Slope Form

Lets solve some word problems where we need to write the equation of a straight line in point-slope form.



Example 6

Marciel rented a moving truck for the day. Marciel only remembers that the rental truck company charges \$40 per day and some amount of cents per mile. Marciel drives 46 miles and the final amount of the bill (before tax) is \$63. What is the amount per mile the truck rental company charges per day? Write an equation in point-slope form that describes this situation. How much would it cost to rent this truck if Marciel drove 220 miles?

Lets define our variables:

x = distance in miles

y = cost of the rental truck in dollars

We see that we are given the y -intercept and the point $(46, 63)$.

Peter pays a flat fee of \$40 for the day. This is the y intercept.

He pays \$63 for 46 miles this is the coordinate point $(46, 63)$.

Start with the point-slope form of the line. $(y - y_0) = m(x - x_0)$

Plug in the coordinate point. $63 - y_0 = m(46 - x_0)$

Plug in point $(0, 40)$. $63 - 40 = m(46 - 0)$

Solve for the slope. $23 = m(46) \rightarrow m = \frac{23}{46} = 0.5$


The slope is : 0.5 dollars per mile

So, the truck company charges 50 cents per mile. ($\$0.5 = 50$ cents) Equation of line is : $y = 0.5x + 40$

To answer the question of 220 miles we plug in $x = 220$.

Solution

$$y - 40 = 0.5(220) \Rightarrow y = \$150$$



Example 7

Anne got a job selling window shades. She receives a monthly base salary and a \$6 commission for each window shade she sells. At the end of the month, she adds up her sales and she figures out that she sold 200 window shades and made \$2500. Write an equation in point-slope form that describes this situation. How much is Annes monthly base salary?

Lets define our variables

x = number of window shades sold

y = Annes monthly salary in dollars

We see that we are given the slope and a point on the line:

Anne gets \$6 for each shade, so the slope = 6 dollars/shade.

She sold 200 shades and made \$2500, so the point is $(200, 2500)$.

Start with the point-slope form of the line. $y - y_0 = m(x - x_0)$

Plug in the coordinate point. $y - y_0 = 6(x - x_0)$

Plug in point $(200, 2500)$. $y - 2500 = 6(x - 200)$

Annes base salary is found by plugging in $x = 0$. We obtain $y - 2500 = -1200 \Rightarrow y = \1300

Solution

Annes monthly base salary is \$1300.

Lesson Summary

- The **point-slope form** of an equation for a line is: $y - y_0 = m(x - x_0)$.
- If you are **given the slope and a point** on the line:

1. Simply plug the point and the slope into the equation.
- If you are **given the slope and y-intercept** of a line:
 1. Plug the value of m into the equation
 2. Plug the y-intercept point into the equation $y_0 = y\text{-intercept}$ and $x_0 = 0$.
- If you are **given two points** on the line:
 1. Use the two points to find the slope using the slope formula $m = \frac{y_2 - y_1}{x_2 - x_1}$.
 2. Plug the value of m into the equation.
 3. Plug either of the points into the equation as (x_0, y_0) .
- The **functional notation of point-slope form** is $f(x) - f(x_0) = m(x - x_0)$.

Review Questions

Write the equation of the line in point-slope form.

1. The line has slope $-\frac{1}{10}$ and goes through point $(10, 2)$.
2. The line has slope -75 and goes through point $(0, 125)$.
3. The line has slope 10 and goes through point $(8, -2)$.
4. The line goes through the points $(-2, 3)$ and $(-1, -2)$.
5. The line contains points $(10, 12)$ and $(5, 25)$.
6. The line goes through points $(2, 3)$ and $(0, 3)$.
7. The line has a slope $\frac{3}{5}$ and a yintercept -3 .
8. The line has a slope -6 and a y-intercept 0.5 .

Write the equation of the line in point-slope form.

9. $m = -\frac{1}{5}$ and point $(0, 7)$
10. $m = -12$ and point $(-2, 5)$
11. The line contains points $(-7, 5)$ and $(3, -4)$
12. The line contains points $(6, 0)$ and $(0, 6)$
13. $m = 3$ and contains point $(2, -9)$
14. $m = -\frac{9}{5}$ and contains point $(0, 32)$
15. Nadia is placing different weights on a spring and measuring the length of the stretched spring. She finds that for a 100 gram weight the length of the stretched spring is 20 cm and for a 300 gram weight the length of the stretched spring is 25 cm. Write an equation in point-slope form that describes this situation. What is the unstretched length of the spring?
16. Andrew is a submarine commander. He decides to surface his submarine to periscope depth. It takes him 20 minutes to get from a depth of 400 feet to a depth of 50 feet. Write an equation in point-slope form that describes this situation. What was the submarines depth five minutes after it started surfacing?

Review Answers

1. $y - 2 = -\frac{1}{10}(x - 10)$
2. $y - 125 = -75x$
3. $y + 2 = 10(x - 8)$
4. $y + 2 = -5(x + 1)$ or $y - 3 = -5(x + 2)$
5. $y - 25 = -\frac{13}{5}(x - 5)$ or $y - 12 = -\frac{13}{5}(x - 10)$
6. $y - 3 = 0$
7. $y + 3 = \frac{3}{5}x$
8. $y - 0.5 = -6x$

9. $y - 7 = -\frac{1}{5}x$

10. $y - 5 = -12(x + 2)$

11. $y - 5 = -\frac{9}{10}(x + 7)$ or $y + 4 = -\frac{9}{10}(x - 3)$

12. $y = -1(x - 6)$ or $y - 6 = -x$

13. $y + 9 = 3(x - 2)$

14. $y - 32 = \frac{9}{5}x$

15. $y - 20 = \frac{1}{40}(x - 100)$ unstretched length = 17.5 cm

16. $y - 50 = -17.5(x - 20)$ or $y - 400 = -17.5x$ depth = 312.5 feet

4.3 Linear Equations in Standard Form

Learning Objectives

- Write equivalent equations in standard form.
- Find the slope and y -intercept from an equation in standard form.
- Write equations in standard form from a graph.
- Solve real-world problems using linear models in standard form.

Introduction

In this section, we are going to talk about the standard form for the equation of a straight line. The following linear equation is said to be in standard form.

$$ax + by = c$$

Here a , b and c are constants that have no factors in common and the constant a is a non-negative value. Notice that the b in the standard form is different than the b in the slope-intercept form. There are a few reasons why standard form is useful and we will talk about these in this section. The first reason is that standard form allows us to write equations for vertical lines which is not possible in slope-intercept form.

For example, let's find the equation of the line that passes through points $(2, 6)$ and $(2, 9)$.

Let's try the slope-intercept form $y = mx + b$

We need to find the slope $m = \frac{9-6}{2-2} = \frac{3}{0}$. The slope is undefined because we cannot divide by zero.

The point-slope form $y - y_0 = m(x - x_0)$ also needs the slope, so we cannot write an equation for this line in either the slope-intercept or the point-slope form.

Since we have two points in a plane, we know that a line passes through these two points, but how do we find the equation of that line? It turns out that this line has no y value in it. Notice that the value of x in both points is two for the different values of y , so we can say that it does not matter what y is because x will always equal two. Here is the equation in standard form.

$$1 \cdot x + 0 \cdot y = 2 \text{ or } x = 2$$

The line passing through point $(2, 6)$ and $(2, 9)$ is a **vertical line** passing through $x = 2$. Note that the equation of a horizontal line would have no x variable, since y would always be the same regardless of the value of x . For example, a **horizontal line** passing through point $(0, 5)$ has this equation in standard form.

$$0 \cdot x + 1 \cdot y = 5 \text{ or } y = 5$$

Write Equivalent Equations in Standard Form

So far you have learned how to write equations of lines in slope-intercept form and point-slope form. Now you will see how to rewrite equations in standard form.

Example 1

Rewrite the following equations in standard form.

a) $y = 5x - 7$

b) $y - 2 = -3(x + 3)$

c) $y = \frac{2}{3}x + \frac{1}{2}$

Solution

We need to rewrite the equations so that all the variables are on one side of the equation and the coefficient of x is not negative.

a) $y = 5x - 7$

Subtract y from both sides.

$$0 = 5x - y - 7$$

Add 7 to both sides.

$$7 = 5x - y$$

The equation in standard form is :

$$5x - y = 7$$

b) $y - 2 = -3(x + 3)$

Distribute the -3 on the right-hand-side.

$$y - 2 = -3x - 9$$

Add $3x$ to both sides.

$$y + 3x - 2 = -9$$

Add 2 to both sides.

$$y + 3x = -7$$

The equation in standard form is :

$$y + 3x = -7$$

c) $y = \frac{2}{3}x + \frac{1}{2}$

Find the common denominator for all terms in the equation. In this case, the common denominator equals 6.

Multiply all terms in the equation by 6.

$$6 \left(y = \frac{2}{3}x + \frac{1}{2} \right) \Rightarrow 6y = 4x + 3$$

Subtract $6y$ from both sides.

$$0 = 4x - 6y + 3$$

Subtract 3 from both sides.

$$-3 = 4x - 6y$$

The equation in standard form is :

$$4x - 6y = -3$$

Find the Slope and

The slope-intercept form and the point-slope form of the equation for a straight line both contain the slope of the equation explicitly, but the standard form does not. Since the slope is such an important feature of a line, it is useful to figure out how you would find the slope if you were given the equation of the line in standard form.

$$ax + by = c$$

Lets rewrite this equation in slope-intercept form by solving the equation for y .

Subtract ax from both sides.

$$by = -ax + c$$

Divide all terms by b .

$$y = -\frac{a}{b}x + \frac{c}{b}$$

If we compare with the slope-intercept form $y = mx + b$, we see that the slope, $m = -\frac{a}{b}$ and the y -intercept $= \frac{c}{b}$. Again, notice that the b in the standard form is different than the b in the slope-intercept form.

Example 2

Find the slope and the y -intercept of the following equations written in standard form:

a) $3x + 5y = 6$

b) $2x - 3y = -8$

c) $x - 5y = 10$

Solution

The slope $m = -\frac{a}{b}$ and the y -intercept $= \frac{c}{b}$.

a) $3x + 5y = 6$ $m = -\frac{3}{5}$ and y -intercept $= \frac{6}{5}$

b) $2x - 3y = -8$ $m = \frac{2}{3}$ and y -intercept $= \frac{8}{3}$

c) $x - 5y = 10$ $m = \frac{1}{5}$ and y -intercept $= \frac{10}{-5} = -2$

Write Equations in Standard Form From a Graph

If we are given a graph of a straight line, it is fairly simple to write the equation in slope-intercept form by reading the slope and y -intercept from the graph. Lets now see how to write the equation of the line in standard form if we are given the graph of the line.

First, remember that to graph an equation from standard form we can use the cover-up method to find the intercepts of the line. For example, lets graph the line given by the equation $3x - 2y = 6$.

To find the x -intercept, cover up the y term (remember, x -intercept is where $y = 0$).

$$3x - \cancel{2y} = 6$$

$$3x = 6 \Rightarrow x = 2$$

The x -intercept is $(2, 0)$


To find the y -intercept, cover up the x term (remember, y -intercept is where $x = 0$).

$$\cancel{3x} - 2y = 6$$

$$-2y = 6 \Rightarrow y = -3$$

The y -intercept is $(0, -3)$

We plot the intercepts and draw a line through them that extends in both directions.




Now we want to apply this process in reverse. If we have the graph of the line, we want to write the equation of the line in standard form.


Example 3

Find the equation of the line and write in standard form.


a)



b)



c)



Solution

a) We see that the x -intercept is $(3, 0) \Rightarrow x = 3$ and the y -intercept is $(0, -4) \Rightarrow y = -4$.

We saw that in standard form $ax + by = c$,

if we cover up the y term, we get $ax = c$

if we cover up the x term, we get $by = c$

We need to find the numbers that when multiplied with the intercepts give the same answer in both cases. In this case, we see that multiplying $x = 3$ by 4 and multiplying $y = -4$ by -3 gives the same result.

$$(x = 3) \times 4 \Rightarrow 4x = 12 \text{ and } (y = -4) \times (-3) \Rightarrow -3y = 12$$

Therefore, $a = 4$, $b = -3$ and $c = 12$ and the standard form is:

$$4x - 3y = 12$$

b) We see that the x -intercept is $(3, 0) \Rightarrow x = 3$ and the y -intercept is $(0, 3) \Rightarrow y = 3$.

The values of the intercept equations are already the same, so $a = 1$, $b = 1$ and $c = 3$. The standard form is:

$$x + y = 3$$

c) We see that the x -intercept is $(\frac{3}{2}, 0) \Rightarrow x = \frac{3}{2}$ and the y -intercept is $(0, 4) \Rightarrow y = 4$.

Lets multiply the x -intercept equation by 2 $\Rightarrow 2x = 3$

Then we see we can multiply the x -intercept again by 4 and the y -intercept by 3.

$$\Rightarrow 8x = 12 \text{ and } 3y = 12$$

The standard form is $8x + 3y = 12$.

Solve Real-World Problems Using Linear Models in Standard Form

Here are two examples of real-world problems where the standard form of the equation is useful.

Example 4



Nimitha buys fruit at her local farmers market. This Saturday, oranges cost \$2per pound and cherries cost \$3per pound. She has \$12to spend on fruit. Write an equation in standard form that describes this situation. If she buys 4 pounds of oranges, how many pounds of cherries can she buy?

Solution

Lets define our variables

x = pounds of oranges

y =pounds of cherries

The equation that describes this situation is: $2x + 3y = 12$

If she buys 4 pounds of oranges, we plug $x = 4$ in the equation and solve for y .

$$2(4) + 3y = 12 \Rightarrow 3y = 12 - 8 \Rightarrow 3y = 4 \Rightarrow y = \frac{4}{3}$$

Nimitha can buy $1\frac{1}{3}$ pounds of cherries.



Example 5

Jethro skateboards part of the way to school and walks for the rest of the way. He can skateboard at 7miles per hour and he can walk at 3miles per hour. The distance to school is 6miles. Write an equation in standard form that describes this situation. If Jethro skateboards for $\frac{1}{2}$ an hour, how long does he need to walk to get to school?

Solution

Lets define our variables.

x = hours Jethro skateboards

y = hours Jethro walks

The equation that describes this situation is $7x + 3y = 6$

If Jethro skateboards $\frac{1}{2}$ an hour, we plug $x = 0.5$ in the equation and solve for y .

$$7(0.5) + 3y = 6 \Rightarrow 3y = 6 - 3.5 \Rightarrow 3y = 2.5 \Rightarrow y = \frac{5}{6}$$

Jethro must walk $\frac{5}{6}$ of an hour.

Lesson Summary

- A linear equation in the form $ax + by = c$ is said to be in **standard form**. Where a , b and c are constants (b is different than the y -intercept b) and a is non-negative.
- Given an equation in standard form, $ax + by = c$, the **slope**, $a = -\frac{a}{b}$, and the **y -intercept** $= \frac{c}{b}$.
- The **cover-up method** is useful for graphing an equation in standard form. To find the y -intercept, cover up the x term and solve the remaining equation for y . Likewise, to find the x -intercept, cover up the y term and solve the remaining equation for x .

Review Questions





Rewrite the following equations in standard form.

1. $y = 3x - 8$
2. $y - 7 = -5(x - 12)$
3. $2y = 6x + 9$
4. $y = \frac{9}{4}x + \frac{1}{4}$
5. $y + \frac{3}{5} = \frac{2}{3}(x - 2)$
6. $3y + 5 = 4(x - 9)$

Find the slope and y -intercept of the following lines.

7. $5x - 2y = 15$
8. $3x + 6y = 25$
9. $x - 8y = 12$
10. $3x - 7y = 20$
11. $9x - 9y = 4$
12. $6x + y = 3$

Find the equation of each line and write it in standard form.



17. Andrew has two part time jobs. One pays \$6 per hour and the other pays \$10 per hour. He wants to make \$366 per week. Write an equation in standard form that describes this situation. If he is only allowed to work 15 hour per week at the \$10 per hour job, how many hours does he need to work per week in his \$6 per hour job in order to achieve his goal?
18. Anne invests money in two accounts. One account returns 5% annual interest and the other returns 7% annual interest. In order not to incur a tax penalty, she can make no more than \$400 in interest per year. Write an equation in standard form that describes this problem. If she invests \$5000 in the 5% interest account, how much money does she need to invest in the other account?

Review Answers

1. $3x - y = 8$
2. $5x + y = 67$
3. $6x - 2y = -9$
4. $9x - 4y = -1$
5. $10x - 15y = 29$
6. $4x - 3y = 41$
7. $m = (5/2), b = -15/2$
8. $m = -(1/2), b = 25/6$
9. $m = (1/8), b = -3/2$
10. $m = (3/7), b = -20/7$
11. $m = 1, b = -4/9$
12. $m = -6, b = 3$
13. $x - 2y = 4$
14. $6x + 5y = -30$
15. $10x + 14y = 35$
16. $3x - 8y = -24$
17. $x =$ number of hours per week worked at \$6 per hour job $y =$ number of hours per week worked at \$10 per hour job Equation $6x + 10y = 366$ Answer 36 hours
18. $x =$ amount of money invested at 5% annual interest $y =$ amount of money invested at 7% annual interest Equation $5x + 7y = 40000$ Answer \$2142.86


4.4 Equations of Parallel and Perpendicular Lines

Learning Objectives

- Determine whether lines are parallel or perpendicular.
- Write equations of perpendicular lines.
- Write equations of parallel lines.
- Investigate families of lines.

Introduction

In this section, you will learn how **parallel lines** are related to each other on the coordinate plane. You will also learn how **perpendicular lines** are related to each other. Lets start by looking at a graph of two parallel lines.



The two lines will never meet because they are parallel. We can clearly see that the two lines have different y -intercepts, more specifically 6 and -4 .


How about the slopes of the lines? Are they related in any way? Because the lines never meet, they must rise at the same rate. This means that the slopes of the two lines are the same.

Indeed, if we calculate the slopes of the lines, we find the following results.

$$\text{Line A: } m = \frac{6-2}{0-(-2)} = \frac{4}{2} = 2$$

$$\text{Line B: } m = \frac{0-(-4)}{2-0} = \frac{4}{2} = 2$$

For Parallel Lines: the slopes are the same, $m_1 = m_2$, and the y -intercepts are different.




Now lets look at a graph of two perpendicular lines.

We find that we cant say anything about the y -intercepts. In this example, they are different, but they would be the same if the lines intersected at the y -intercept.

Now we want to figure out if there is any relationship between the slopes of the two lines.

First of all we see that the *slopes must have opposite signs, one negative and one positive.*



To find the slope of line **A**, we pick two points on the line and draw the blue (upper) right triangle. The legs of the triangle represent the rise and the run.

Looking at the figure $m_1 = \frac{b}{a}$

To find the slope of line **B**, we pick two points on the line and draw the red (lower) right triangle. If we look at the figure, we see that the two triangles are identical, only rotated by 90° .

Looking at the diagram $m_2 = -\frac{a}{b}$

For Perpendicular Lines: the slopes are negative reciprocals of each other. $m_1 = -\frac{1}{m_2}$ or $m_1 m_2 = -1$

Determine Whether Lines are Parallel or Perpendicular

You can find whether lines are parallel or perpendicular by comparing the slopes of the lines. If you are given points on the line you can find the slope using the formula. If you are given the equations of the lines, rewrite each equation in a form so that it is easy to read the slope, such as the slope-intercept form.

Example 1

Determine whether the lines are parallel or perpendicular or neither.

- One line passes through points $(2, 11)$ and $(-1, 2)$; another line passes through points $(0, -4)$ and $(-2, -10)$.
- One line passes through points $(-2, -7)$ and $(1, 5)$; another line passes through points $(4, 1)$ and $(-8, 4)$.
- One line passes through points $(3, 1)$ and $(-2, -2)$; another line passes through points $(5, 5)$ and $(4, -6)$.

Solution

Find the slope of each line and compare them.

$$a) m_1 = \frac{2-11}{-1-2} = \frac{-9}{-3} = 3 \text{ and } m_2 = \frac{-10-(-4)}{-2-0} = \frac{-6}{-2} = 3$$

The slopes are equal, so the lines are parallel.

$$b) m_1 = \frac{5-(-7)}{1-(-2)} = \frac{12}{3} = 4 \text{ and } m_2 = \frac{4-1}{-8-4} = \frac{3}{-12} = -\frac{1}{4}$$

The slopes are negative reciprocals of each other, so the lines are perpendicular.

$$c) m_1 = \frac{-2-1}{-2-3} = \frac{-3}{-5} = \frac{3}{5} \text{ and } m_2 = \frac{-6-5}{4-5} = \frac{-11}{-1} = 11$$

The slopes are not the same or negative reciprocals of each other, so the lines are neither parallel nor perpendicular.

Example 2

Determine whether the lines are parallel or perpendicular or neither.

- Line 1: $3x + 4y = 2$ Line 2: $8x - 6y = 5$
- Line 1: $2x = y - 10$ Line 2: $y = -2x + 5$
- Line 1: $7y + 1 = 7x$ Line 2: $x + 5 = y$

Solution

Write each equation in slope-intercept form.

$$a) \text{ Line 1 } 3x + 4y = 2 \Rightarrow 4y = -3x + 2 \Rightarrow y = -\frac{3}{4}x + \frac{1}{2} \Rightarrow \text{slope} = -\frac{3}{4}$$

$$\text{Line 2: } 8x - 6y = 5 \Rightarrow 8x - 5 = 6y \Rightarrow y = \frac{8}{6}x - \frac{5}{6} \Rightarrow y = \frac{4}{3}x - \frac{5}{6} \Rightarrow \text{slope} = \frac{4}{3}$$

The slopes are negative reciprocals of each other, so the lines are perpendicular to each other.

$$\text{b) Line 1 } 2x = y - 10 \Rightarrow y = 2x + 10 \Rightarrow \text{slope} = 2$$

$$\text{Line 2 } y = -2x + 5 \Rightarrow \text{slope} = -2$$

The slopes are not the same or negative reciprocals of each other, so the lines are neither parallel nor perpendicular.

$$\text{c) Line 1 } 7y + 1 = 7x \Rightarrow 7y = 7x - 1 \Rightarrow y = x - \frac{1}{7} \Rightarrow \text{slope} = 1$$

$$\text{Line 2: } x + 5 = y \Rightarrow y = x + 5 \Rightarrow \text{slope} = 1$$

The slopes are the same so the lines are parallel.

Write Equations of Perpendicular Lines

We can use the properties of perpendicular lines to write an equation of a line perpendicular to a given line. You will be given the equation of a line and asked to find the equation of the perpendicular line passing through a specific point. Here is the general method for solving a problem like this.

- Find the slope of the given line from its equation. You might need to rewrite the equation in a form such as the slope-intercept form.
- Find the slope of the perpendicular line by writing the negative reciprocal of the slope of the given line.
- Use the slope and the point to write the equation of the perpendicular line in point-slope form.

Example 3

Find the equation perpendicular to the line $y = -3x + 5$ that passes through point $(2, 6)$.

Solution

Find the slope of the given line $y = -3x + 5$ has a slope $= -3$.

The slope of the perpendicular line is the negative reciprocal $m = \frac{1}{3}$

Now, we are trying to find the equation of a line with slope $m = \frac{1}{3}$ that passes through point $(2, 6)$.

Use the point-slope form with the slope and the point $y - 6 = \frac{1}{3}(x - 2)$

The equation of the line could also be written as $y = \frac{1}{3}x + \frac{16}{3}$

Example 4

Find the equation of the line perpendicular to $x - 5y = 15$ that passes through the point $(-2, 5)$.

Solution

Rewrite the equation in slope-intercept form $x - 5y = 15 \Rightarrow -5y = -x + 15 \Rightarrow y = \frac{1}{5}x - 3$

The slope of the given line is $m = \frac{1}{5}$ and the slope of the perpendicular is the negative reciprocal or $m = -5$. We are looking for a line with a slope $m = -5$ that passes through the point $(-2, 5)$.

Use the point-slope form with the slope and the point $y - 5 = -5(x + 2)$

The equation of the line could also be written as $y = -5x - 5$

Example 5

Find the equation of the line perpendicular to $y = -2$ that passes through the point $(4, -2)$.

Solution

The equation is already in slope intercept form but it has an x term of $0y = 0x - 2$. This means the slope is $m = 0$.

We'd like a line with slope that is the negative reciprocal of 0. The reciprocal of 0 is $m = \frac{1}{0} = \text{undefined}$. Hmm... It seems like we have a problem. But, look again at the desired slope in terms of the definition of slope $m = \frac{\text{rise}}{\text{run}} = \frac{1}{0}$. So our desired line will move 0 units in x for every 1 unit it rises in y . This is a vertical line, so the solution is the vertical line that passes through $(4, -2)$. This is a line with an x coordinate of 4 at every point along it.

The equation of the line is: $x = 4$

Write Equations of Parallel Lines

We can use the properties of parallel lines to write an equation of a line parallel to a given line. You will be given the equation of a line and asked to find the equation of the parallel line passing through a specific point. Here is the general method for solving a problem like this.

- Find the slope of the given line from its equation. You might need to rewrite the equation in a form such as the slope-intercept form.
- The slope of the parallel line is the same as that of the given line.
- Use the slope and the point to write the equation of the perpendicular line in slope-intercept form or point-slope form.

Example 6

Find the equation parallel to the line $y = 6x - 9$ that passes through point $(-1, 4)$.

Solution

Find the slope of the given line $y = 6x - 9$ has a slope $= 6$.

Since parallel lines have the same slope, we are trying to find the equation of a line with slope $m = 6$ that passes through point $(-1, 4)$.

Start with the slope-intercept form.	$y = mx + b$
Plug in the slope	$y = 6x + b$
Plug in point $(-1, 4)$.	$4 = 6(-1) + b \Rightarrow b = 4 + 6 \Rightarrow b = 10$

The equation of the line is $y = 6x + 10$.

Example 7

Find the equation of the line parallel to $7 - 4y = 0$ that passes through the point $(9, 2)$.

Solution

Rewrite the equation in slope-intercept form.

$$7 - 4y = 0 \Rightarrow 4y - 7 = 0 \Rightarrow 4y = 7 \Rightarrow y = \frac{7}{4} \Rightarrow y = 0x + \frac{7}{4}$$

The slope of the given line is $m = 0$. This is a horizontal line.

Since the slopes of parallel lines are the same, we are looking for a line with slope $m = 0$ that passes through the point $(9, 2)$.

Start with the slope-intercept form.	$y = 0x + b$
Plug in the slope.	$y = 0x + b$
Plug in point (9, 2).	$2 = 0(9) + b$
	$\Rightarrow b = 2$

The equation of the line is $y = 2$.

Example 8

Find the equation of the line parallel to $6x - 5y = 12$ that passes through the point $(-5, -3)$.

Solution

Rewrite the equation in slope-intercept form.

$$3x - 5y = 12 \Rightarrow 5y = 6x - 12 \Rightarrow y = \frac{6}{5}x - \frac{12}{5}$$

The slope of the given line is $m = \frac{6}{5}$.


Since the slopes of parallel lines are the same, we are looking for a line with slope $m = \frac{6}{5}$ that passes through the point $(-5, -3)$.

Start with the slope-intercept form.	$y = mx + b$
Plug in the slope.	$y = \frac{6}{5}x + b$
Plug in point $(-5, -3)$.	$-3 = \frac{6}{5}(-5) + b \Rightarrow -3 = -6 + b \Rightarrow b = 3$

The equation of the line is: $y = \frac{6}{5}x + 3$

Investigate Families of Lines

A straight line has two very important properties, its **slope** and its **y-intercept**. The slope tells us how steeply the line rises or falls, and the y-intercept tells us where the line intersects the y-axis. In this section, we will look at two families of lines. A **family of lines** is a set of lines that have something in common with each other. Straight lines can belong to two types of families. One where the slope is the same and one where the y-intercept is the same.




Family 1

Keep slope unchanged and vary the y-intercept.

The figure to the right shows the family of lines $y = -2x + b$.

All the lines have a slope of -2 but the value of b is different for each of the lines.

Notice that in such a family all the lines are parallel. All the lines look the same but they are shifted up and down the y -axis. As b gets larger the line rises on the y -axis and as b gets smaller the line goes lower on the y -axis. This behavior is often called a **vertical shift**.



Family 2

Keep the y -intercept unchanged and vary the slope.

The figure to the right shows the family of lines $y = mx + 2$.

All lines have a y -intercept of two but the value of the slope is different for each of the lines. The lines start with $y = 2$ (red line) which has a slope of zero. They get steeper as the slope increases until it gets to the line $x = 0$ (purple line) which has an undefined slope.

Example 9


Write the equation of the family of lines satisfying the given condition:

- Parallel to the x -axis
- Through the point $(0, -1)$
- Perpendicular to $2x + 7y - 9 = 0$
- Parallel to $x + 4y - 12 = 0$

Solution


a) All lines parallel to the x -axis will have a slope of zero. It does not matter what the y -intercept is.

The family of lines is $y = 0 \cdot x + b$ or $y = b$.



b) All lines passing through the point $(0, -1)$ have the same y -intercept, $b = -1$.

The family of lines is $y = mx - 1$.



c) First we need to find the slope of the given line.


Rewrite $2x + 7y - 9 = 0$ in slope-intercept form $y = -\frac{2}{7}x + \frac{9}{7}$.

The slope is $-\frac{2}{7}$.

The slope of our family of lines is the negative reciprocal of the given slope $m = \frac{7}{2}$.

All the lines in this family have a slope of $m = \frac{7}{2}$ but different y -intercepts.

The family of lines is $y = \frac{7}{2}x + b$.




d) First we need to find the slope of the given line.

Rewrite $x + 4y - 12 = 0$ in slope-intercept form $y = -\frac{1}{4}x + 3$.

The slope is $m = -\frac{1}{4}$.

All the lines in the family have a slope of $m = -\frac{1}{4}$ but different y -intercepts.

The family of lines is $y = -\frac{1}{4}x + b$.



Lesson Summary

- **Parallel lines** have the same slopes, $m_1 = m_2$, but different y -intercepts.
- **Perpendicular lines** have slopes which are the negative reciprocals of each other.

$$m_1 = -\frac{1}{m_2} \text{ or } m_1 m_2 = -1$$

- **To find the line parallel (or perpendicular)** to a specific line which passes through a given point:

1. Find the slope of the given line from its equation.
2. Compute the slope parallel (or perpendicular) to the line.
3. Use the computed slope and the specified point to write the equation of the new line in point-slope form.
4. Transform from point-slope form to another form if required.

- **A family of lines** is a set of lines that have something in common with each other. There are two types of line families. One where the slope is the same and one where the y -intercept is the same.

Review Questions

Determine whether the lines are parallel, perpendicular or neither.

1. One line passes through points $(-1, 4)$ and $(2, 6)$; another line passes through points $(2, -3)$ and $(8, 1)$.
2. One line passes through points $(4, -3)$ and $(-8, 0)$; another line passes through points $(-1, -1)$ and $(-2, 6)$.

3. One line passes through points $(-3, 14)$ and $(1, -2)$; another line passes through points $(0, -3)$ and $(-2, 5)$.
4. One line passes through points $(3, 3)$ and $(-6, -3)$; another line passes through points $(2, -8)$ and $(-6, 4)$.
5. Line 1: $4y + x = 8$ Line 2: $12y + 3x = 1$
6. Line 1: $5y + 3x + 1$ Line 2: $6y + 10x = -3$
7. Line 1: $2y - 3x + 5 = 0$ Line 2: $y + 6x = -3$
8. Find the equation of the line parallel to $5x - 2y = 2$ that passes through point $(3, -2)$.
9. Find the equation of the line perpendicular to $y = -\frac{2}{5}x - 3$ that passes through point $(2, 8)$.
10. Find the equation of the line parallel to $7y + 2x - 10 = 0$ that passes through the point $(2, 2)$.
11. Find the equation of the line perpendicular to $y + 5 = 3(x - 2)$ that passes through the point $(6, 2)$. Write the equation of the family of lines satisfying the given condition.
12. All lines pass through point $(0, 4)$.
13. All lines are perpendicular to $4x + 3y - 1 = 0$.
14. All lines are parallel to $y - 3 = 4x + 2$.
15. All lines pass through point $(0, -1)$.

Review Answers

1. parallel
2. neither
3. parallel
4. perpendicular
5. parallel
6. perpendicular
7. neither
8. $y = \frac{5}{2}x - \frac{19}{2}$
9. $y = \frac{5}{2}x + 3$
10. $y = -\frac{2}{7}x + \frac{18}{7}$
11. $y = -\frac{1}{3}x + 4$
12. $y = mx + 4$
13. $y = \frac{3}{4}x + b$
14. $y = 4x + b$
15. $y = mx - 1$

CHAPTER

5**Introduction to Functions
and Modeling****Chapter Outline**

- 5.1 FUNCTIONS AS RULES AND TABLES**
 - 5.2 FUNCTIONS AS GRAPHS**
 - 5.3 LINEAR FUNCTION GRAPHS**
 - 5.4 FITTING A LINE TO DATA**
 - 5.5 PREDICTING WITH LINEAR MODELS**
 - 5.6 PROBLEM SOLVING STRATEGIES: USE A LINEAR MODEL**
-

5.1 Functions as Rules and Tables

Learning Objectives

- Identify the domain and range of a function.
- Make a table for a function.
- Write a function rule.
- Represent a real-world situation with a function.

Introduction


A **function** is a rule for relating two or more variables. For example, the price a person pays for phone service may depend on the number of minutes he/she talks on the phone. We would say that the cost of phone service is a *function* of the number of minutes she talks. Consider the following situation.

Josh goes to an amusement park where he pays \$2 per ride.

There is a relationship between the number of rides on which Josh goes and the total cost for the day. To figure out the cost you multiply the number of rides by two. A **function** is the rule that takes us from the number of rides to the cost. Functions usually, *but not always* are rules based on mathematical operations. You can think of a function as a box or a machine that contains a mathematical operation.



A set of numbers is fed into the function box. Those numbers are changed by the given operation into a set of numbers that come out from the opposite side of the box. We can input different values for the number of rides and obtain the cost.




The input is called the **independent variable** because its value can be any possible number. The output results from applying the operation and is called the **dependent variable** because its value depends on the input value.

Often functions are more complicated than the one in this example. Functions usually contain more than one mathematical operation. Here is a situation that is slightly more complicated.

Jason goes to an amusement park where he pays \$8 admission and \$2 per ride.


This function represents the total amount Jason pays. The rule for the function is "multiply the number of rides by 2 and add 8."



We input different values for the number of rides and we arrive at different outputs (costs).



These flow diagrams are useful in visualizing what a function is. However, they are cumbersome to use in practice. We use the following short-hand notation instead.



First, we define the variables.

x = the number of rides Josh goes on

y = the total amount of money Jason paid at the amusement park.

So, x represents the input and y represents the output. The notation: 2 times the number of rides plus 8. This can be written as a function.

$$f(x) = 2x + 8$$

The output is given by the formula $f(x) = 2x + 8$. The notations y and $f(x)$ are used interchangeably but keep in mind that y represents output value and $f(x)$ represents the mathematical operations that gets us from the input to the output.

Identify the Domain and Range of a Function

In the last example, we saw that we can input the number of rides into the function to give us the total cost for going to the amusement park. The set of all values that are possible for the input is called the **domain** of the function. The set of all values that are possible for the output is called the **range** of function. In many situations the **domain** and **range** of a function is the set of all real numbers, but this is not always the case. Let's look at our amusement park example.

Example 1

Find the domain and range of the function that describes the situation:

Jason goes to an amusement park where he pays \$8 admission and \$2 per ride.

Solution

Here is the function that describes this situation.

$$f(x) = 2x + 8 = y$$

In this function, x is the number of rides and y is the total cost. To find the domain of the function, we need to determine which values of x make sense as the input.

- The values have to be zero or positive because Jason can't go on a negative number of rides.
- The values have to be integers because, for example, Jason could not go on 2.25 rides.
- Realistically, there must be a maximum number of rides that Jason can go on because the park closes, he runs out of money, etc. However, since we are not given any information about this we must consider that all non-negative integers could be possible regardless of how big they are.

Answer For this function, the domain is the set of all non-negative integers.

To find the range of the function we must determine what the values of y will be when we apply the function to the input values. The domain is the set of all non-negative integers $(0, 1, 2, 3, 4, 5, 6, \dots)$. Next we plug these values into the function for x .

$$f(x) = 2x + 8 = y$$

Then, $y = 8, 10, 12, 14, 16, 18, 20, \dots$

Answer The range of this function is the set of all even integers greater than or equal to 8.

Example 2

Find the domain and range of the following functions.

a) A ball is dropped from a height and it bounces up to 75% of its original height.

b) $y = x^2$

Solution

a) Lets define the variables:

x = original height

y = bounce height

Here is a function that describes the situation. $y = f(x) = 0.75x$.

The variable x can take any real value greater than zero.

The variable y can also take any real value greater than zero.

Answer The domain is the set of all real numbers greater than zero.

The range is the set of all real numbers greater than zero.

b) Since we dont have a word-problem attached to this equation we can assume that we can use any real number as a value of x .

Since $y = x^2$, the value of y will always be non-negative whether x is positive, negative, or zero.

Answer The domain of this function is all real numbers.

The range of this function is all non-negative real numbers

As we saw, for a function, the variable x is called the **independent variable** because it can be any of the values from the domain. The variable y is called the **dependent variable** because its value depends on x . Any symbols can be used to represent the dependent and independent variables. Here are three different examples.

$$y = f(x) = 3x$$

$$R = f(w) = 3w$$

$$v = f(t) = 3t$$

These expressions all represent the same function. The dependent variable is three times the independent variable. In practice, the symbols used for the independent and dependent variables are based on common usage. For example: t for time, d for distance, v for velocity, etc. The standard symbols to use are y for the dependent variable and x for the independent variable.

A Function:

- Only accepts numbers from the domain.
- For each input, there is exactly one output. All the outputs form the range.

Make a Table For a Function

A table is a very useful way of arranging the data represented by a function. We can match the input and output values and arrange them as a table. Take the amusement park example again.

Jason goes to an amusement park where he pays \$8 admission and \$2 per ride.

We saw that to get from the input to the output we perform the operations $2 \times \text{input} + 8$. For example, we input the values 0, 1, 2, 3, 4, 5, 6, and we obtain the output values 8, 10, 12, 14, 16, 18, 20. Next, we can make the following table of values.

x	y
0	8
1	10
2	12
3	14
4	16
5	18
6	20

A table allows us organize our data in a compact manner. It also provides an easy reference for looking up data, and it gives us a set of coordinate points that we can plot to create a graphical representation of the function.

Example 3

Make a table of values for the following functions.

a) $f(x) = 5x - 9$ Use the following numbers for input values: $-4, -3, -2, -1, 0, 1, 2, 3, 4$.

b) $f(x) = \frac{1}{x}$ Use the following numbers for input values: $-1, -0.5, -0.2, -0.1, -0.01, 0.01, 0.1, 0.2, 0.5, 1$.

Solution

Make a table of values by filling the first column with the input values and the second column with the output values calculated using the given function.

a)

x	$f(x) = 5x - 9 = y$
-4	$5(-4) - 9 = -29$
-3	$5(-3) - 9 = -24$
-2	$5(-2) - 9 = -19$
-1	$5(-1) - 9 = -14$
0	$5(0) - 9 = -9$
1	$5(1) - 9 = -4$
2	$5(2) - 9 = 1$
3	$5(3) - 9 = 6$
4	$5(4) - 9 = 11$

b)

x	$f(x) = \frac{1}{x} = y$
-1	$\frac{1}{-1} = -1$
-0.5	$\frac{1}{-0.5} = -2$
-0.2	$\frac{1}{-0.2} = -5$
-0.1	$\frac{1}{-0.1} = -10$
-0.01	$\frac{1}{-0.01} = -100$
0.01	$\frac{1}{0.01} = 100$
0.1	$\frac{1}{0.1} = 10$
0.2	$\frac{1}{0.2} = 5$
0.5	$\frac{1}{0.5} = 2$
1.0	$\frac{1}{1} = 1$

You are not usually given the input values of a function. These are picked based on the particular function or circumstance. We will discuss how we pick the input values for the table of values throughout this book.

Write a Function Rule

In many situations, we collect data by conducting a survey or an experiment. Then we organize the data in a table of values. Most often, we would like to find the function rule or formula that fits the set of values in the table. This way we can use the rule to predict what could happen for values that are not in the table.

Example 4

Write a function rule for the table.

Number of CDs	2	4	6	8	10
Cost(\$)	24	48	72	86	120

Solution

You pay \$24 for 2 CDs, \$48 for 4 CDs, \$120 for 10 CDs. That means that each CD costs \$12.

We can write the function rule.

Cost = \$12 \times number of CDs or $f(x) = 12x$

Example 5

Write a function rule for the table.

x	-3	-2	-1	0	1	2	3
y	3	2	1	0	1	2	3

Solution

You can see that a negative number turns in the same number but a positive and a non-negative number stays the same. This means that the output values are obtained by applying the absolute value function to the input values:
 $f(x) = |x|$.

Writing a functional rule is probably the hardest thing to do in mathematics. In this book, you will write functional rules mostly for linear relationships which are the simplest type of function.

Represent a Real-World Situation with a Function

Lets look at a few real-world situations that can be represented by a function.

Example 5

Maya has an internet service that currently has a monthly access fee of \$11.95 and a connection fee of \$0.50 per hour. Represent her monthly cost as a function of connection time.

Solution

Define Let x = the number of hours Maya spends on the internet in one month

Let y = Mayas monthly cost


Translate There are two types of cost flat fee of \$11.95 and charge per hour of \$0.50

The total cost = flat fee + hourly fee \times number of hours

Answer The function is $y = f(x) = 11.95 + 0.50x$

Example 6

Alfredo wants a deck build around his pool. The dimensions of the pool are 12 feet \times 24 feet. He does not want to spend more than a set amount and the decking costs \$3 per square foot. Write the cost of the deck as a function of the width of the deck.

**Solution**

Define Let x = width of the deck

Let y = cost of the deck

Make a sketch and label it

Translate You can look at the decking as being formed by several rectangles and squares. We can find the areas of all the separate pieces and add them together:

$$\text{Area of deck} = 12x + 12x + 24x + 24x + x^2 + x^2 + x^2 + x^2 + 72x + 4x^2$$

To find the total cost we multiply the area by the cost per square foot.

Answer $f(x) = 3(72x + 4x^2) = 216x + 12x^2$

Example 7

A cell phone company sells two million phones in their first year of business. The number of phones they sell doubles each year. Write a function that gives the number of phones that are sold per year as a function of how old the company is.

Solution

Define Let x = age of company in years

Let y = number of phones that are sold per year

Make a table

Age (years)	1	2	3	4	5	6	7
Number of phones (millions)	2	4	8	16	32	64	128

Write a function rule

The number of phones sold per year doubles every year. We start with one million the first year:

Year1 :	2 million
Year2 :	$2 \times 2 = 4$ million
Year3 :	$2 \times 2 \times 2 = 8$ million
Year4 :	$2 \times 2 \times 2 \times 2 = 16$ million

We can keep multiplying by two to find the number of phones sold in the next years. You might remember that when we multiply a number by itself several times we can use exponential notation.

$$2 = 2^1$$

$$2 \times 2 = 2^2$$

$$2 \times 2 \times 2 = 2^3$$

In this problem, the exponent represents the age of the company.

Answer $y = f(x) = 2^x$

Review Questions

Identify the domain and range of the following functions.

- Dustin charges \$10 per hour for mowing lawns.
- Maria charges \$25 per hour for tutoring math, with a minimum charge of \$15.
- $f(x) = 15x - 12$
- $f(x) = 2x^2 + 5$
- $f(x) = \frac{1}{x}$
- What is the range of the function $y = x^2 - 5$ when the domain is $-2, -1, 0, 1, 2$?
- What is the range of the function $y = 2x - \frac{3}{4}$ when the domain is $-2.5, 1.5, 5$?
- Angie makes \$6.50 per hour working as a cashier at the grocery store. Make a table of values that shows her earning for input values 5, 10, 15, 20, 25, 30.
- The area of a triangle is given by: $A = \frac{1}{2}bh$. If the height of the triangle is 8 centimeters, make a table of values that shows the area of the triangle for heights 1, 2, 3, 4, 5, and 6 centimeters.
- Make a table of values for the function $f(x) = \sqrt{2x + 3}$ for input values $-1, 0, 1, 2, 3, 4, 5$.

11. Write a function rule for the table

x	3	4	5	6
y	9	16	25	36

12. Write a function rule for the table

hours	0	1	2	3
cost	15	20	25	30

13. Write a function rule for the table

x	0	1	2	3
y	24	12	6	3

14. Write a function that represents the number of cuts you need to cut a ribbon in x number of pieces.
15. Solomon charges a \$40 flat rate and \$25 per hour to repair a leaky pipe. Write a function that represents the total fee charge as a function of hours worked. How much does Solomon earn for a 3 hour job?
16. Rochelle has invested \$2500 in a jewelry making kit. She makes bracelets that she can sell for \$12.50 each. How many bracelets does Rochelle need to make before she breaks even?

Review Answers

- Domain: non-negative rational numbers; Range: non-negative rational numbers.
- Domain: non-negative rational numbers; Range: rational numbers greater than 15.
- Domain: all real numbers; Range: all real numbers.
- Domain: all real numbers; Range: real number greater than or equal to 5.
- Domain: all real numbers except 0; Range: all real numbers except 0.
- $-1, -4, -5$
- $-2, 0, \frac{7}{4}$
-

hours	5	10	15	20	25	30
earnings	\$32.50	\$65	\$97.50	\$130	\$162.50	\$195

- 9.

height (cm)	1	2	3	4	5	6
Area	4	8	12	16	20	24

- 10.

x	-1	0	1	2	3	4	5
y	1	1.73	2.24	2.65	3	3.32	3.6


- $y = x^2$
- $y = 15 + 5x$
- $y = \frac{24}{2x}$
- $f(x) = x - 1$
- $y = 40 + 25x$; \$115
- 200 bracelets

5.2 Functions as Graphs

Learning Objectives

- Graph a function from a rule or table.
- Write a function rule from a graph.
- Analyze the graph of a real world situation.
- Determine whether a relation is a function.

Introduction






We represent functions graphically by plotting points on a **Coordinate Plane** (this is also sometimes called the **Cartesian plane**). The coordinate plane is a grid formed by a horizontal number line and a vertical number line that cross at a point called the **origin**. The origin is point $(0,0)$ and it is the starting location. In order to plot points on the grid, you are told how many units you go right or left and how many units you go up or down from the origin. The horizontal line is called the x -**axis** and the vertical line is called the y -**axis**. The arrows at the end of each axis indicate that the plane continues past the end of the drawing.

From a function, we can gather information in terms of pairs of points. For each value of the independent variable in the domain, the function is used to calculate the value of the dependent variable. We call these pairs of points **coordinate points** or x, y values and they are written as (x, y) .

To graph a coordinate point such as $(4, 2)$ we start at the origin.

Then we move 4 units to the right.


And then we move 2 units up from the last location.



Example 1

Plot the following coordinate points on the Cartesian plane.

- (a) $(5, 3)$
- (b) $(-2, 6)$
- (c) $(3, -4)$
- (d) $(-5, -7)$

**Solution**

We show all the coordinate points on the same plot.

Notice that:

For a positive x value we move to the right.

For a negative x value we move to the left.

For a positive y value we move up.

For a negative y value we move down.

The x -axis and y -axis divide the coordinate plane into four **quadrants**. The quadrants are numbered counter-clockwise starting from the upper right. The plotted point for (a) is in the **First** quadrant, (b) is in the **Second** quadrant, (c) is in the **Fourth** quadrant, and (d) is in the **Third** quadrant.

Graph a Function From a Rule or Table

Once a rule is known or if we have a table of values that describes a function, we can draw a graph of the function. A table of values gives us coordinate points that can be plotted on the Cartesian plane.

Example 2


Graph the function that has the following table of values.

x	-2	-1	0	1	2
y	6	8	10	12	14

Solution

The table gives us five sets of coordinate points $(-2, 6)$, $(-1, 8)$, $(0, 10)$, $(1, 12)$, $(2, 14)$.

To graph the function, we plot all the coordinate points. Since we are not told the domain of the function or the context where it appears we can assume that the domain is the set of all real numbers. To show that the function holds for all values in the domain, we connect the points with a smooth line. Also, we must realize that the line continues infinitely in both directions.


**Example 3**

Graph the function that has the following table of values.

Side of the Square	0	1	2	3	4
Area of the Square	0	1	4	9	16


The table gives us five sets of coordinate points: $(0,0)$, $(1,1)$, $(2,4)$, $(3,9)$, $(4,16)$.

To graph the function, we plot all the coordinate points. Since we are not told the domain of the function, we can assume that the domain is the set of all non-negative real numbers. To show that the function holds for all values in the domain, we connect the points with a smooth curve. The curve does not make sense for negative values of the independent variable so it stops at $x = 0$ but it continues infinitely in the positive direction.

**Example 4**

Graph the function that has the following table of values.

Number of Balloons	12	13	14	15	16
Cost	41	44	47	50	53



This function represents the total cost of the balloons delivered to your house. Each balloon is \$3 and the store delivers if you buy a dozen balloons or more. The delivery charge is a \$5 flat fee.

Solution

The table gives us five sets of coordinate points $(12,41)$, $(13,44)$, $(14,47)$, $(15,50)$, $(16,53)$.

To graph the function, we plot all the coordinate points. Since the x -values represent the number of balloons for 12 balloons or more, the domain of this function is all integers greater than or equal to 12. In this problem, the points are not connected by a line or curve because it does not make sense to have non-integer values of balloons.

In order to draw a graph of a function given the function rule, we must first make a table of values. This will give us a set of coordinate points that we can plot on the Cartesian plane. Choosing the values of the independent variables for the table of values is a skill you will develop throughout this course. When you pick values here are some of the things you should keep in mind.

- Pick only values from the domain of the function.
- If the domain is the set of real numbers or a subset of the real numbers, the graph will be a continuous curve.
- If the domain is the set of integers or a subset of the integers, the graph will be a set of points not connected by a curve.
- Picking integers is best because it makes calculations easier, but sometimes we need to pick other values to capture all the details of the function.
- Often we start with a set of values. Then after drawing the graph, we realize that we need to pick different values and redraw the graph.



Example 5

Graph the following function $f(x) = |x - 2|$

Solution

Make a table of values. Pick a variety of negative and positive integer values for the independent variable. Use the function rule to find the value of the dependent variable for each value of the independent variable. Then, graph each of the coordinate points.

x	$y = f(x) = x - 2 $
-4	$ -4 - 2 = -6 = 6$
-3	$ -3 - 2 = -5 = 5$
-2	$ -2 - 2 = -4 = 4$
-1	$ -1 - 2 = -3 = 3$
0	$ 0 - 2 = -2 = 2$
1	$ 1 - 2 = -1 = 1$
2	$ 2 - 2 = 0 = 0$
3	$ 3 - 2 = 1 = 1$
4	$ 4 - 2 = 2 = 2$
5	$ 5 - 2 = 3 = 3$
6	$ 6 - 2 = 4 = 4$
7	$ 7 - 2 = 5 = 5$
8	$ 8 - 2 = 6 = 6$



It is wise to work with a lot of values when you begin graphing. As you learn about different types of functions, you will find that you will only need a few points in the table of values to create an accurate graph.


Example 6

Graph the following function: $f(x) = \sqrt{x}$

Solution

Make a table of values. We cannot use negative numbers for the independent variable because we can't take the square root of a negative number. The square root doesn't give real answers for negative inputs. The domain is all positive real numbers, so we pick a variety of positive integer values for the independent variable. Use the function rule to find the value of the dependent variable for each value of the independent variable.

x	$y = f(x) = \sqrt{x}$
0	$\sqrt{0} = 0$
1	$\sqrt{1} = 1$
2	$\sqrt{2} \approx 1.41$
3	$\sqrt{3} \approx 1.73$
4	$\sqrt{4} = 2$
5	$\sqrt{5} \approx 2.24$
6	$\sqrt{6} \approx 2.45$
7	$\sqrt{7} \approx 2.65$
8	$\sqrt{8} \approx 2.83$
9	$\sqrt{9} = 3$



Note that the range is all positive real numbers.


Example 7

The post office charges 41cents to send a letter that is one ounce or less and an extra 17cents for any amount up to and including an additional ounce. This rate applies to letters up to 3.5ounces.

Solution

Make a table of values. We cannot use negative numbers for the independent variable because it does not make sense to have negative weight. We pick a variety of positive integer values for the independent variable but we also need to pick some decimal values because prices can be decimals too. This will give us a clear picture of the function. Use the function rule to find the value of the dependent variable for each value of the independent variable.

x	y
0	0
0.2	41
0.5	41
0.8	41
1	41
1.2	58
1.5	58
1.8	58
2	58
2.2	75
2.5	75
2.8	75
3.0	75
3.2	92
3.5	92




Write a Function Rule from a Graph

Sometimes you will need to find the equation or rule of the function by looking at the graph of the function. From a graph, you can read pairs of coordinate points that are on the curve of the function. The coordinate points give values of dependent and independent variables that are related to each other by the rule. However, we must make sure that this rule works for all the points on the curve. In this course you will learn to recognize different kinds of functions. There will be specific methods that you can use for each type of function that will help you find the function rule. For now we will look at some simple examples and find patterns that will help us figure out how the dependent and independent variables are related.

Example 8

The graph to the right shows the distance that an ant covers over time. Find the function rule that shows how distance and time are related to each other.



Solution

We make table of values of several coordinate points to see if we can identify a pattern of how they are related to each other.

Time	0	1	2	3	4	5	6
Distance	0	1.5	3	4.5	6	7.5	9


We can see that for every second the distance increases by 1.5 feet. We can write the function rule as:

$$\text{Distance} = 1.5 \times \text{time}$$

The equation of the function is $f(x) = 1.5x$

Example 9

Find the function rule that describes the function shown in the graph.



Solution:

We make a table of values of several coordinate points to see if we can identify a pattern of how they are related to each other.

x	-4	-3	-2	-1	0	1	2	3	4
y	8	4.5	2	.5	0	.5	2	4.5	8

We notice that the values of y are half of perfect squares. Re-write the table of values as:


x	-4	-3	-2	-1	0	1	2	3	4
y	$\frac{16}{2}$	$\frac{9}{2}$	$\frac{4}{2}$	$\frac{1}{2}$	$\frac{0}{2}$	$\frac{1}{2}$	$\frac{4}{2}$	$\frac{9}{2}$	$\frac{16}{2}$

We can see that to obtain y , we square x and divide by 2.

The function rule is $y = \frac{1}{2}x^2$ and the equation of the function is $f(x) = \frac{1}{2}x^2$.

Example 10

Find the function rule that shows what is the volume of a balloon at different times.



Solution

We make table of values of several coordinate points to see if we can identify a pattern of how they are related to each other.

Time	-1	0	1	2	3	4	5
Volume	10	5	2.5	1.2	0.6	0.3	0.15

We can see that for every day the volume of the balloon is cut in half. Notice that the graph shows negative time. The negative time can represent what happened on days before you started measuring the volume.

$$\text{Day0 : Volume} = 5$$

$$\text{Day1 : Volume} = 5 \cdot \frac{1}{2}$$

$$\text{Day2 : Volume} = 5 \cdot \frac{1}{2} \cdot \frac{1}{2}$$


$$\text{Day3 : Volume} = 5 \cdot \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2}$$

The equation of the function is $f(x) = 5\left(\frac{1}{2}\right)^x$

Analyze the Graph of a Real-World Situation

Graphs are used to represent data in all areas of life. You can find graphs in newspapers, political campaigns, science journals and business presentations.

Here is an example of a graph you might see reported in the news. Most mainstream scientists believe that increased emissions of greenhouse gases, particularly carbon dioxide, are contributing to the warming of the planet. This graph shows how carbon dioxide levels have increased as the world has industrialized.



From this graph, we can find the concentration of carbon dioxide found in the atmosphere in different years.

1900	285 part per million
1930	300 part per million
1950	310 parts per million
1990	350 parts per million

We can find approximate function rules for these types of graphs using methods that you learn in more advanced math classes. The function $f(x) = 0.0066x^2 - 24.9x + 23765$ approximates this graph very well.


Determine Whether a Relation is a Function

You saw that a function is a relation between the independent and the dependent variables. It is a rule that uses the values of the independent variable to give the values of the dependent variable. A function rule can be expressed in words, as an equation, as a table of values and as a graph. All representations are useful and necessary in understanding the relation between the variables. Mathematically, a function is a special kind of relation.

In a function, for each input there is exactly one output.

This usually means that each x -value has only one y -value assigned to it. But, not all functions involve x and y .

Consider the relation that shows the heights of all students in a class. The domain is the set of people in the class and the range is the set of heights. Each person in the class cannot be more than one height at the same time. This relation is a function because for each person there is exactly one height that belongs to him or her.



Notice that in a function, a value in the range can belong to more than one element in the domain, so more than one person in the class can have the same height. The opposite is not possible, one person cannot have multiple heights.

Example 11

Determine if the relation is a function.

- a) $(1, 3), (-1, -2), (3, 5), (2, 5), (3, 4)$
 b) $(-3, 20), (-5, 25), (-1, 5), (7, 12), (9, 2)$
 c)

x	2	4	6	8	10
y	41	44	47	50	53

d)

x	2	1	0	1	2
y	12	10	8	6	4

Solution

The easiest way to figure out if a relation is a function is to look at all the x -values in the list or the table. If a value of x appears more than once and the y -values are different then the relation is not a function.

- a) $(1, 3), (-1, -2), (3, 5), (2, 5), (3, 4)$

You can see that in this relation there are two different y -values that belong to the x -value of 3. This means that this relation is **not** a function.

- b) $(-3, 20), (-5, 25), (-1, 5), (7, 12), (9, 2)$

Each value of x has exactly one y -value. The relation is a function.

c)

x	2	4	6	8	10
y	4	4	4	4	4


Each value of x appears only once. The relation is a function.

d)


x	2	1	0	1	1	2
y	12	10	8	6	4	4

In this relation there are two y -values that belong to the x -value of 2 and two y -values that belong to the x -value of 1. The relation *is not* a function.


When a relation is represented graphically, we can determine if it is a function by using the **vertical line test**. If you can draw a vertical line that crosses the graph in more than one place, then the relation is not a function. Here are some examples.




Not a function. It fails the vertical line test.



A function. No vertical line will cross more than one point on the graph.



A function. No vertical line will cross more than one point on the graph.




Not a function. It fails the vertical line test.

Review Questions

1. Plot the coordinate points on the Cartesian plane.

- (a) $(4, -4)$
- (b) $(2, 7)$
- (c) $(-3, -5)$
- (d) $(6, 3)$
- (e) $(-4, 3)$

2. Give the coordinates for each point in the Cartesian plane.



3. Graph the function that has the following table of values.

(a)

x	-10	-5	0	5	10
y	-3	-0.5	2	4.5	7

(b)

side of cube (in)	0	1	2	3
volume(in^3)	0	1	8	27

(c)

time (hours)	-2	-1	0	1	2
distance from town center (miles)	50	25	0	25	50

4. Graph the following functions.

(a) Brandon is a member of a movie club. He pays a \$50 annual membership and \$8 per movie.

(b) $f(x) = (x - 2)^2$

(c) $f(x) = 3 \cdot 2^x$

5. Determine whether each relation is a function:

(a) $(1, 7), (2, 7), (3, 8), (4, 8), (5, 9)$

(b) $(1, 1), (1, -1), (4, 2), (4, -2), (9, 3), (9, -3)$


(c)

x	-4	-3	-2	-1	0
y	16	9	4	1	0

(d)


Age	20	25	25	30	35
Number of jobs by that age	3	4	7	4	2

6. Write the function rule for each graph.




7. The students at a local high school took The Youth Risk Behavior Survey. The graph below shows the percentage of high school students who reported that they were current smokers. A person qualifies as a current smoker if he/she has smoked one or more cigarettes in the past 30 days. What percentage of high-school students were current smokers in the following years?

- (a) 1991
- (b) 1996
- (c) 2004
- (d) 2005




8. The graph below shows the average life-span of people based on the year in which they were born. This information comes from the National Vital Statistics Report from the Center for Disease Control. What is the average life-span of a person born in the following years?

- (a) 1940
- (b) 1955
- (c) 1980
- (d) 1995





9. The graph below shows the median income of an individual based on his/her number of years of education. The top curve shows the median income for males and the bottom curve shows the median income for females. (Source: US Census, 2003.) What is the median income of a male that has the following years of education?


- (a) 10 years of education
- (b) 17 years of education
- (c) What is the median income of a female that has the same years of education?
- (d) 10 years of education
- (e) 17 years of education



10. Use the vertical line test to determine whether each relation is a function.




Review Answers




2. (a) $(-6, 4)$;
 (b) $(7, 6)$;
 (c) $(-8, -2)$;
 (d) $(4, -7)$;
 (e) $(5, 0)$


3. (a)




(b)




(c)




4. (a)



(b)



(c)



5. (a) function

- (b) not a function
 - (c) function
 - (d) not a function
6. (a) $f(x) = \frac{1}{2}|x|$
(b) $f(x) = \sqrt{x}$
7. (a) 27.5%
(b) 35.6%
(c) 22.2%
(d) 23%
8. (a) 63 years
(b) 69 years
(c) 74 years
(d) 76 years
9. (a) \$19,500
(b) \$56,000
(c) \$10,000
(d) \$35,000
10. (a) function
(b) not a function

5.3 Linear Function Graphs


Learning Objectives

- Recognize and use function notation.
- Graph a linear function.
- Change slope and intercepts of function graphs.
- Analyze graphs of real-world functions.

Introduction Functions

So far we have used the term **function** to describe many of the equations we have been graphing, but the concept of a function is extremely important in mathematics. Not all equations are functions. In order to be a function, the relationship between two variables, x and y , must map each x -value to **exactly one** y -value.

Visually this means the graph of y versus x must pass the **vertical line test** meaning that a vertical line drawn through the graph of the function must never intersect the graph in more than one place.



Use Function Notation

When we write functions we often use the notation $f(x) =$ in place of $y =$. $f(x) =$ is read " f of x ".

Example 1

Rewrite the following equations so that y is a function of x and written $f(x)$.

a. $y = 2x + 5$

b. $y = -0.2x + 7$

c. $x = 4y - 5$

d. $9x + 3y = 6$

Solution

a. Simply replace y with $f(x)$. $f(x) = 2x + 5$

b. $f(x) = -0.2x + 7$

c. Rearrange to isolate y .

$$\begin{array}{ll}
 x = 4y - 5 & \text{Add 5 to both sides.} \\
 x + 5 = 4y & \text{Divide by 4.} \\
 \frac{x + 5}{4} = y & \\
 f(x) = \frac{x + 5}{4} &
 \end{array}$$


d. Rearrange to isolate y .

$$\begin{array}{ll}
 9x + 3y = 6 & \text{Subtract } 9x \text{ from both sides.} \\
 3y = 6 - 9x & \text{Divide by 3.} \\
 y = \frac{6 - 9x}{3} = 2 - 3x & \\
 f(x) = 2 - 3x &
 \end{array}$$

You can think of a function as a machine made up from a number of separate processes. For example, you can look at the function $3x + 2$ and break it down to the following instructions.


- Take a number
- Multiply it by 3
- Add 2

We can visualize these processes like this:



In this case, the number we chose was 2. Multiplied by 3 it becomes 6. When we add 2 our output is 8.

Let's try that again. This time we will put -3 through our machine to get 7.



On the bottom of this process tree you can see what happens when we put the letter n (the variable used to represent *any* number) through the function. We can write the results of these processes.

- $f(2) = 8$
- $f(-3) = -7$
- $f(n) = 3n + 2$

Example 2

A function is defined as $f(x) = 6x - 36$. Evaluate the following:

- $f(2)$
- $f(0)$
- $f(36)$

d. $f(z)$

e. $f(p)$

Solution

a. Substitute $x = 2$ into the function $f(x)f(2) = 6 \cdot 2 - 36 = 12 - 36 = -24$

b. Substitute $x = 0$ into the function $f(x)f(0) = 6 \cdot 0 - 36 = 0 - 36 = -36$

c. Substitute $x = 36$ into the function $f(x)f(36) = 6 \cdot 36 - 36 = 216 - 36 = 180$

d. Substitute $x = z$ into the function $f(x)f(z) = 6z + 36$


e. Substitute $x = p$ into the function $f(x)f(p) = 6p + 36$

Graph a Linear Function

You can see that the notation $f(x) =$ and $y =$ are interchangeable. This means that we can use all the concepts we have learned so far to graph functions.

Example 3

Graph the function $f(x) = \frac{3x+5}{4}$

**Solution**

We can write this function in **slope intercept** form ($y = mx + b$ form).


$$f(x) = \frac{3}{4}x + \frac{5}{4} = 0.75x + 1.25$$

So our graph will have a y -intercept of $(0, 1.25)$ and a slope of 0.75 .

- Remember that this slope **rises** by 3 units for every 4 units we move right.

Example 4

Graph the function $f(x) = \frac{7(5-x)}{5}$

**Solution**

This time we will solve for the x and y intercepts.

To solve for y -intercept substitute $x = 0$.

$$f(0) = \frac{7(5-0)}{5} = \frac{35}{5} = 7$$

To solve for x -intercept substitute use $f(x) = 0$.

$$0 = \frac{7(5-x)}{5}$$

Multiply by 5 and distribute 7.

$$5 \cdot 0 = 35 - 7x$$

Add $7x$ to both sides:

$$7x = 35$$

$$x = 5$$

Our graph has intercepts $(0, 7)$ and $(5, 0)$.

Arithmetic Progressions

You may have noticed that with linear functions, when you increase the x value by one unit, the y value increases by a fixed amount. This amount is equal to the slope. For example, if we were to make a table of values for the function $f(x) = 2x + 3$ we might start at $x = 0$ then add one to x for each row.

x	$f(x)$
0	3
1	5
2	7
3	9
4	11

Look at the values for $f(x)$. They go up by two (the slope) each time. When we consider continually adding a fixed value to numbers, we get sequences like 3, 5, 7, 9, 11.... We call these **arithmetic progressions**. They are characterized by the fact that each number is greater than (or lesser than) than the preceding number by a fixed amount. This amount is called the **common difference**. The common difference can be found by taking two consecutive terms in a sequence and subtracting the first from the second.

Example 5

Find the common difference for the following arithmetic progressions:

- 7, 11, 15, 19...
- 12, 1, -10, -21...
- 7, , 12, , 17...

Solution

a.

$$11 - 7 = 4$$


$$15 - 11 = 4$$

$$19 - 15 = 4.$$

The common difference is 4.

b. $1 - 12 = -11$. The common difference is -11 .

c. There are not two consecutive terms here, but we know that to get the term after 7, we would add the common difference. Then to get to 12, we would add the common difference again. Twice the common difference is $12 - 7 = 5$. So the common difference is 2.5.



Arithmetic sequences and linear functions are very closely related. You just learned that to get to the next term in an arithmetic sequence you add the common difference to last term. We have seen that with linear functions the function increases by the value of the slope every time the x -value is increased by one. As a result, arithmetic sequences and linear functions look very similar.

The graph to the right shows the arithmetic progression $-2, 0, 2, 4, 6 \dots$ with the function $y = 2x - 4$. The fundamental difference between the two graphs is that an arithmetic sequence is **discrete** while a linear function is **continuous**.

- **Discrete** means that the sequence has x values only at distinct points (the 1st term, 2nd term, etc). The domain is not all real numbers (often it is whole numbers).
- **Continuous** means that the function has values for all possible values of x , the integers and also all of the numbers in between. The domain is all real numbers.

We can write a formula for an arithmetic progression. We will define the first term as a_1 and d as the common difference. The sequence becomes the following.


$$a_1, a_1 + d, a_1 + 2d, a_1 + 3d, \dots, a_1 + n \cdot d$$

- To find the second term (a_2) we take the first term (a_1) and add d .
- To find the third term (a_3) we take the first term (a_1) and add $2d$.
- To find the n th term (a_n) we take the first term (a_1) and add $(n - 1)d$.

Analyze Graphs of Real-World Functions

Example 6

Use the diagram below to determine the three decades since 1940 in which the infant mortality rate decreased most.



Let's make a table of the infant mortality rate in the years 1940, 1950, 1960, 1970, 1980, 1990, 2000.

TABLE 5.1: Infant Mortality Rates: United States,

Year	Mortality rate (per 100,000)	change over decade
1940	47	N/A
1950	30	-17
1960	26	-4
1970	20	-6
1980	13	-7
1990	9	-4
2000	7	-2

Solution




The best performing decades were the 1940s (1940 – 1950) with a drop of 17 deaths per 100,000. The 1970s (1970 – 1980) with a drop of 7 deaths per 100,000. The 1960s (1960 – 1970) with a drop of 6 deaths per 100,000.


Lesson Summary

- In order for an equation to be a **function**, the relationship between the two variables, x and y , must map each x -value to *exactly* one y -value, or $y = f(x)$.
- The graph of a function of y versus x must pass the **vertical line test**. Any vertical line will only cross the graph of the function in one place.
- The sequence of $f(x)$ values for a linear function form an arithmetic progression. Each number is greater than (or less than) the preceding number by a fixed amount, or **common difference**.

Review Questions

1. When an object falls under gravity, it gains speed at a constant rate of 9.8 m/s every second. An item dropped from the top of the Eiffel Tower, which is 300 meters tall, takes 7.8 seconds to hit the ground. How fast is it moving on impact?
2. A prepaid phone card comes with \$20 worth of calls on it. Calls cost a flat rate of \$0.16 per minute. Write the value of the card as a function of minutes per calls. Use a function to determine the number of minutes you can make with the card.
3. For each of the following functions evaluate:
 - (a) $f(x) = -2x + 3$
 - (b) $f(x) = 0.7x + 3.2$
 - (c) $f(x) = \frac{5(2-x)}{11}$
 - i. $f(-3)$
 - ii. $f(7)$
 - iii. $f(0)$
 - iv. $f(z)$
4. Determine whether the following could be graphs of **functions**.





5. The roasting guide for a turkey suggests cooking for 100 minutes plus an additional 8 minutes per pound.
- Write a function for the roasting time the given the turkey weight in pounds (x).
 - Determine the time needed to roast a 10 lb turkey.
 - Determine the time needed to roast a 27 lb turkey.
 - Determine the maximum size turkey you could roast in $4\frac{1}{2}$ hours .
6. Determine the missing terms in the following arithmetic progressions.
- $\{-11, 17, \quad, 73\}$
 - $\{2, \quad, -4\}$
 - $\{13, \quad, \quad, 0\}$

Review Answers

- 76.44 m/s
- $f(x) = 2000 - 16x$
125 minutes
- 9
 - 11
 - 3
 - $f(z) = -2z + 3$
 - 1.1
 - 8.1
 - 3.2
 - $f(z) = 0.7z + 3.2$
 - 2.27
 - 2.27
 - 0.909
 - $f(z) = \frac{10}{11} - \frac{5}{11}z$
- yes
 - no
 - no
 - yes
- $f(x) = 8x + 100$
 - 180 min = 3 hrs
 - 316 min = 5 hrs 16 min
 - 21.25 lbs.
- 45
 - 1
 - 9.75, 6.5, 3.25

5.4 Fitting a Line to Data

Learning Objectives

- Make a scatter plot.
- Fit a line to data and write an equation for that line.
- Perform linear regression with a graphing calculator.
- Solve real-world problems using linear models of scattered data.

Introduction

Often in application problems, the relationship between our dependent and independent variables is linear. That means that the graph of the dependent variable vs. independent variable will be a straight line. In many cases we don't know the equation of the line but we have data points that were collected from measurements or experiments. The goal of this section is to show how we can find an equation of a line from data points collected from experimental measurements.

Make a Scatter Plot


A **scatter plot** is a plot of all the ordered pairs in the table. This means that a scatter plot is a relation, and not necessarily a function. Also, the scatter plot is discrete, as it is a set of distinct points. Even when we expect the relationship we are analyzing to be linear, we should not expect that all the points would fit perfectly on a straight line. Rather, the points will be scattered about a straight line. There are many reasons why the data does not fall perfectly on a line such as **measurement error** and **outliers**.

Measurement error is always present as no measurement device is perfectly accurate. In measuring length, for example, a ruler with millimeter markings will be more accurate than a ruler with just centimeter markings.

An **outlier** is an accurate measurement that does not fit with the general pattern of the data. It is a statistical fluctuation like rolling a die ten times and getting the six side all ten times. It can and will happen, but not very often.

Example 1


Make a scatter plot of the following ordered pairs: $(0, 2)$, $(1, 4.5)$, $(2, 9)$, $(3, 11)$, $(4, 13)$, $(5, 18)$, $(6, 19.5)$



Solution


We make a scatter plot by graphing all the ordered pairs on the coordinate axis.

Fit a Line to Data



Notice that the points look like they might be part of a straight line, although they would not fit perfectly on a straight line. If the points were perfectly lined up it would be quite easy to draw a line through all of them and find the equation of that line. However, if the points are scattered, we try to find a line that best fits the data.

You see that we can draw many lines through the points in our data set. These lines have equations that are very different from each other. We want to use the line that is closest to **all** the points on the graph. The best candidate in our graph is the red line **A**. We want to minimize the sum of the distances from the point to the line of fit as you can see in the figure below.



Finding this line mathematically is a complex process and is not usually done by hand. We usually eye-ball the line or find it exactly by using a graphing calculator or computer software such as Excel. The line in the graph above is eye-balled, which means we drew a line that comes closest to all the points in the scatter plot.


When we use the line of best fit we are assuming that there is a continuous linear function that will approximate the discrete values of the scatter plot. We can use this to interpret unknown values.

Write an Equation for a Line of Best Fit

Once you draw the line of best fit, you can find its equation by using two points on the line. Finding the equation of the line of best fit is also called **linear regression**.

Caution: Make sure you don't get caught making a common mistake. In many instances the line of best fit will not pass through many or any of the points in the original data set. This means that you can't just use two random points from the data set. **You need to use two points that are on the line.**

We see that two of the data points are very close to the line of best fit, so we can use these points to find the equation of the line (1, 4.5) and (3, 11).



Start with the slope-intercept form of a line $y = mx + b$.

Find the slope $m = \frac{11 - 4.5}{3 - 1} = \frac{6.5}{2} = 3.25$

Then $y = 3.25x + b$

Plug (3, 11) into the equation. $11 = 3.25(3) + b \Rightarrow b = 1.25$

The equation for the line that fits the data best is $y = 3.25x + 1.25$.

Perform Linear Regression with a Graphing Calculator

Drawing a line of fit can be a good approximation but you can't be sure that you are getting the best results because you are guessing where to draw the line. Two people working with the same data might get two different equations because they would be drawing different lines. To get the most accurate equation for the line, we can use a graphing calculator. The calculator uses a mathematical algorithm to find the line that minimizes the sum of the squares.

Example 2

Use a graphing calculator to find the equation of the line of best fit for the following data $(3, 12), (8, 20), (1, 7), (10, 23), (5, 18), (8,$

Solution

L1	L2	L3	L4
3	12		
8	20		
1	7		
10	23		
5	18		
8			

Step 1 Input the data in your calculator.

Press [STAT] and choose the [EDIT] option.

Input the data into the table by entering the x values in the first column and the y values in the second column.

EDIT	RES	TESTS
1:1-Var	1:1-Var	1:1-Var
2:2-Var	2:2-Var	2:2-Var
3:1-Var	3:1-Var	3:1-Var
4:LinReg(ax+b)	4:LinReg(ax+b)	4:LinReg(ax+b)
5:QuadReg	5:QuadReg	5:QuadReg
6:CubicReg	6:CubicReg	6:CubicReg
7:QuartReg	7:QuartReg	7:QuartReg

Step 2 Find the equation of the line of best fit.

Press [STAT] again use right arrow to select [CALC] at the top of the screen.

Chose option number 4: $LinReg(ax + b)$ and press [ENTER]

The calculator will display $LinReg(ax + b)$

Press [ENTER] and you will be given the a and b values.

LinReg
a=2.01
b=5.94

Here a represents the slope and b represents the y -intercept of the equation. The linear regression line is $y = 2.01x + 5.94$.

STAT	PL1	PL2	PL3
1:On	1:On	1:On	1:On
2:Off	2:Off	2:Off	2:Off
3:Type	3:Type	3:Type	3:Type
4:Format	4:Format	4:Format	4:Format
5:Mark	5:Mark	5:Mark	5:Mark

Step 3 Draw the scatter plot.

To draw the scatter plot press [STATPLOT] [2nd] [Y=].

Choose Plot 1 and press [ENTER].

Press the On option and choose the Type as scatter plot (the one highlighted in black).



Make sure that the X list and Y list names match the names of the columns of the table in Step 1.

Choose the box or plus as the mark since the simple dot may make it difficult to see the points.

Press [**GRAPH**] and adjust the window size so you can see all the points in the scatter plot.



Step 4 Draw the line of best fit through the scatter plot.

Press [**Y=**]

Enter the equation of the line of best fit that you just found $Y_1 = 2.01X + 5.94$

Press [**GRAPH**].

Solve Real-World Problems Using Linear Models of Scattered Data

In a real-world problem, we use a data set to find the equation of the line of best fit. We can then use the equation to predict values of the dependent or independent variables. The usual procedure is as follows.

1. Make a scatter plot of the given data.
2. Draw a line of best fit.
3. Find an equation of a line either using two points on the line or the TI-83/84 calculator.
4. Use the equation to answer the questions asked in the problem.



Example 3


Gal is training for a 5 K race (a total of 5000 meters, or about 3.1 miles). The following table shows her times for each month of her training program. Assume here that her times will decrease in a straight line with time (does that seem like a good assumption?) Find an equation of a line of fit. Predict her running time if her race is in August.

TABLE 5.2:

Month	Month number	Average time (minutes)
January	0	40
February	1	38
March	2	39
April	3	38
May	4	33
June	5	30

Solution

Lets make a scatter plot of Gals running times. The independent variable, x , is the month number and the dependent variable, y , is the running time in minutes. We plot all the points in the table on the coordinate plane.



Draw a line of fit.

Choose two points on the line (0, 41) and (4, 34).


Find the equation of the line.

$$m = \frac{34 - 41}{4 - 0} = -\frac{7}{4} = -1\frac{3}{4}$$

$$y = -\frac{7}{4}x + b$$

$$41 = -\frac{7}{4}(0) + b \Rightarrow b = 41$$

$$y = -\frac{7}{4}x + 41$$



In a real-world problem, the slope and yintercept have a physical significance.

$$\text{Slope} = \frac{\text{number of minutes}}{\text{month}}$$

Since the slope is negative, the number of minutes Gal spends running a 5K race decreased as the months pass. The slope tells us that Gals running time decreases by $\frac{7}{4}$ or 1.75 minutes per month.

The yintercept tells us that when Gal started training, she ran a distance of 5K in 41 minutes , which is just an estimate, since the actual time was 40 minutes .

The problem asks us to predict Gals running time in August. Since June is assigned to month number five, then August will be month number seven. We plug $x = 7$ into the equation of the line of best fit.

$$y = -\frac{7}{4}(7) + 41 = -\frac{49}{4} + 41 = -\frac{49}{4} + \frac{164}{4} = \frac{115}{4} = 28\frac{3}{4}$$

The equation predicts that Gal will be running the 5K race in 28.75 minutes .

In this solution, we eye-balled a line of best fit. Using a graphing calculator, we found this equation for a line of fit $y = -2.2x + 43.7$.

If we plug $x = 7$ in this equation, we get $y = -2.2(7) + 43.7 = 28.3$. This means that Gal ran her race in 28.3 minutes . You see that the graphing calculator gives a different equation and a different answer to the question. The graphing calculator result is more accurate but the line we drew by hand still gives a good approximation to the result.

Example 4

Baris is testing the burning time of BriteGlo candles. The following table shows how long it takes to burn candles of different weights. Assume its a linear relation and we can then use a line to fit the data. If a candle burns for 95 hours , what must be its weight in ounces?




TABLE 5.3: Candle Burning Time Based on Candle Weight

Candle weight (oz)	Time (hours)
2	15
3	20
4	35
5	36
10	80
16	100
22	120
26	180

Solution

Lets make a scatter plot of the data. The independent variable, x , is the candle weight in ounces and the dependent variable, y , is the time in hours it takes the candle to burn. We plot all the points in the table on the coordinate plane.



Then we draw the line of best fit.

Now pick two points on the line $(0,0)$ and $(30,200)$.

Find the equation of the line:

$$m = \frac{200}{30} = \frac{20}{3}$$

$$y = \frac{20}{3}x + b$$

$$0 = \frac{20}{3}(0) + b \Rightarrow b = 0$$

$$y = \frac{20}{3}x$$

In this problem the slope is burning time divided by candle weight. A slope of $\frac{20}{3} = 6\frac{2}{3}$ tells us for each extra ounce of candle weight, the burning time increases by $6\frac{2}{3}$ hours .

A yintercept of zero tells us that a candle of weight 0 oz will burn for 0 hours .

The problem asks for the weight of a candle that burns 95 hours . We are given the value of $y = 95$. We need to use the equation to find the corresponding value of x .

$$y = \frac{20}{3}x \Rightarrow \frac{20}{3}x \Rightarrow x = \frac{285}{20} = \frac{57}{4} = 14\frac{1}{4}$$

A candle that burns 95 hours weighs 14.25 oz.

The graphing calculator gives the linear regression equation as $y = 6.1x + 5.9$ and a result of 14.6 oz.

Notice that we can use the line of best fit to estimate the burning time for a candle of any weight.

Lesson Summary

- A **scatter plot** is a plot of all ordered pairs of experimental measurements.
- **Measurement error** arises from inaccuracies in the measurement device. All measurements of continuous values contain measurement error.
- An **outlier** is an experimental measurement that does not fit with the general pattern of the data.
- For experimental measurements with a linear relationship, you can draw a **line of best fit** which minimizes the distance of each point to the line. Finding the line of best fit is called **linear regression**. A statistics class can teach you the math behind linear regression. For now, you can estimate it visually or use a graphing calculator.

Review Questions

For each data set, draw the scatter plot and find the equation of the line of best fit for the data set by hand.

1. (57, 45) (65, 61) (34, 30) (87, 78) (42, 41) (35, 36) (59, 35) (61, 57) (25, 23) (35, 34)
2. (32, 43) (54, 61) (89, 94) (25, 34) (43, 56) (58, 67) (38, 46) (47, 56) (39, 48)
3. (12, 18) (5, 24) (15, 16) (11, 19) (9, 12) (7, 13) (6, 17) (12, 14)
4. (3, 12) (8, 20) (1, 7) (10, 23) (5, 18) (8, 24) (2, 10)

For each data set, use a graphing calculator to find the equation of the line of best fit.

5. (57, 45) (65, 61) (34, 30) (87, 78) (42, 41) (35, 36) (59, 35) (61, 57) (25, 23) (35, 34)
6. (32, 43) (54, 61) (89, 94) (25, 34) (43, 56) (58, 67) (38, 46) (47, 56) (95, 105) (39, 48)
7. (12, 18) (3, 26) (5, 24) (15, 16) (11, 19) (0, 27) (9, 12) (7, 13) (6, 17) (12, 14)
8. Shiva is trying to beat the samosa eating record. The current record is 53.5 samosas in 12 minutes.

The following table shows how many samosas he eats during his daily practice for the first week of his training. Will he be ready for the contest if it occurs two weeks from the day he started training? What are the meanings of the slope and the y intercept in this problem?

TABLE 5.4:

Day	No. of Samosas
1	30
2	34
3	36
4	36
5	40
6	43
7	45

9. Nitisha is trying to find the elasticity coefficient of a Superball. She drops the ball from different heights and measures the maximum height of the resulting bounce. The table below shows her data. Draw a scatter plot and find the equation. What is the initial height if the bounce height is 65 cm? What are the meanings of the slope and the yintercept in this problem?

TABLE 5.5:

Initial height (cm)	Bounce height (cm)
30	22
35	26
40	29
45	34
50	38
55	40
60	45
65	50
70	52


10. The following table shows the median California family income from 1995 to 2002 as reported by the US Census Bureau. Draw a scatter plot and find the equation. What would you expect the median annual income of a Californian family to be in year 2010? What are the meanings of the slope and the yintercept in this problem?

TABLE 5.6:


Year	Income
1995	53807
1996	55217
1997	55209
1998	55415
1999	63100
2000	63206
2001	63761
2002	65766

Review Answers


1. $y = 0.9x - 0.8$




2. $y = 1.05x + 6.1$




3. $y = -0.86x + 24.3$




4. $y = 2x + 6$




5. $y = .8x + 3.5$




6. $y = .96x + 10.83$



7. $y = -.8x + 25$



8. $y = 2.5x + 27.5$



Solution

$y = 57.5$. Shiva will beat the record.

5.5 Predicting with Linear Models

Learning Objectives

- Collect and organize data.
- Interpolate using an equation.
- Extrapolate using an equation.
- Predict using an equation.

Introduction

Numerical information appears in all areas of life. You can find it in newspapers, magazines, journals, on the television or on the internet. In the last section, we saw how to find the equation of a line of best fit and how to use this equation to make predictions. The line of best fit is a good method if the relationship between the dependent and the independent variables is linear. In this section, you will learn other methods that help us estimate data values. These methods are useful in linear and non-linear relationships equally. The methods you will learn are **linear interpolation** which is useful if the information you are looking for is between two known points and **linear extrapolation** which is useful for estimating a value that is either less than or greater than the known values.

Collect and Organize Data

Data can be collected through **surveys** or **experimental measurements**.

Surveys are used to collect information about a population. Surveys of the population are common in political polling, health, social science and marketing research. A survey may focus on opinions or factual information depending on its purpose.

Experimental measurements are data sets that are collected during experiments.

The information collected by the US Census Bureau (www.census.gov) or the Center for Disease Control (www.cdc.gov) are examples of data gathered using surveys. The US Census Bureau collects information about many aspects of the US population. The census takes place every ten years and it polls the population of the United States.

Lets say we are interested in how the median age for first marriages has changed during the 20th century.

Example 1

Median age at first marriage

The US Census gives the following information about the median age at first marriage for males and females.



In 1890, the median age for males was 26.1 and for females it was 22.0.

In 1900, the median age for males was 25.9 and for females it was 21.9.

In 1910, the median age for males was 25.1 and for females it was 21.6.

In 1920, the median age for males was 24.6 and for females it was 21.2.

In 1930, the median age for males was 24.3 and for females it was 21.3.

In 1940, the median age for males was 24.3 and for females it was 21.5.

In 1950, the median age for males was 22.8 and for females it was 20.3.

In 1960, the median age for males was 22.8 and for females it was 20.3.

In 1970, the median age for males was 23.2 and for females it was 20.8.

In 1980, the median age for males was 24.7 and for females it was 22.0.

In 1990, the median age for males was 26.1 and for females it was 23.9.

In 2000, the median age for males was 26.8 and for females it was 25.1.

This is not a very efficient or clear way to display this information. Some better options are organizing the data in a table or a scatter plot.


A table of the data would look like this.

TABLE 5.7: Median Age of Males and Females at First Marriage by Year

Year	Median Age of Males	Median Age of Females
1890	26.1	22.0
1900	25.9	21.9
1910	25.1	21.6
1920	24.6	21.2
1930	24.3	21.3
1940	24.3	21.5
1950	22.8	20.3
1960	22.8	20.3
1970	23.2	20.8
1980	24.7	22.0
1990	26.1	23.9
2000	26.8	25.1

A scatter plot of the data would look like this.

Median Age of Males and Females at First Marriage by Year



The Center for Disease Control collects information about the health of the American people and behaviors that might lead to bad health. The next example shows the percent of women that smoke during pregnancy.



Example 2

Pregnant women and smoking

The CDC has the following information.

In the year 1990, 18.4 percent of pregnant women smoked.

In the year 1991, 17.7 percent of pregnant women smoked.

In the year 1992, 16.9 percent of pregnant women smoked.

In the year 1993, 15.8 percent of pregnant women smoked.

In the year 1994, 14.6 percent of pregnant women smoked.

In the year 1995, 13.9 percent of pregnant women smoked.

In the year 1996, 13.6 percent of pregnant women smoked.

In the year 2000, 12.2 percent of pregnant women smoked.

In the year 2002, 11.4 percent of pregnant women smoked.

In the year 2003, 10.4 percent of pregnant women smoked.

In the year 2004, 10.2 percent of pregnant women smoked.

Lets organize this data more clearly in a table and in a scatter plot.


Here is a table of the data.

TABLE 5.8: Percent of Pregnant Women Smokers by Year

Year	Percent of pregnant women smokers
1990	18.4
1991	17.7
1992	16.9
1993	15.8
1994	14.6
1995	13.9
1996	13.6
2000	12.2
2002	11.4
2003	10.4
2004	10.2

Here is a scatter plot of the data.

Percent of Pregnant Women Smokers by Year



Interpolate Using an Equation

Linear interpolation is often used to fill the gaps in a table. Example one shows the median age of males and females at the time of their first marriage. However, the information is only available at ten year intervals. We know the median age of marriage every ten years from 1890 to 2000, but we would like to estimate the median age of marriage for the years in between. Example two gave us the percentage of women smoking while pregnant. But,

there is no information collected for 1997, 1998, 1999 and 2001 and we would like to estimate the percentage for these years. Linear interpolation gives you an easy way to do this.

The strategy for linear interpolation is to use a straight line to connect the known data points (we are assuming that the data would be continuous between the two points) on either side of the unknown point. Then we use that equation to estimate the value we are looking for.


Example 3

Estimate the median age for the first marriage of a male in the year 1946.

TABLE 5.9: Median Age of Males and Females at First Marriage by Year (excerpt)

Year	Median age of males	Median age of females
...
1940	24.3	21.5
1950	22.8	20.3
...

The table to the left shows only the data for the years 1950 and 1940 because we want to estimate a data point between these two years.



We connect the two points on either side of 1946 with a straight line and find its equation.

$$\begin{aligned} \text{Slope} \quad m &= \frac{22.8 - 24.3}{1950 - 1940} = \frac{-1.5}{10} = -0.15 \\ y &= -0.15x + b \\ 24.3 &= -0.15(1940) + b \\ b &= 315.3 \\ \text{Equation} \quad y &= -0.15x + 315.3 \end{aligned}$$

To estimate the median age of marriage of males in year 1946 we plug $x = 1946$ in the equation.

$$y = -0.15(1946) + 315.3 = 23.4 \text{ years old}$$

Example 4

Estimate the percentage of pregnant women that were smoking in the year 1998.


TABLE 5.10: Percent of Pregnant Women Smokers by Year (excerpt)

Year	Percent of Pregnant Women Smokers
...	...
1996	13.6
2000	12.2
...	...

The table to the left shows only the data for year 1996 and 2000 because we want to estimate a data point between these two years.

Connect the points on either side of 1998 with a straight line and find the equation of that line.

Slope	$m = \frac{12.2 - 13.6}{2000 - 1996} = \frac{-1.4}{4} = -0.35$ $y = -0.35x + b$ $12.2 = -0.35(2000) + b$ $b = 712.2$
Equation	$y = -0.35x + 712.2$



To estimate the percentage of pregnant women who smoked in year 1998 we plug $x = 1998$ into the equation.

$$y = -0.35(1998) + 712.2 = 12.9\%$$

For non-linear data, linear interpolation is often not accurate enough for our purposes. If the points in the data set change by a large amount in the interval in which you are interested, then linear interpolation may not give a good estimate. In that case, it can be replaced by **polynomial interpolation** which uses a curve instead of a straight line to estimate values between points.

Extrapolating: How to Use it and When Not to Use it

Linear extrapolation can help us estimate values that are either higher or lower than the range of values of our data set. The strategy is similar to linear interpolation. However you only use a subset of the data, rather than all of the data. For linear data, you are ALWAYS more accurate by using the best fit line method of the previous section. For non-linear data, it is sometimes useful to extrapolate using the last two or three data points in order to estimate a y -value that is higher than the data range. To estimate a value that is higher than the points in the data set, we connect the last two data points with a straight line and find its equation. Then we can use this equation to estimate the value we are trying to find. To estimate a value that is lower than the points in the data set, we follow the same procedure. But we use the first two points of our data instead.

Example 5

Winning Times

The winning times for the women's 100 meter race are given in the following table³. Estimate the winning time in the year 2010. Is this a good estimate?




TABLE 5.11:

Winner	Country	Year	Time (seconds)
Mary Lines	UK	1922	12.8
Leni Schmidt	Germany	1925	12.4
Gerturd Glasitsch	Germany	1927	12.1
Tollien Schuurman	Netherlands	1930	12.0
Helen Stephens	USA	1935	11.8
Lulu Mae Hymes	USA	1939	11.5
Fanny Blankers-Koen	Netherlands	1943	11.5
Marjorie Jackson	Australia	1952	11.4
Vera Krepkina	Soviet Union	1958	11.3
Wyomia Tyus	USA	1964	11.2
Barbara Ferrell	USA	1968	11.1
Ellen Strophal	East Germany	1972	11.0
Inge Helten	West Germany	1976	11.0
Marlies Gohr	East Germany	1982	10.9
Florence Griffith Joyner	USA	1988	10.5

Solution

We start by making a scatter plot of the data. Connect the last two points on the graph and find the equation of the line.

Winning Times for the Womens 100 meter Race by Year

Slope

$$m = \frac{10.5 - 10.9}{1988 - 1982} = \frac{-0.4}{6} = -0.067$$

$$y = -0.067x + b$$

$$10.5 = -0.067(1988) + b$$

$$b = 143.7$$

Equation $y = -0.067x + 143.7$

The winning time in year 2010 is estimated to be:

$$y = -0.067(2010) + 143.7 = \underline{9.03 \text{ seconds}}$$

³ Source: http://en.wikipedia.org/wiki/World_Record_progression_100_m_women.


How accurate is this estimate? It is likely that it's not very accurate because 2010 is a long time from 1988. This example demonstrates the weakness of linear extrapolation. Estimates given by linear extrapolation are never as good as using the equation from the best fit line method. In this particular example, the last data point clearly does not fit in with the general trend of the data so the slope of the extrapolation line is much steeper than it should be.

As a historical note, the last data point corresponds to the winning time for Florence Griffith Joyner in 1988. After her race, she was accused of using performance-enhancing drugs but this fact was never proven. In addition, there is a question about the accuracy of the timing because some officials said that the tail wind was not accounted for in this race even though all the other races of the day were impacted by a strong wind.

Predict Using an Equation

Linear extrapolation was not a good method to use in the last example. A better method for estimating the winning time in 2010 would be the use of linear regression (i.e. *best fit line method*) that we learned in the last section. Lets apply that method to this problem.

Winning Times for the Womens 100 meter Race by Year



We start by drawing the line of best fit and finding its equation. We use the points (1982, 10.9) and (1958, 11.3).

The equation is $y = -0.017x + 43.9$

In year 2010, $y = -0.017(2010) + 43.9 = 9.73$ seconds

This shows a much slower decrease in winning times than linear extrapolation. This method (fitting a line to all of the data) is always more accurate for linear data and approximate linear data. However, the line of best fit in this case will not be useful in the future. For example, the equation predicts that around the year 2582 the time will be about zero seconds, and in years that follow the time will be negative!

Lesson Summary

- A **survey** is a method of collecting information about a population.
- **Experimental measurements** are data sets that are collected during experiments.
- **Linear interpolation** is used to estimate a data value between two experimental measurements. To do so, compute the line through the two adjacent measurements, then use that line to estimate the intermediate value.
- **Linear extrapolation** is used to estimate a data value either above or below the experimental measurements. Again, find the line defined by the two closest points and use that line to estimate the value.
- The **most accurate method** of estimating data values from a linear data set is to perform linear regression and estimate the value from the best-fit line.

Review Questions

1. Use the data from Example one (*Median age at first marriage*) to estimate the age at marriage for females in 1946. Fit a line, by hand, to the data before 1970.
2. Use the data from Example one (*Median age at first marriage*) to estimate the age at marriage for females in 1984. Fit a line, by hand, to the data from 1970 on in order to estimate this accurately.
3. Use the data from Example one (*Median age at first marriage*) to estimate the age at marriage for males in 1995. Use linear interpolation between the 1990 and 2000 data points.
4. Use the data from Example two (*Pregnant women and smoking*) to estimate the percent of pregnant smokers in 1997. Use linear interpolation between the 1996 and 2000 data points.
5. Use the data from Example two (*Pregnant women and smoking*) to estimate the percent of pregnant smokers in 2006. Use linear extrapolation with the final two data points.

6. Use the data from Example five (*Winning times*) to estimate the winning time for the female 100 meter race in 1920. Use linear extrapolation because the first two or three data points have a different slope than the rest of the data.
7. The table below shows the highest temperature vs. the hours of daylight for the 15th day of each month in the year 2006 in San Diego, California. Estimate the high temperature for a day with 13.2 hours of daylight using linear interpolation.

TABLE 5.12:

Hours of daylight	High temperature (F)
10.25	60
11.0	62
12	62
13	66
13.8	68
14.3	73
14	86
13.4	75
12.4	71
11.4	66
10.5	73
10	61

- 8.
9. Using the table above to estimate the high temperature for a day with 9 hours of daylight using linear extrapolation. Is the prediction accurate? Find the answer using line of best fit.

Review Answers

1. About 21 years
2. 22.8 years
3. 26.5 years
4. 13.25 percent
5. 9.8 percent
6. 13.1 seconds
7. 70.5 F
8. 65 F. Prediction is not very good since we expect cooler temperatures for less daylight hours. The best fit line method of linear regression predicts 58.5 F.

5.6 Problem Solving Strategies: Use a Linear Model

Learning Objectives

- Read and understand given problem situations.
- Develop and apply the strategy: use a linear model.
- Plan and compare alternative approaches to solving problems.
- Solve real-world problems using selected strategies as part of a plan.

Introduction

In this chapter, we have been estimating values using straight lines. When we use linear interpolation, linear extrapolation or predicting results using a line of best fit, it is called **linear modeling**. In this section, we will look at a few examples where data sets occurring in real-world problems can be modeled using linear relationships. From previous sections remember our problem solving plan:.

Step 1

Understand the problem

Read the problem carefully. Once the problem is read, list all the components and data that are involved. This is where you will be assigning your variables

Step 2

Devise a plan Translate

Come up with a way to solve the problem. Set up an equation, draw a diagram, make a chart or construct a table as a start to solving your problem.

Step 3

Carry out the plan Solve

This is where you solve the equation you came up with in Step 2.

Step 4

Look Check and Interpret

Check to see if you used all your information and that the answer makes sense.

Example 1

Dana heard something very interesting at school. Her teacher told her that if you divide the circumference of a circle by its diameter you always get the same number. She tested this statement by measuring the circumference and diameter of several circular objects. The following table shows her results.

From this data, estimate the circumference of a circle whose diameter is 12 inches . What about 25 inches ? 60 inches ?

TABLE 5.13: Diameter and Circumference of Various Objects

Object	Diameter (inches)	Circumference (inches)
Table	53	170
Soda can	2.25	7.1

TABLE 5.13: (continued)

Object	Diameter (inches)	Circumference (inches)
Cocoa tin	4.2	12.6
Plate	8	25.5
Straw	.25	1.2
Propane tank	13.3	39.6
Hula Hoop	34.25	115

Solution

Lets use the problem solving plan.

Step 1


We define our variables.

x = diameter of the circle in inches

y = circumference of the circle in inches

We want to know the circumference when the diameter is 12, 25 or 60 inches .

Step 2 We can find the answers either by using the line of best fit or by using linear interpolation or extrapolation. We start by drawing the scatter plot of the data.

**Step 3** Line of best fit

Estimate a line of best fit on the scatter plot.

Find the equation using points $(.25, 1.2)$ and $(8, 25.5)$.

Slope

$$m = \frac{25.5 - 1.2}{8 - .25} = \frac{24.3}{7.75} = 3.14$$

$$y = 3.14x + b$$

$$1.2 = 3.14(.25) + b \Rightarrow b = 0.42$$

Equation

$$y = 3.14x + 0.42$$

$$\text{Diameter} = 12 \text{ inches} \Rightarrow y = 3.14(12) + 0.42 = \underline{38.1 \text{ inches}}$$

$$\text{Diameter} = 25 \text{ inches} \Rightarrow y = 3.14(25) + 0.42 = \underline{78.92 \text{ inches}}$$

$$\text{Diameter} = 60 \text{ inches} \Rightarrow y = 3.14(60) + 0.42 = \underline{188.82 \text{ inches}}$$

In this problem the slope = 3.14. This number should be very familiar to you it is the number rounded to the hundredths place. Theoretically, the circumference of a circle divided by its diameter is always the same and it equals 3.14 or π .

You are probably more familiar with the formula $C = \pi \cdot d$.

Note: The calculator gives the line of best fit as $y = 3.25x - 0.57$, so we can conclude that we luckily picked two values that gave the correct slope of 3.14. Our line of best fit shows that there was more measurement error in other points.

Step 4 Check and Interpret

The circumference of a circle is πd and the diameter is simply d . If we divide the circumference by the diameter we will get π . The slope of the line is 3.14, which is very close to the exact value of π . There is some error in the estimation because we expect the y -intercept to be zero and it is not.

The reason the line of best fit method works the best is that the data is very linear. All the points are close to the straight line but there is some slight measurement error. The line of best fit averages the error and gives a good estimate of the general trend.

Note: The linear interpolation and extrapolation methods give estimates that aren't as accurate because they use only two points in the data set. If there are measurement errors in the points that are being used, then the estimates will lose accuracy. Normally, it is better to compute the line of best fit with a calculator or computer.

Example 2

A cylinder is filled with water to a height of 73 centimeters. The water is drained through a hole in the bottom of the cylinder and measurements are taken at two second intervals. The table below shows the height of the water level in the cylinder at different times.

a) Find the water level at 15 seconds.

b) Find the water level at 27 seconds.

Water Level in Cylinder at Various Times

TABLE 5.14: Water Level in Cylinder at Various Times

Time (seconds)	Water level (cm)
0.0	73
2.0	63.9
4.0	55.5
6.0	47.2
8.0	40.0
10.0	33.4
12.0	27.4
14.0	21.9
16.0	17.1
18.0	12.9
20.0	9.4
22.0	6.3
24.0	3.9

TABLE 5.14: (continued)

Time (seconds)	Water level (cm)
26.0	2.0
28.0	0.7
30.0	0.1

Solution

Lets use the problem solving plan.

Step 1


Define our variables

x = time in seconds

y = water level in centimeters

We want to know the water level at time 15, 27 and -5 seconds.

Step 2 We can find the answers either by using the line of best fit or by using linear interpolation or extrapolation. We start by drawing the scatter plot of the data.

**Step 3 Method 1** Line of best fit

Draw an estimate of the line of best fit on the scatter plot. Find the equation using points (6, 47.2) and (24, 3.9).

Slope

$$m = \frac{3.9 - 47.2}{24 - 6} = \frac{-43.3}{18} = -2.4$$

$$y = -2.4x + b$$

$$47.2 = -2.4(6) + b \Rightarrow b = 61.6$$

Equation

$$y = -2.4x + 61.6$$


$$\text{Time} = 15 \text{ seconds} \Rightarrow y = -2.4(15) + 61.6 = \underline{25.6\text{cm}}$$

$$\text{Time} = 27 \text{ seconds} \Rightarrow y = -2.4(27) + 61.6 = \underline{-3.2 \text{ cm}}$$

The line of best fit does not show us accurate estimates for the height. The data points do not appear to fit a linear trend so the line of best fit is close to very few data points.

Method 2: Linear interpolation or linear extrapolation.

We use linear interpolation to find the water level for the times 15 and 27 seconds, because these points are between the points we know.



Time = 15 seconds

Connect points (14, 21.9) and (16, 17.1) and find the equation of the straight line.

$$m = \frac{17.1 - 21.9}{16 - 14} = \frac{-4.8}{2} = -2.4 \Rightarrow -2.4y = -2.4x + b \Rightarrow 21.9 = -2.4(14) + b \Rightarrow b = 55.5$$

Equation $y = -2.4x + 55.5$

Plug in $x = 15$ and obtain $y = -2.4(15) + 55.5 = 19.5 \text{ cm}$

Time = 27 seconds

Connect points (26, 2) and (28, 0.7) and find the equation of the straight line.

$$m = \frac{0.7 - 2}{28 - 26} = \frac{-1.3}{2} = -.65$$

$$y = -.65x + b \Rightarrow 2 = -.65(26) + b \Rightarrow b = 18.9$$

Equation $y = -.65x + 18.9$

Plug in $x = 27$ and obtain $y = -.65(27) + 18.9 = 1.35$ cm

We use linear extrapolation to find the water level for time -5 seconds because this point is smaller than the points in our data set.

Step 4 Check and Interpret

In this example, the linear interpolation and extrapolation method gives better estimates of the values that we need to solve the problem. Since the data is not linear, the line of best fit is not close to many of the points in our data set. The linear interpolation and extrapolation methods give better estimates because we do not expect the data to change greatly between the points that are known.

Lesson Summary

- Using linear interpolation, linear extrapolation or prediction using a line of best fit is called **linear modeling**.
- The four steps of the **problem solving plan** are:
 1. Understand the problem
 2. Devise a plan Translate
 3. Carry out the plan Solve
 4. Look Check and Interpret

Review Questions

The table below lists the predicted life expectancy based on year of birth (US Census Bureau).

Use this table to answer the following questions.

1. Make a scatter plot of the data
2. Use a line of best fit to estimate the life expectancy of a person born in 1955.
3. Use linear interpolation to estimate the life expectancy of a person born in 1955.
4. Use a line of best fit to estimate the life expectancy of a person born in 1976.
5. Use linear interpolation to estimate the life expectancy of a person born in 1976.
6. Use a line of best fit to estimate the life expectancy of a person born in 2012.
7. Use linear extrapolation to estimate the life expectancy of a person born in 2012.
8. Which method gives better estimates for this data set? Why?

TABLE 5.15:

Birth Year	Life expectancy in years
1930	59.7
1940	62.9
1950	68.2
1960	69.7
1970	70.8
1980	73.7
1990	75.4
2000	77

- 9.
10. The table below lists the high temperature for the first day of the month for year
11. 2006
12. in San Diego, California (Weather Underground). Use this table to answer the following questions.

13. Draw a scatter plot of the data
14. Use a line of best fit to estimate the temperature in the middle of the 4th month (month 4.5).
15. Use linear interpolation to estimate the temperature in the middle of the 4th month (month 4.5).
16. Use a line of best fit to estimate the temperature for month 13 (January 2007).
17. Use linear extrapolation to estimate the temperature for month 13 (January 2007).
18. Which method gives better estimates for this data set? Why?


TABLE 5.16:

Month number	Temperature (F)
1	63
2	66
3	61
4	64
5	71
6	78
7	88
8	78
9	81
10	75
11	68
12	69


19.

Review Answers

1. Equation of line of best fit using points (1940, 62.9) and (1990, 75.4) $y = .25x - 422.1$.



2. 66.7 years
3. $y = .15x - 224.3$, 69.0 years
4. 71.9 years
5. $y = .29x - 500.5$, 72.5 years
6. 80.9 years
7. $y = .16x - 243$, 78.9 years
8. A line of best fit gives better estimates because data is linear.
9. Equation of line of best fit using points (2, 66) and (10, 75) $y = 1.125x + 63.75$



10. 68.8 F
11. $y = 7x + 36$, 67.5 F
12. 78.4 F
13. $y = x + 57$, 70 F
14. Linear interpolation and extrapolation give better estimates because data is not linear.

CHAPTER 6**Linear Inequalities****Chapter Outline**

- 6.1 SOLVING ONE-STEP INEQUALITIES**
 - 6.2 MULTI-STEP INEQUALITIES**
 - 6.3 COMPOUND INEQUALITIES**
 - 6.4 ABSOLUTE VALUE EQUATIONS**
 - 6.5 ABSOLUTE VALUE INEQUALITIES**
 - 6.6 LINEAR INEQUALITIES IN TWO VARIABLES**
-

6.1 Solving One-Step Inequalities

Learning Objectives

- Write and graph inequalities in one variable on a number line.
- Solve an inequality using addition.
- Solve an inequality using subtraction.
- Solve an inequality using multiplication.
- Solve an inequality using division.
- Multiply or divide an inequality by a negative number.

Introduction

Inequalities are similar to equations in that they show a relationship between two expressions. We solve and graph inequalities in a similar way to equations. However, there are some differences that we will talk about in this chapter. The main difference is that for linear inequalities the answer is an interval of values whereas for a linear equation the answer is most often just one value.


When writing inequalities we use the following symbols

$>$	is greater than
\geq	is greater than or equal to
$<$	is less than
\leq	is less than or equal to

Write and Graph Inequalities in One Variable on a Number Line

Let's start with the simple inequality $x > 3$

We read this inequality as “ x is greater than 3”. The solution is the set of all real numbers that are greater than three. We often represent the solution set of an inequality by a number line graph.



Consider another simple inequality $x \leq 4$

We read this inequality as “ x is less than or equal to 4”. The solution is the set of all real numbers that equal four or less than four. We graph this solution set on the number line.



In a graph, we use an empty circle for the endpoint of a strict inequality ($x > 3$) and a filled circle if the equal sign is included ($x \leq 4$).

Example 1

Graph the following inequalities on the number line.

a) $x < -3$


b) $x \geq 6$

c) $x > 0$


d) $x \leq 8$

Solution


a) The inequality $x < -3$ represents all real numbers that are less than -3 . The number -3 is not included in the solution and that is represented by an open circle on the graph.



b) The inequality $x \geq 6$ represents all real numbers that are greater than or equal to six. The number six is included in the solution and that is represented by a closed circle on the graph.



c) The inequality $x > 0$ represents all real numbers that are greater than zero. The number zero is not included in the solution and that is represented by an open circle on the graph.




d) The inequality $x \leq 8$ represents all real numbers that are less than or equal to eight. The number eight is included in the solution and that is represented by a closed circle on the graph.


**Example 2**

Write the inequality that is represented by each graph.

a)




b)



c)



d)

**Solution:**

$$d) x \geq c) x < 65 \quad b) x > 540 \quad a) x \leq -12$$

Inequalities appear everywhere in real life. Here are some simple examples of real-world applications.


Example 3

Write each statement as an inequality and graph it on the number line.


- You must maintain a balance of at least \$2500 in your checking account to get free checking.
- You must be at least 48 inches tall to ride the “Thunderbolt” Rollercoaster.
- You must be younger than 3 years old to get free admission at the San Diego Zoo.
- The speed limit on the interstate is 65 miles per hour.

Solution:


a) The inequality is written as $x \geq 2500$. The words “at least” imply that the value of \$2500 is included in the solution set.



b) The inequality is written as $x \geq 48$. The words “at least” imply that the value of 48 inches is included in the solution set.



c) The inequality is written as $x < 3$.



d) Speed limit means the highest allowable speed, so the inequality is written as $x \leq 65$.

**Solve an Inequality Using Addition**

To solve an inequality we must isolate the variable on one side of the inequality sign. To isolate the variable, we use the same basic techniques used in solving equations. For inequalities of this type:

$$x - a < b \text{ or } x - a > b$$

We isolate the x by adding the constant a to both sides of the inequality.

Example 4

Solve each inequality and graph the solution set.

$$a) x - 3 < 10$$

$$b) x - 1 > -10$$

c) $x - 1 \leq -5$


d) $x - 20 \geq 14$

Solution:

a)

To solve the inequality
Add 3 to both sides of the inequality.
Simplify


$$\begin{aligned}x - 3 &< 10 \\x - 3 + 3 &< 10 + 3 \\x &< 13\end{aligned}$$



b)

To solve the inequality
Add 12 to both sides of the inequality
Simplify


$$\begin{aligned}x - 1 &> -10 \\x - 12 + 12 &> -10 + 12 \\x &> 2\end{aligned}$$



c)

To solve the inequality
Add 1 to both sides of the inequality
Simplify to obtain

$$\begin{aligned}x - 1 &\leq -5 \\x - 1 + 1 &\leq -5 + 1 \\x &\leq -4\end{aligned}$$



d)

To solve the inequality
Add 20 to both sides of the inequality :
Simplify

$$\begin{aligned}x - 20 &\leq 14 \\x - 20 + 20 &\leq 14 + 20 \\x &\leq 34\end{aligned}$$

**Solve an Inequality Using Subtraction**

For inequalities of this type:

$x + 1 < b$ or $x + 1 > b$

We isolate the x by subtracting the constant a on both sides of the inequality.

Example 5

Solve each inequality and graph the solution set.

a) $x + 2 < 7$

b) $x + 8 \leq -7$

c) $x + 4 > 13$

d) $x + 5 \geq \frac{3}{4}$

Solution:


a)

To solve the inequality	$x + 2 < 7$
Subtract 2 on both sides of the inequality	$x + 2 - 2 < 7 - 2$
Simplify to obtain	$x < 5$




b)

To solve the inequality	$x + 8 \leq -7$
Subtract 8 on both sides of the inequality	$x + 8 - 8 \leq -7 - 8$
Simplify to obtain :	$x \leq -15$




c)

To solve the inequality	$x + 4 > 13$
Subtract 4 on both sides of the inequality	$x + 4 - 4 > 13 - 4$
Simplify	$x > 9$



d)

To solve the inequality	$x + 5 \geq \frac{3}{4}$
Subtract 5 on both sides of the inequality	$x + 5 - 5 \geq -\frac{3}{4} - 5$
Simplify to obtain :	$x \geq -5\frac{3}{4}$



Solve an Inequality Using Multiplication

Consider the problem

$$\frac{x}{5} \leq 3$$

To find the solution we multiply both sides by 5.

$$5 \cdot \frac{x}{5} \leq 3 \cdot 5$$

We obtain

$$x \leq 15$$

The answer of an inequality can be expressed in four different ways:

1. **Inequality notation** The answer is simply expressed as $x < 15$.
2. **Set notation** The answer is $\{x | x < 15\}$. You read this as “the set of all values of x , such that x is a real number less than 15”.
3. **Interval notation** uses brackets to indicate the range of values in the solution.

The interval notation solution for our problem is $(-\infty, 15)$. Interval notation also uses the concept of **infinity** ∞ and **negative infinity** $-\infty$. Round or **open brackets** “(” and “)” indicate that the number next to the bracket is not included in the solution set. Square or **closed brackets** “[” and “]” indicate that the number next to the bracket is included in the solution set.


1. **Solution graph** shows the solution on the real number line. A closed circle on a number indicates that the number is included in the solution set. While an open circle indicates that the number is not included in the set. For our example, the solution graph is drawn here.



Example 6


- a) $[-4, 6]$ says that the solutions is all numbers between -4 and 6 **including** -4 and 6 .

FIGURE 6.1




- b) $(8, 24)$ says that the solution is all numbers between 8 and 24 but **does not include** the numbers 8 and 24.

FIGURE 6.2




- c) $[3, 12)$ says that the solution is all numbers between 3 and 12, **including** 3 but **not including** 12.

FIGURE 6.3




- d) $(-5, \infty)$ says that the solution is all numbers greater that -5 , **not including** -5 .

FIGURE 6.4



e) $(-\infty, \infty)$ says that the solution is all real numbers.

FIGURE 6.5



Solving an Inequality Using Division


We solve the inequality $2x \geq 12$ by dividing both sides by 2. Then we simplify

Original problem.	$2x \geq 12$
Divide both sides by 2.	$\frac{2x}{2} \geq \frac{12}{2}$
Simplify	$x \geq 6$

Let's write the solution in the four different notations you just learned:

Inequality notation	$x \geq 6$
Set notation	$\{x x \geq 6\}$
Interval notation	$[6, \infty)$

Solution graph



Multiplying and Dividing an Inequality by a Negative Number

We solve an inequality in a similar way to solving a regular equation. We can add or subtract numbers on both sides of the inequality. We can also multiply or divide **positive** numbers on both sides of an inequality without changing the solution.

Something different happens if we multiply or divide by **negative** numbers. **In this case, the inequality sign changes direction.**

For example, to solve $-3x < 9$

We divide both sides by $[-3]$. The inequality sign changes from $<$ to $>$ because we divide by a negative number. $\frac{-3x}{-3} > \frac{9}{-3}$

$$x > -3$$

We can explain why this happens with a simple example. We know that two is less than three, so we can write the inequality.

$$2 < 3$$

If we multiply both numbers by -1 we get -2 and -3 , but we know that -2 is greater than -3 .

$$-2 > -3$$

You see that multiplying both sides of the inequality by a negative number caused the inequality sign to change direction. This also occurs if we divide by a negative number.

Example 7

Solve each inequality. Give the solution in inequality notation and interval notation.

a) $4x < 24$

b) $-9x \geq -\frac{3}{5}$

c) $-5x \leq 21$

d) $12 > -30$

Solution:

a)

Original problem.	$4x < 24$
Divide both sides by 4.	$\frac{4x}{4} < \frac{24}{4}$
Simplify	$x < 6$ or $(-\infty, 6)$ Answer

b)

Original problem :	$-9x \geq -\frac{3}{5}$
Divide both sides by -9 .	$\frac{-9}{-9} \leq \frac{-\frac{3}{5}}{-9} \cdot \frac{1}{-\frac{1}{9}}$ Direction of the inequality is changed
Simplify.	$x \geq \frac{1}{15}$ or $\left[\frac{1}{15}, \infty\right)$ Answer

Original problem :	$-5x \leq 21$
Divide both sides by -5 .	$\frac{-5x}{-5} \geq \frac{21}{-5}$ Direction of the inequality is changed
Simplify.	$x \geq -\frac{21}{5}$ or $\left[-\frac{21}{5}, \infty\right)$ Answer

d)

Original problem	$12x > -30$
Divide both sides by 12.	$\frac{12x}{12} > \frac{-30}{12}$
Simplify.	$x > -\frac{5}{2}$ or $\left(-\frac{5}{2}, \infty\right)$ Answer

Example 8

Solve each inequality. Give the solution in inequality notation and solution graph.

a) $\frac{x}{2} > 40$

b) $\frac{x}{-3} \leq -12$

c) $\frac{x}{25} < \frac{3}{2}$

d) $\frac{x}{-7} \geq 9$


Solution

a)

Original problem	$\frac{x}{2} > 40$
Multiply both sides by 2.	$2 \cdot \frac{x}{2} > 40 \cdot 2$ Direction of inequality is NOT changed
Simplify.	$x > -80$ Answer

b)

Original problem	$\frac{x}{-3} \leq -12$
Multiply both sides by -3 .	$-3 \cdot \frac{x}{-3} \geq -12 \cdot (-3)$ Direction of inequality is changed
Simplify.	$x \geq 36$ Answer




c)

Original problem	$\frac{x}{25} < \frac{3}{2}$
Multiply both sides by 25.	$25 \cdot \frac{x}{25} < \frac{3}{2} \cdot 25$ direction of inequality is NOT changed
Simplify.	$x < \frac{75}{2}$ or $x < 37.5$ Answer



d)

Original problem	$\frac{x}{-7} \geq 9$
Multiply both sides by -7 .	$-7 \cdot \frac{x}{-7} \leq 9 \cdot (-7)$ Direction of inequality is changed
Simplify.	$x \leq -63$ Answer



Lesson Summary

- The answer to an **inequality** is often an **interval of values**. Common **inequalities** are:
 - $>$ is greater than
 - \geq is greater than or equal to
 - $<$ is less than
 - \leq is less than or equal to
- Solving inequalities with **addition** and **subtraction** works just like solving an equation. To solve, we isolate the variable on one side of the equation.
- There are four ways to represent an inequality:


1. **Equation notation** $x \geq 2$

2. **Set notation** $x \geq 2$

3. **Interval notation** $[2, \infty)$

Closed brackets “[” and “]” mean inclusive, parentheses “(” and “)” mean exclusive.


4. **Solution graph**




- When multiplying or dividing both sides of an inequality by a negative number, you need to **reverse the inequality**.

Review Questions 1


1. Write the inequality represented by the graph.




2. Write the inequality represented by the graph.



3. Write the inequality represented by the graph.



4. Write the inequality represented by the graph.



Graph each inequality on the number line.

5. $x < -35$
6. $x > -17$
7. $x \geq 20$
8. $x \leq 3$

Solve each inequality and graph the solution on the number line.

9. $x - 5 < 35$
10. $x + 15 \geq -60$
11. $x - 2 \leq 1$
12. $x - 8 > -20$
13. $x + 11 > 13$
14. $x + 65 < 100$
15. $x - 32 \leq 0$
16. $x + 68 \geq 75$

Review Questions 2

Solve each inequality. Give the solution in inequality notation and solution graph.

1. $3x \leq 6$
2. $\frac{x}{5} > -\frac{3}{10}$
3. $-10x > 250$
4. $\frac{x}{-7} \geq -5$

Solve each inequality. Give the solution in inequality notation and interval notation.


5. $9x > -\frac{3}{4}$
6. $-\frac{x}{15} > \leq 5$
7. $620x > 2400$
8. $\frac{x}{20} \geq -\frac{7}{40}$

Solve each inequality. Give the solution in inequality notation and set notation.

9. $-0.5x \leq 7.5$
10. $75x \geq 125$
11. $\frac{x}{-3} > -\frac{10}{9}$
12. $\frac{x}{-15} < 8$

Review Answers 1

1. $x \geq 1$
2. $x < -10$
3. $x \leq -10$
4. $x > 30$




Review Answers 2


1. $x \leq 2$ or



2. $x > -\frac{3}{2}$ or



3. $x < -25$ or



4. $x \leq 35$ or



5. $x > -\frac{1}{12}$ or $(-\frac{1}{12}, \infty)$

6. $x \geq -75$ or $[-75, \infty)$

7. $x < 3.9$ or $(-\infty, 3.9)$

8. $x \geq -\frac{7}{2}$ or $[-\frac{7}{2}, \infty)$

9. $x \geq -15$ or x is a real number $| x \geq -15$

10. $x \geq \frac{5}{3}$ or x is a real number $| x \geq \frac{5}{3}$

11. $x < -\frac{10}{3}$ or x is a real number $| x < -\frac{10}{3}$

12. $x > -120$ or x is a real number $| x > -120$

6.2 Multi-Step Inequalities

Learning Objectives

- Solve a two-step inequality.
- Solve a multi-step inequality.
- Identify the number of solutions of an inequality.
- Solve real-world problems using inequalities.

Introduction

In the last two sections, we considered very simple inequalities which required one-step to obtain the solution. However, most inequalities require several steps to arrive at the solution. As with solving equations, we must use the order of operations to find the correct solution. In addition **remember that when we multiply or divide the inequality by a negative number the direction of the inequality changes.**

The general procedure for solving multi-step inequalities is as follows.

1. Clear parenthesis on both sides of the inequality and collect like terms.
2. Add or subtract terms so the variable is on one side and the constant is on the other side of the inequality sign.
3. Multiply and divide by whatever constants are attached to the variable. Remember to change the direction of the inequality if you multiply or divide by a negative number.

Solve a Two-Step Inequality

Example 1

Solve each of the following inequalities and graph the solution set.

a) $6x - 5 < 10$

b) $-9x < -5x - 15$

c) $-\frac{9x}{5} \leq 24$

Solution


a)

Original problem :	$6x - 5 < 10$
Add 5 to both sides : .	$6x - 5 + 5 < 10 + 5$
Simplify.	$6x < 15$
Divide both sides by 6.	$\frac{6x}{6} < \frac{15}{6}$
Simplify.	$x < \frac{5}{2}$ Answer




b)

Original problem.	$-9x \leq -5x - 15$
Add $5x$ to both sides.	$-9x + 5x \leq -5x + 5x - 15$
Simplify.	$-4x < -15$
Divide both sides by -4 .	$\frac{-4x}{-4} > \frac{-15}{-4}$ Inequality sign was flipped
Simplify.	$x > \frac{15}{4}$ Answer



c)

Original problem.	$-9x \leq 24$
Multiply both sides by 5.	$\frac{-9x}{5} \cdot 5 \leq 24 \cdot 5$
Simplify.	$-9x \leq 120$
Divide both sides by -9 .	$\frac{-9x}{-9} > \frac{120}{-9}$ Inequality sign was flipped
Simplify.	$x \geq -\frac{40}{3}$ Answer



Solve a Multi-Step Inequality

Example 2

Each of the following inequalities and graph the solution set.


a) $\frac{9x}{5} - 7 \geq -3x + 12$

b) $-25x + 12 \leq -10x - 12$

Solution


a)

Original problem	$\frac{9x}{5} - 7 \geq -3x + 12$
Add 3x to both sides.	$\frac{9x}{5} + 3x - 7 \geq -3x + 3x + 12$
Simplify.	$\frac{24x}{5} - 7 \geq 12$
Add 7 to both sides.	$\frac{24x}{5} - 7 + 7 \geq 12 + 7$
Simplify.	$\frac{24x}{5} - 7 \geq 19$
Multiply 5 to both sides.	$5 \cdot \frac{24x}{5} \geq 5 \cdot 19$
Simplify.	$24x \geq 95$
Divide both sides by 24.	$\frac{24x}{24} \geq \frac{95}{24}$
Simplify.	$x \geq \frac{95}{24}$ Answer



b)

Original problem	$-25x + 12 \leq -10x - 12$
Add 5 to both sides.	$-25x + 10x + 12 \leq -10x + 10x - 12$
Simplify.	$-15x + 12 \leq -12$
Subtract 12 from both sides.	$-15x + 12 - 12 \leq -12 - 12$
Simplify.	$-15x \leq -24$
Divide both sides by -15 .	$\frac{-15x}{-15} \geq \frac{-24}{-15}$ Inequality sign was flipped
Simplify.	$x \geq \frac{8}{5}$ Answer



Example 3

Solve the following inequalities.

- a) $4x - 2(3x - 9) \leq -4(2x - 9)$
- b) $\frac{5x-1}{4} > -2(x+5)$

Solution

a)

Original problem	$4x - 2(3x - 9) \leq -4(2x - 9)$
Simplify parentheses.	$4x - 6x + 18 \leq -8x + 36$
Collect like terms.	$-2x + 18 \leq -8x + 36$
Add $8x$ to both sides.	$-2x + 8x + 18 \leq -8x + 8x + 36$
Simplify.	$-6x + 18 \leq 36$
Subtract 18 from both sides.	$-6x + 18 - 18 \leq 36 - 18$
Simplify.	$6x \leq 18$
Divide both sides by 6.	$\frac{6x}{6} \leq \frac{18}{6}$
Simplify.	$x \leq 3$ Answer

b)

Original problem	$\frac{5x - 1}{4} > -2(x + 5)$
Simplify parenthesis.	$\frac{5x - 1}{4} > -2x - 10$
Multiply both sides by 4.	$4 \cdot \frac{5x - 1}{4} > 4(-2x - 10)$
Simplify.	$5x - 1 > -8x - 40$
Add $8x$ to both sides.	$5x + 8x - 1 > -8x + 8x - 40$
Simplify.	$13x - 1 > -40$
Add 1 to both sides.	$13x - 1 + 1 > -40 + 1$
Simplify.	$13x > -39$
Divide both sides by 13.	$\frac{13x}{13} > \frac{-39}{13}$
Simplify.	$x > -3$ Answer

Identify the Number of Solutions of an Inequality

Inequalities can have:

- A set that has an infinite number of solutions.
- No solutions
- A set that has a discrete number of solutions.

Infinite Number of Solutions

The inequalities we have solved so far all have an infinite number of solutions. In the last example, we saw that the inequality

$$\frac{5x-1}{4} > -2(x+5) \text{ has the solution } x > -3$$

This solution says that all real numbers greater than -3 make this inequality true. You can see that the solution to this problem is an infinite set of numbers.

No solutions

Consider the inequality
This simplifies to

$$x - 5 > x + 6$$
$$-5 > 6$$

This statements is not true for any value of x . We say that this inequality has no solution.

Discrete solutions

So far we have assumed that the variables in our inequalities are real numbers. However, in many real life situations we are trying to solve for variables that represent integer quantities, such as number of people, number of cars or number of ties.



Example 4

Raul is buying ties and he wants to spend \$200 or less on his purchase. The ties he likes the best cost \$50. How many ties could he purchase?

Solution

Let x = the number of ties Raul purchases.

We can write an inequality that describes the purchase amount using the formula.


$$(\text{number of ties}) \times (\text{price of a tie}) \leq \$200 \text{ or } 50x \leq 200$$

We simplify our answer. $x \leq 4$

This solution says that Raul bought four or less ties. Since ties are discrete objects, the solution set consists of five numbers $\{0, 1, 2, 3, 4\}$.

Solve Real-World Problems Using Inequalities

Sometimes solving a word problem involves using an inequality.



Example 5

In order to get a bonus this month, Leon must sell at least 120 newspaper subscriptions. He sold 85 subscriptions in the first three weeks of the month. How many subscriptions must Leon sell in the last week of the month?

Solution

Step 1

We know that Leon sold 85 subscriptions and he must sell at least 120 subscriptions.

We want to know the least amount of subscriptions he must sell to get his bonus.

Let x = the number of subscriptions Leon sells in the last week of the month.

Step 2

The number of subscriptions per month must be greater than 120.

We write

$$85 + x \geq 120$$

Step 3

We solve the inequality by subtracting 85 from both sides $x \geq 35$

Answer Leon must sell 35 or more subscriptions in the last week to get his bonus.

Step 4:

To check the answer, we see that $85 + 35 = 120$. If he sells 35 or more subscriptions the number of subscriptions sold that month will be 120 or more.

**Example 6**

Virena's Scout Troup is trying to raise at least \$650 this spring. How many boxes of cookies must they sell at \$4.50 per box in order to reach their goal?

Solution

Step 1

Virena is trying to raise at least \$650

Each box of cookies sells for \$4.50

Let x = number of boxes sold

The inequality describing this problem is:

$$4.50x \geq 650.$$

Step 3

We solve the inequality by dividing both sides by 4.50


$$x \geq 1444.44$$

Answer We round up the answer to 145 since only whole boxes can be sold.

Step 4

If we multiply 145 by \$4.50 we obtain \$652.50. If Virena's Troop sells more than 145 boxes, they raise more than \$650.

The answer checks out.



Example 7

The width of a rectangle is 20 inches . What must the length be if the perimeter is at least 180 inches?

Solution

Step 1

width = 20 inches

Perimeter is at least 180 inches

What is the smallest length that gives that perimeter?

Let x = length of the rectangle

Step 2

Formula for perimeter is $\text{Perimeter} = 2 \times \text{length} + 2 \times \text{width}$

Since the perimeter must be at least 180 inches , we have the following equation.

$$2x + 2(20) \geq 180$$

Step 3

We solve the inequality.

Simplify.

$$2x + 40 \geq 180$$

Subtract 40 from both sides.

$$2x \geq 140$$

Divide both sides by 2.

$$x \geq 70$$

Answer The length must be at least 70 inches .

Step 4

If the length is at least 70 inches and the width is 20 inches, then the perimeter can be found by using this equation.

$$2(70) + 2(20) = 180 \text{ inches}$$

The answer check out.

Lesson Summary

- The **general procedure** for solving multi-step inequalities is as follows.
 3. Multiply and divide by whatever constants are attached to the variable. Remember to change the direction of the inequality if you multiply or divide by a negative number.
 2. Add or subtract terms so the variable is on one side and the constant is on the other side of the inequality sign.
 1. Clear parentheses on both sides of the inequality and collect like terms.
- Inequalities can have **multiple solutions**, **no solutions**, or **discrete solutions**.


Review Questions

Solve the following inequalities and give the solution in set notation and show the solution graph.


1. $4x + 3 < -1$
2. $2x < 7x - 36$
3. $5x > 8x + 27$
4. $5 - x < 9 + x$
5. $4 - 6x \leq 2(2x + 3)$
6. $5(4x + 3) \geq 9(x - 2) - x$
7. $2(2x - 1) + 3 < 5(x + 3) - 2x$
8. $8x - 5(4x + 1) \geq -1 + 2(4x - 3)$
9. $2(7x - 2) - 3(x + 2) < 4x - (3x + 4)$
10. $\frac{2}{3}x - \frac{1}{2}(4x - 1) \geq x + 2(x - 3)$
11. At the San Diego Zoo, you can either pay \$22.75 for the entrance fee or \$71 for the yearly pass which entitles you to unlimited admission. At most how many times can you enter the zoo for the \$22.75 entrance fee before spending more than the cost of a yearly membership?
12. Proteek's scores for four tests were 82, 95, 86 and 88. What will he have to score on his last test to average at least 90 for the term?

Review Answers


1. $\{x \mid x \text{ is a real number, } x < -1\}$



2. $\{x \mid x \text{ is a real number, } x > \frac{36}{5}\}$



3. $\{x \mid x \text{ is a real number, } x < -9\}$



4. $\{x \mid x \text{ is a real number, } x > -2\}$



5. $\{x \mid x \text{ is a real number, } x \geq -\frac{1}{5}\}$




6. $\{x \mid x \text{ is a real number, } x \geq -\frac{33}{12}\}$



7. $\{x \mid x \text{ is a real number, } x < 14\}$



8. $\{x \mid x \text{ is a real number, } x \leq \frac{1}{10}\}$



9. $\{x \mid x \text{ is a real number, } x < \frac{3}{5}\}$



10. $\{x \mid x \text{ is a real number, } x \leq \frac{3}{2}\}$



11. At most 3 times.

12. At least 99.

6.3 Compound Inequalities

Learning Objectives

- Write and graph compound inequalities on a number line.
- Solve a compound inequality with "and".
- Solve a compound inequality with "or".
- Solve compound inequalities using a graphing calculator (TI family).
- Solve real-world problems using compound inequalities

Introduction

In this section, we will solve compound inequalities. In previous sections, we obtained solutions that gave the variable either as greater than or as less than a number. In this section we are looking for solutions where the variable can be in two or more intervals on the number line.

There are two types of compound inequalities:

1. Inequalities joined by the word "and".

The solution is a set of values greater than a number *and* less than another number.

$$a < x < b$$

In this case we want values of the variable for which *both* inequalities are true.

2. Inequalities joined by the word "or".

The solution is a set of values greater than a number or less than another number.

$$x < a \text{ or } x > b$$

In this case, we want values for the variable in which *at least one* of the inequalities is true.

Write and Graph Compound Inequalities on a Number Line


Example 1

Write the inequalities represented by the following number line graphs.


a)




b)



c)



d)

**Solution**

a) The solution graph shows that the solution is any value between -40 and 60 , including -40 but not 60 . Any value in the solution set satisfies both inequalities.

$$x \geq -40 \text{ and } x < 60$$

This is usually written as the following compound inequality.

$$-40 \leq x < 60$$

b) The solution graph shows that the solution is any value greater than 1 (not including 1) or any value less than -2 (not including -2). You can see that there can be no values that can satisfy both these conditions at the same time. We write:

$$x > 1 \text{ or } x < -2$$

c) The solution graph shows that the solution is any value greater than 4 (including 4) or any value less than -1 (including -1). We write:

$$x \geq 4 \text{ or } x \leq -1$$

d) The solution graph shows that the solution is any value less than 25 (not including 25) and any value greater than -25 (not including -25). Any value in the solution set satisfies both conditions.

$$x > -25 \text{ and } x < 25$$

This is usually written as $-25 < x < 25$.

Example 2

Graph the following compound inequalities on the number line.

a) $-4 \leq x \leq 6$

b) $x < 0$ or $x > 2$

c) $x \geq -8$ or $x \leq -20$


d) $-15 < x \leq 85$

Solution


a) The solution is all numbers between -4 and 6 including both -4 and 6 .



b) The solution is either numbers less than 0 or numbers greater than 2 not including 0 or 2 .



c) The solution is either numbers greater than or equal to -8 or less than or equal to -20 .



d) The solution is numbers between -15 and 85 , not including -15 but including 85 .



Solve a compound Inequality With "and"

When we solve compound inequalities, we separate the inequalities and solve each of them separately. Then, we combine the solutions at the end.

Example 3

Solve the following compound inequalities and graph the solution set.

a) $-2 < 4x - 5 \leq 11$


b) $3x - 5 < x + 9 \leq 5x + 13$

Solution

a) First, we rewrite the compound inequality as two separate inequalities with *and*. Then solve each inequality separately.

$$\begin{array}{lll} -2 < 4x - 5 & & 4x - 5 \leq 11 \\ 3 < 4x & \text{and} & 4x \leq 16 \\ \frac{3}{4} < x & & x \leq 4 \end{array}$$

Answer $\frac{3}{4} < x$ and $x \leq 4$. This can be written as $\frac{3}{4} < x \leq 4$.



b) Rewrite the compound inequality as two separate inequalities by using *and*. Then solve each inequality separately.

$$\begin{array}{lll} 3x - 5 < x + 9 & & x + 9 \leq 5x + 13 \\ 2x < 14 & \text{and} & -4 \leq 4x \\ x < 7 & & -1 \leq x \text{ or } x \geq -1 \end{array}$$

Answer $x < 7$ and $x \geq -1$. This can be written as $-1 \leq x < 7$.



Solve a Compound Inequality With "or"

Consider the following example.

Example 4

Solve the following compound inequalities and graph the solution set.

a) $9 - 2x \leq 3$ or $3x + 10 \leq 6 - x$


b) $\frac{x-2}{6} \leq 2x - 4$ or $\frac{x-2}{6} > x + 5$

Solution

a) Solve each inequality separately.

$$\begin{array}{l} 9 - 2x \leq 3 \\ -2x \leq -6 \\ x \geq 3 \end{array} \qquad \text{or} \qquad \begin{array}{l} 3x + 10 \leq 6 - x \\ 4x \leq -4 \\ x \leq -1 \end{array}$$


Answer $x \geq 3$ or $x \leq -1$



b) Solve each inequality separately.

$$\begin{array}{l} \frac{x-2}{6} \leq 2x - 4 \\ x - 2 \leq 6(2x - 4) \\ x - 2 \leq 12x - 24 \\ 22 \leq 11x \\ 2 \leq x \end{array} \qquad \text{or} \qquad \begin{array}{l} \frac{x-2}{6} > x + 5 \\ x - 2 > 6(x + 5) \\ x - 2 > 6x + 30 \\ -32 > 5x \\ -6.4 > x \end{array}$$

Answer $x \geq 2$ or $x < -6.4$



Solve Compound Inequalities Using a Graphing Calculator (TI-83/84 family)

This section explains how to solve simple and compound inequalities with a graphing calculator.


Example 5

Solve the following inequalities using the graphing calculator.

a) $5x + 2(x - 3) \geq 2$

b) $7x - 2 < 10x + 1 < 9x + 5$

c) $3x + 2 \leq 10$ or $3x + 2 \geq 15$


**Solution**

a) $5x + 2(x - 3) \geq 2$

Step 1 Enter the inequality.


Press the [Y=] button.

Enter the inequality on the first line of the screen.



$$Y_1 = 5x + 2(x - 3) \geq 2$$

The \geq symbol is entered by pressing [TEST] [2nd] [MATH] and choose option 4.



Step 2 Read the solution.

Press the **[GRAPH]** button.

Because the calculator translates a true statement with the number 1 and a false statement with the number 0, you will see a step function with the y -value jumping from 0 to 1. The solution set is the values of x for which the graph shows $y = 1$.

X	Y1
1.13	0
1.14	0
1.15	1
1.16	1
1.17	1
1.18	1
1.19	1

X=1.14


Note: You need to press the **[WINDOW]** key or the **[ZOOM]** key to adjust window to see full graph.

The solution is $x \geq \frac{8}{7} = 1.42857\dots$, which is why you can see the y value changing from 0 to 1 at 1.14.

b) $7x - 2 < 10x + 1 < 9x + 5$

This is a compound inequality $7x - 2 < 10x + 1$ and $10x + 1 < 9x + 5$.


To enter a compound inequality:



Press the [Y=] button.

Enter the inequality as $Y_1 = (7x - 2 < 10x + 1) \text{ AND } (10x + 1 < 9x + 5)$

To enter the [AND] symbol press [TEST], choose [LOGIC] on the top row and choose option 1.




The resulting graph looks as shown at the right.

The solution are the values of x for which $y = 1$.

In this case $-1 < x < 4$.


c) $3x + 2 \leq 10$ or $3x + 2 \geq 15$

This is a compound inequality $3x + 2 \leq 10$ or $3x + 2 \geq 15$



Press the [Y=] button.

Enter the inequality as $Y_1 = (3x + 2 \leq 10) \text{ OR } (3x + 2 \geq 15)$ To enter the [OR] symbol press [TEST], choose [LOGIC] on the top row and choose option 2.



The resulting graph looks as shown at the right. The solution are the values of x for which $y = 1$. In this case, $x \leq 2.7$ or $x \geq 4.3$.

Solve Real-World Problems Using Compound Inequalities

Many application problems require the use of compound inequalities to find the solution.

Example 6

The speed of a golf ball in the air is given by the formula $v = -32t + 80$, where t is the time since the ball was hit. When is the ball traveling between 20 ft/sec and 30 ft/sec?

**Solution***Step 1*

We want to find the times when the ball is traveling between 20 ft/sec and 30 ft/sec.

Step 2

Set up the inequality $20 \leq v \leq 30$

Step 3

Replace the velocity with the formula $v = -32t + 80$.

$$20 \leq -32t + 80 \leq 30$$

Separate the compound inequality and solve each separate inequality.

$$\begin{array}{rcl}
 20 \leq -32t + 80 & & -32t + 80 \leq 30 \\
 32t \leq 60 & \text{and} & 50 \leq 32t \\
 t \leq 1.875 & & 1.56 \leq t
 \end{array}$$

Answer $1.56 \leq t \leq 1.875$

Step 4 To check plug in the minimum and maximum values of t into the formula for the speed.


For $t = 1.56$, $v = -32t + 80 = -32(1.56) + 80 = 30$ ft/sec

For $t = 1.875$, $v = -32t + 80 = -32(1.875) + 80 = 20$ ft/sec

So the speed is between 20 and 30 ft/sec . The answer checks out.

Example 7

William's pick-up truck gets between 18 to 22 miles per gallon of gasoline. His gas tank can hold 15 gallons of gasoline. If he drives at an average speed of 40 miles per hour how much driving time does he get on a full tank of gas?



Solution

Step 1 We know

The truck gets between 18 and 22 miles/gallon

There are 15 gallons in the truck's gas tank

William drives at an average of 40 miles/hour

Let t = driving time

Step 2 We use dimensional analysis to get from time per tank to miles per gallon.

$$\frac{t \text{ hours}}{1 \text{ tank}} \times \frac{1 \text{ tank}}{15 \text{ gallons}} \times \frac{40 \text{ miles}}{1 \text{ hours}} = \frac{40t}{15} \frac{\text{miles}}{\text{gallon}}$$

Step 3 Since the truck gets between 18 to 22 miles/gallon, we set up the compound inequality.

$$18 \leq \frac{40t}{15} \leq 22$$

Separate the compound inequality and solve each inequality separately.

$$\begin{array}{rcl} 18 \leq \frac{40t}{15} & & \frac{40t}{15} \leq 22 \\ 270 \leq 40t & \text{and} & 40t \leq 330 \\ 6.75 \leq t & & t \leq 8.25 \end{array}$$

Answer $6.75 \leq t \leq 8.25$. Andrew can drive between 6.75 and 8.25 hours on a full tank of gas.

Step 4

For $t = 6.75$, we get $\frac{40t}{15} = \frac{40(6.75)}{15} = 18$ miles per gallon.


For $t = 8.25$, we get $\frac{40t}{15} = \frac{40(8.25)}{15} = 22$ miles per gallon.

Lesson Summary

- **Compound inequalities** combine two or more inequalities with "**and**" or "**or**".
- "**And**" combinations mean the only solutions for both inequalities will be solutions to the compound inequality.
- "**Or**" combinations mean solutions to either inequality will be solutions to the compound inequality.

Review Questions

Write the compound inequalities represented by the following graphs.





Solve the following compound inequalities and graph the solution on a number line.


5. $-5 \leq x - 4 \leq 13$
6. $1 \leq 3x + 4 \leq 4$
7. $-12 \leq 2 - 5x \leq 7$
8. $\frac{3}{4} \leq 2x + 9 \leq \frac{3}{2}$
9. $-2\frac{2x-1}{3} < -1$
10. $4x - 1 \geq 7$ or $\frac{9x}{2} < 3$
11. $3 - x < -4$ or $3 - x > 10$
12. $\frac{2x+3}{4} < 2$ or $-\frac{x}{5} + 3\frac{2}{5}$
13. $2x - 7 \leq -3$ or $2x - 3 > 11$
14. $4x + 3 \leq 9$ or $-5x + 4 \leq -12$
15. To get a grade of B in her Algebra class, Stacey must have an average grade greater than or equal to 80 and less than 90. She received the grades of 92, 78, 85 on her first three tests. Between which scores must her grade fall if she is to receive a grade of B for the class?

Review Answers

1. $-40 \leq x \leq 70$
2. $x < -2$ or $x \geq 5$
3. $-8 < x < 0$
4. $x \leq -2$ or $x > 1.5$
5. $-1 \leq x \leq 17$




6. $-\frac{4}{3} \leq x \leq -\frac{1}{3}$




7. $-1 \leq x \leq \frac{14}{5}$




8. $-\frac{33}{8} \leq x \leq -\frac{15}{4}$




9. $-\frac{5}{2} \leq x < -1$




10. $x \geq 2$ or $x < \frac{2}{3}$



11. $x > 7$ or $x < -7$



12. $x < \frac{5}{2}$ or $x > 13$



13. $x \leq 2$ or $x > 7$



14. $x < \frac{3}{2}$ or $x \geq \frac{16}{5}$



15. $65 \leq x < 105$


6.4 Absolute Value Equations

Learning Objectives

- Solve an absolute value equation.
- Analyze solutions to absolute value equations.
- Graph absolute value functions.
- Solve real-world problems using absolute value equations.

Introduction

The **absolute value** of a number is its distance from zero on a number line. There are always two numbers on the number line that are the same distance from zero. For instance, the numbers 4 and -4 are both a distance of 4 units away from zero.



$|4|$ represents the distance from 4 to zero which equals 4.

$|-4|$ represents the distance from -4 to zero which also equals 4.

In fact, for any real number x ,

$|x| = x$ if x is not negative (that is, including $x = 0$.)

$|x| = -x$ if x is negative.

Absolute value has no effect on a positive number but changes a negative number into its positive inverse.

Example 1

Evaluate the following absolute values.

a) $|25|$

b) $|-120|$

c) $|-3|$

d) $|55|$

e) $|\frac{-5}{4}|$

Solution:

a) $|25| = 25$ Since 25 is a positive number the absolute value does not change it.

b) $|-120| = 120$ Since -120 is a negative number the absolute value makes it positive.


c) $|-3| = 3$ Since -3 is a negative number the absolute value makes it positive.

d) $|55| = 55$ Since 55 is a positive number the absolute value does not change it.

e) $|\frac{-5}{4}| = \frac{5}{4}$ Since $\frac{-5}{4}$ is a negative number the absolute value makes it positive.

Absolute value is very useful in finding the distance between two points on the number line. The distance between any two points a and b on the number line is $|a - b|$ or $|b - a|$.

For example, the distance from 3 to -1 on the number line is $|3 - (-1)| = |4| = 4$.



We could have also found the distance by subtracting in the reverse order, $|-1 - 3| = |-4| = 4$.

This makes sense because the distance is the same whether you are going from 3 to -1 or from -1 to 3.

Example 2

Find the distance between the following points on the number line.

- a) 6 and 15
- b) -5 and 8
- c) -3 and -12

Solutions

Distance is the absolute value of the difference between the two points.

- a) Distance = $|6 - 15| = |-9| = 9$
- b) Distance = $|-5 - 8| = |-13| = 13$
- c) Distance = $|-3 - (-12)| = |9| = 9$


Remember: When we computed the change in x and the change in y as part of the slope computation, these values were positive or negative, depending on the direction of movement. In this discussion, “distance” means a positive distance only.

Solve an Absolute Value Equation

We now want to solve equations involving absolute values. Consider the following equation.

$$|x| = 8$$

This means that the distance from the number x to zero is 8. There are two possible numbers that satisfy this condition 8 and -8 .



When we solve absolute value equations we always consider 2 possibilities.

1. The expression inside the absolute value sign is not negative.
2. The expression inside the absolute value sign is negative.

Then we solve each equation separately.

Example 3

Solve the following absolute value equations.

a) $|3| = 3$

b) $|10| = 10$

Solutiona) There are two possibilities $x = 3$ and $x = -3$.b) There are two possibilities $x = 10$ and $x = -10$.**Analyze Solutions to Absolute Value Equations****Example 4***Solve the equation and interpret the answers.***Solution**

We consider two possibilities. The expression inside the absolute value sign is not negative or is negative. Then we solve each equation separately.

$$x - 4 = 5$$

$$x = 9$$


and

$$x - 4 = -5$$

$$x = -1$$

Answer $x = 9$ and $x = -1$.

Equation $|x - 4| = 5$ can be interpreted as “what numbers on the number line are 5 units away from the number 4?” If we draw the number line we see that there are two possibilities 9 and -1 .

**Example 5***Solve the equation $|x + 3| = 2$ and interpret the answers.***Solution**

Solve the two equations.

$$x + 3 = 2$$

$$x = -1$$


and

$$x + 3 = -2$$

$$x = -5$$

Answer $x = -5$ and $x = -1$.

Equation $|x + 3| = 2$ can be re-written as $|x - (-3)| = 2$. We can interpret this as “what numbers on the number line are 2 units away from -3 ?” There are two possibilities -5 and -1 .



Example 6

Solve the equation $|2x - 7| = 6$ and interpret the answers.


Solution

Solve the two equations.

$$\begin{array}{l} 2x - 7 = -6 \\ 2x = 13 \\ x = \frac{13}{2} \end{array} \qquad \text{and} \qquad \begin{array}{l} 2x - 7 = 6 \\ 2x = 13 \\ x = \frac{13}{2} \end{array}$$

Answer $x = \frac{13}{2}$ and $x = \frac{1}{2}$

The interpretation of this problem is clearer if the equation $|2x - 7| = 6$ was divided by 2 on both sides. We obtain $|x - \frac{7}{2}| = 3$. The question is “What numbers on the number line are 3 units away from $\frac{7}{2}$?” There are two possibilities $\frac{13}{2}$ and $\frac{1}{2}$.


**Graph Absolute Value Functions**

You will now learn how to graph absolute value functions. Consider the function:

$$y = |x - 1|$$

Let's graph this function by making a table of values.

x	$y = x - 1 $
-2	$y = -2 - 1 = -3 = 3$
-1	$y = -1 - 1 = -2 = 2$
0	$y = 0 - 1 = -1 = 1$
1	$y = 1 - 1 = 0 = 0$
2	$y = 2 - 1 = 1 = 1$
3	$y = 3 - 1 = 2 = 2$
4	$y = 4 - 1 = 3 = 3$



You can see that the graph of an absolute value function makes a big “V”. It consists of two line rays (or line segments), one with positive slope and one with negative slope joined at the **vertex** or **cusp**.

We saw in previous sections that to solve an absolute value equation we need to consider two options.

1. The expression inside the absolute value is not negative.
2. The expression inside the absolute value is negative.

The graph of $y = |x - 1|$ is a combination of two graphs.

Option 1

$$y = x - 1$$

when $x - 1 \geq 0$

Option 2

$$y = -(x - 1) \text{ or } y = -x + 1$$

when $x - 1 < 0$

These are both graphs of straight lines.

The two straight lines meet at the vertex. We find the vertex by setting the expression inside the absolute value equal to zero.

$$x - 1 = 0 \text{ or } x = 1$$

We can always graph an absolute value function using a table of values. However, we usually use a simpler procedure.

Step 1 Find the vertex of the graph by setting the expression inside the absolute value equal to zero and solve for x .

Step 2 Make a table of values that includes the vertex, a value smaller than the vertex and a value larger than the vertex. Calculate the values of y using the equation of the function.

Step 3 Plot the points and connect with two straight lines that meet at the vertex.

Example 7

Graph the absolute value function: $y = |x + 5|$.


Solution

Step 1 Find the vertex $x + 5 = 0$ or $x = -5$ vertex.

Step 2 Make a table of values.

x	$y = x + 5 $
-8	$y = -8 + 5 = -3 = 3$
-5	$y = -5 + 5 = 0 = 0$
-2	$y = -2 + 5 = 3 = 3$

Step 3 Plot the points and draw two straight lines that meet at the vertex.



Example 8

Graph the absolute value function $y = |3x - 12|$.


Solution

Step 1 Find the vertex $3x - 12 = 0$ so $x = 4$ is the vertex.


Step 2 Make a table of values:

x	$y = 3x - 12 $
0	$y = 3(0) - 12 = -12 = 12$
4	$y = 3(4) - 12 = 0 = 0$
8	$y = 3(8) - 12 = 12 = 12$

Step 3 Plot the points and draw two straight lines that meet at the vertex.



Solve Real-World Problems Using Absolute Value Equations



Example 9

A company packs coffee beans in airtight bags. Each bag should weigh 16 ounces but it is hard to fill each bag to the exact weight. After being filled, each bag is weighed and if it is more than 0.25 ounces overweight or underweight it is emptied and repacked. What are the lightest and heaviest acceptable bags?

Solution

Step 1

We know that each bag should weigh 16 ounces.

A bag can weigh 0.25 ounces more or less than 16 ounces.

We need to find the lightest and heaviest bags that are acceptable.

Let x = weight of the coffee bag in ounces.

Step 2

The equation that describes this problem is written as $|x - 16| \leq 0.25$.

Step 3

Consider the positive and negative options and solve each equation separately.

$$x - 16 = 0.25$$

$$x = 16.25$$

and

$$x - 16 = -0.25$$

$$x = 15.75$$

Answer The lightest acceptable bag weighs 15.75 ounces and the heaviest weighs 16.25 ounces.

Step 4

We see that $16.25 - 16 = 0.25$ ounces and $16 - 15.75 = 0.25$ ounces. The answers are 0.25 ounces bigger and smaller than 16 ounces respectively.

The answer checks out.

Lesson Summary

- The absolute value of a number is its distance from zero on a number line.

$|x| = x$ if x is not negative.

$|x| = -x$ if x is negative.

- An equation with an absolute value in it **splits into two equations**.

- The expression within the absolute value is **positive**, then the absolute value signs do nothing and can be omitted.
- The expression within the absolute value is **negative**, then the expression within the absolute value signs must be negated before removing the signs.

Review Questions

Evaluate the absolute values.

- $|250|$
- $|-12|$
- $|\frac{2}{5}|$
- $|\frac{1}{10}|$

Find the distance between the points.

- 12 and -11
- 5 and 22
- -9 and -18
- -2 and 3

Solve the absolute value equations and interpret the results by graphing the solutions on the number line.

- $|x - 5| = 10$
- $|x + 2| = 6$
- $|5x - 2| = 3$
- $|4x - 1| = 19$


Graph the absolute value functions.

- $y = |x + 3|$
- $y = |x - 6|$
- $y = |4x + 2|$
- $y = |\frac{x}{3} - 4|$


- A company manufactures rulers. Their 12 – inch rulers pass quality control if they within $\frac{1}{32}$ inches of the ideal length. What is the longest and shortest ruler that can leave the factory?

Review Answers


1. 250
2. 12
3. $\frac{2}{5}$
4. $\frac{1}{10}$
5. 23
6. 17
7. 9
8. 5
9. 15 and -5





10. 4 and -8




11. 1 and $-\frac{1}{5}$





12. 5 and $-\frac{9}{2}$



13.



14.



17. $11\frac{31}{32}$ and $12\frac{1}{32}$

6.5 Absolute Value Inequalities

Learning Objectives

- Solve absolute value inequalities.
- Rewrite and solve absolute value inequalities as compound inequalities.
- Solve real-world problems using absolute value inequalities.

Introduction

Absolute value inequalities are solved in a similar way to absolute value equations. In both cases, you must consider the two options.

1. The expression inside the absolute value is not negative.
2. The expression inside the absolute value is negative.


Then we solve each inequality separately.

Solve Absolute Value Inequalities

Consider the inequality

$$|x| \leq 3$$

Since the absolute value of x represents the distance from zero, the solutions to this inequality are those numbers whose distance from zero is less than or equal to 3. The following graph shows this solution:




Notice that this is also the graph for the compound inequality $-3 \leq x \leq 3$.

Now consider the inequality

$$|x| > 2$$

Since the absolute value of x represents the distance from zero, the solutions to this inequality are those numbers whose distance from zero are more than 2. The following graph shows this solution.



Notice that this is also the graph for the compound inequality $x < -2$ or $x > 2$.

[Video Example](#)

Example 1

Solve the following inequalities and show the solution graph.

a) $|x| < 6$

b) $|x| \geq 2.5$


Solution

a) $|x| < 5$ represents all numbers whose distance from zero is less than 5.



Answer $-5 < x < 5$

b) $|x| \geq 2.5$ represents all numbers whose distance from zero is more than or equal to 2.5.



Answer $x \leq -2.5$ or $x \geq 2.5$

Rewrite and Solve Absolute Value Inequalities as Compound Inequalities

In the last section you saw that absolute value inequalities are compound inequalities.

Inequalities of the type $|x| < a$ can be rewritten as $-a < x < a$

Inequalities of the type $|x| < b$ can be rewritten as $x < -b$ or $x > b$

To solve an absolute value inequality, we separate the expression into two inequalities and solve each of them individually.

Example 2

Solve the inequality $|x - 3| < 7$ and show the solution graph.

Solution

Rewrite as a compound inequality.

Write as two separate inequalities.

$$x - 3 < 7 \text{ and } x - 3 < 7$$

Solve each inequality

$$x < 10 \text{ and } x > -4$$

The solution graph is

**Example 3**

Solve the inequality $|4x + 6| \leq 13$ and show the solution graph.

Solution

Rewrite as a compound inequality.

Write as two separate inequalities


$$4x + 5 \leq 13 \text{ and } 4x + 5 \geq -13$$

Solve each inequality:

$$4x \leq 8 \text{ and } 4x \geq -18$$

$$x \leq 2 \text{ and } x \geq -\frac{9}{2}$$

The solution graph is



Example 4

Solve the inequality $|x + 12| > 2$ and show the solution graph.

Solution

Rewrite as a compound inequality.

Write as two separate inequalities.

$$x + 12 < -2 \text{ or } x + 12 > 2$$

Solve each inequality

$$x < -14 \text{ or } x > -10$$

The solution graph is



Example 5

Solve the inequality $|8x - 15| \geq 9$ and show the solution graph.

Rewrite as a compound inequality.

Write as two separate inequalities.


$$8x - 15 \leq -9 \text{ or } 8x - 15 \geq 9$$

Solve each inequality

$$8x \leq 6 \text{ or } 8x \geq 24$$

$$x \leq \frac{3}{4} \text{ or } x \geq 3$$

The solution graph is



Solve Real-World Problems Using Absolute Value Inequalities

Absolute value inequalities are useful in problems where we are dealing with a range of values.

Example 6:

The velocity of an object is given by the formula $v = 25t - 80$ where the time is expressed in seconds and the velocity is expressed in feet per seconds. Find the times when the magnitude of the velocity is greater than or equal to 60 feet per second.

Solution

Step 1

We want to find the times when the velocity is greater than or equal to 60 feet per second

Step 2

We are given the formula for the velocity $v = 25t - 80$

Write the absolute value inequality $|25t - 80| \geq 60$

Step 3

Solve the inequality

$$25t - 80 \geq 60 \text{ or } 25t - 80 \leq -60$$

$$25t \geq 140 \text{ or } 25t \leq 20$$

$$t \geq 5.6 \text{ or } t \leq 0.8$$

Answer: The magnitude of the velocity is greater than 60 ft/sec for times less than 0.8 seconds and for times greater than 5.6 seconds.

Step 4 When $t = 0.8$ seconds, $v = 25(0.8) - 80 = -60$ ft/sec. The magnitude of the velocity is 60 ft/sec. The negative sign in the answer means that the object is moving backwards.

When $t = 5.6$ seconds, $v = 25(5.6) - 80 = 60$ ft/sec.

To find where the magnitude of the velocity is greater than 60 ft/sec, check values in each of the following time intervals: $t \leq 0.8$, $0.8 \leq t \leq 5.6$ and $t \geq 5.6$.

Check $t = 0.5$: $v = 25(0.5) - 80 = -67.5$ ft/sec

Check $t = 2$: $v = 25(2) - 80 = -30$ ft/sec

Check $t = 6$: $v = 25(6) - 80 = 70$ ft/sec

You can see that the magnitude of the velocity is greater than 60 ft/sec for $t \geq 5.6$ or $t \leq 0.8$.

The answer checks out.

Lesson Summary

- Like absolute value equations, inequalities with absolute value split into two inequalities. One where the expression within the absolute value is negative and one where it is positive.
- Inequalities of the type $|x| < a$ can be rewritten as $-a < x < a$.
- Inequalities of the type $|x| > b$ can be rewritten as $-x < -b$ or $x > b$.

Review Questions


Solve the following inequalities and show the solution graph.

1. $|x| \leq 6$
2. $|x| > 3.5$
3. $|x| < 12$
4. $|\frac{x}{5}| \leq 6$
5. $|7x| \geq 21$
6. $|x - 5| > 8$
7. $|x + 7| < 3$
8. $|x - \frac{3}{4}| \leq \frac{1}{2}$
9. $|2x - 5| \geq 13$


10. $|5x + 3| < 7$
11. $|\frac{x}{3} - 4| \leq 2$
12. $|\frac{2x}{7} + 9| > \frac{5}{7}$
13. A three month old baby boy weighs an average of 13 pounds . He is considered healthy if he is 2.5 lbs more or less than the average weight. Find the weight range that is considered healthy for three month old boys.

Review Answers

1. $-6 \leq x \leq 6$




2. $x < -3.5$ or $x > 3.5$




3. $-12 < x < 12$




4. $x < -10$ or $x > 10$




5. $x \leq -3$ or $x \geq 3$




6. $x < -3$ or $x > 13$




7. $-10 < x < -4$



8. $\frac{1}{4} \leq x \leq \frac{5}{4}$




9. $x \leq -4$ or $x \geq 9$



10. $-2 < x < \frac{4}{5}$



11. $6 \leq x \leq 18$



12. $x < -34$ or $x > -29$




13. A healthy weight is $10.5 \text{ lb} \leq x \leq 15.5 \text{ lb}$.

6.6 Linear Inequalities in Two Variables

Learning Objectives

- Graph linear inequalities in one variable on the coordinate plane.
- Graph linear inequalities in two variables.
- Solve real-world problems using linear inequalities

Introduction



A linear inequality in two variables takes the form

$$y > mx + b \text{ or } y < mx + b$$

Linear inequalities are closely related to graphs of straight lines. A straight line has the equation $y = mx + b$. When we graph a line in the coordinate plane, we can see that it divides the plane in two halves.

The solution to a linear inequality includes all the points in one of the plane halves. We can tell which half of the plane the solution is by looking at the inequality sign.

$>$ The solution is the half plane above the line.

\geq The solution is the half plane above the line and also all the points on the line.

$<$ The solution is the half plane below the line.

\leq The solution is the half plane below the line and also all the points on the line.


(Above the line means for a given x -coordinate, all points with y -values greater than the y -value are on the line)

For a strict inequality, we draw a **dashed line** to show that the points on the line are not part of the solution.


For an inequality that includes the equal sign, we draw a **solid line** to show that the points on the line are part of the solution.

Here is what you should expect linear inequality graphs to look like.


The solution of $y > mx + b$ is the half plane above the line. The dashed line shows that the points on the line are not part of the solution.




The solution of $y \geq mx + b$ is the half plane above the line and all the points on the line.



The solution of $y < mx + b$ is the half plane below the line.



The solution of $y \leq mx + b$ is the half plane below the line and all the points on the line.



Graph Linear Inequalities in One Variable in the Coordinate Plane


In the last few sections, we graphed inequalities in one variable on the number line. We can also graph inequalities in one variable on the coordinate plane. We just need to remember that when we graph an equation of the type $x = a$ we get a vertical line and when we graph an equation of the type $y = b$ we get a horizontal line.

Example 1


Graph the inequality $x > 4$ on the coordinate plane.

Solution

First, let's remember what the solution to $x > 4$ looks like on the number line.



The solution to this inequality is the set of all real numbers x that are bigger than four but not including four. The solution is represented by a line.



In two dimensions we are also concerned with values of y , and the solution to $x > 4$ consists of all coordinate points for which the value of x is bigger than four. The solution is represented by the half plane to the right of $x = 4$.

The line $x = 4$ is dashed because the equal sign is not included in the inequality and therefore points on the line are not included in the solution.

Example 2

Graph the inequality $y \leq 6$ on the coordinate plane.

Solution

The solution is all coordinate points for which the value of y is less than or equal than 6. This solution is represented by the half plane below the line $y = 6$.


The line $y = 6$ is solid because the equal sign is included in the inequality sign and the points on the line are included in the solution.

Example 3

Graph the inequality $|6| < 5$

Solution

The absolute value inequality $|6| < 5$ can be re-written as $-5 < y < 5$. This is a compound inequality which means $y > -5$ and $y < 5$



In other words, the solution is all the coordinate points for which the value of y is larger than -5 and smaller than 5. The solution is represented by the plane between the horizontal lines $y = -5$ and $y = 5$.

Both horizontal lines are dashed because points on the line are not included in the solution.


Example 4

Graph the inequality $|x| \geq 2$.

Solution

The absolute value inequality $|x| \geq 2$ can be re-written as a compound inequality:

$$x \leq -2 \text{ or } x \geq 2$$



In other words, the solution is all the coordinate points for which the value of x is smaller than or equal to -2 and greater than or equal to 2 . The solution is represented by the plane to the left of the vertical line $x = -2$ and the plane to the right of line $x = 2$.

Both vertical lines are solid because points on the line are included in the solution.

Graph Linear Inequalities in Two Variables

The general procedure for graphing inequalities in two variables is as follows.

Step 1: Re-write the inequality in slope-intercept form $y = mx + b$. Writing the inequality in this form lets you know the direction of the inequality

Step 2 Graph the line of equation $y = mx + b$ using your favorite method. (For example, plotting two points, using slope and y -intercept, using y -intercept and another point, etc.). Draw a dashed line if the equal sign is not included and a solid line if the equal sign is included.

Step 3 Shade the half plane above the line if the inequality is greater than. Shade the half plane under the line if the inequality is less than.

Example 5

Graph the inequality $y \geq 2x - 3$.

Solution

Step 1

The inequality is already written in slope-intercept form $y \geq 2x - 3$.


x	y
-1	$2(-1) - 3 = -5$
0	$2(0) - 3 = -3$
1	$2(1) - 3 = -1$

Step 2

Graph the equation $y = 2x - 3$ by making a table of values.

Step 3

Graph the inequality. We shade the plane above the line because y is greater than. The value $2x - 3$ defines the line. The line is solid because the equal sign is included.

**Example 6**

Graph the inequality $5x - 2y > 4$.

Solution*Step 1*

Rewrite the inequality in slope-intercept form.

$$\begin{aligned} -2y &> -5x + 4 \\ y &> \frac{5}{2}x - 2 \end{aligned}$$

Notice that the inequality sign changed direction due to division of negative sign.


Step 2

Graph the equation $y > \frac{5}{2}x - 2$ by making a table of values.

x	y
-2	$\frac{5}{2}(-2) - 2 = -7$
0	$\frac{5}{2}(0) - 2 = -2$
2	$\frac{5}{2}(2) - 2 = 3$

Step 3

Graph the inequality. We shade the plane **below** the line because the inequality in slope-intercept form is less than. The line is dashed because the equal sign is not included.

**Example 7**

Graph the inequality $y + 4 \leq -\frac{x}{3} + 5$.

Solution

Step 1

Rewrite the inequality in slope-intercept form $y \leq -\frac{x}{3} + 1$


Step 2

Graph the equation $y = -\frac{x}{3} + 1$ by making a table of values.

x	y
-3	$-\frac{(-3)}{3} + 1 = 2$
0	$-\frac{0}{3}(0) + 1 = 1$
3	$-\frac{3}{3} + 1 = 0$

Step 3

Graph the inequality. We shade the plane below the line. The line is solid because the equal sign is included.



Solve Real-World Problems Using Linear Inequalities

In this section, we see how linear inequalities can be used to solve real-world applications.

Example 8

A pound of coffee blend is made by mixing two types of coffee beans. One type costs \$9 per pound and another type costs \$7 per pound. Find all the possible mixtures of weights of the two different coffee beans for which the blend costs \$8.50 per pound or less.

Solution

Let's apply our problem solving plan to solve this problem.

Step 1:

Let x = weight of \$9 per pound coffee beans in pounds

Let y = weight of \$7 per pound coffee beans in pounds

Step 2

The cost of a pound of coffee blend is given by $9x + 7y$.

We are looking for the mixtures that cost \$8.50 or less.

We write the inequality $9x + 7y \leq 8.50$.

Step 3

To find the solution set, graph the inequality $9x + 7y \leq 8.50$.


Rewrite in slope-intercept $y \leq -1.29x + 1.21$.

Graph $y = -1.29x + 1.21$ by making a table of values.

x	y
0	1.21
1	-0.08
2	-1.37

Step 4

Graph the inequality. The line will be solid. We shade below the line.



Notice that we show only the first quadrant of the coordinate plane because the weight values should be positive.

The blue-shaded region tells you all the possibilities of the two bean mixtures that will give a total less than or equal to \$8.50.

Example 9

Julian has a job as an appliance salesman. He earns a commission of \$60 for each washing machine he sells and \$130 for each refrigerator he sells. How many washing machines and refrigerators must Julian sell in order to make \$1000 or more in commission?

Solution Let's apply our problem solving plan to solve this problem.

Step 1

Let x = number of washing machines Julian sells

Let y = number of refrigerators Julian sells

Step 2

The total commission is given by the expression $60x + 130y$.

We are looking for total commission of \$1000 or more. We write the inequality. $60x + 130y \geq 1000$.

Step 3

To find the solution set, graph the inequality $60x + 130y \geq 1000$.


Rewrite it in slope-intercept $y \geq -.46x + 7.7$.

Graph $y = -.46x + 7.7$ by making a table of values.

x	y
0	7.7
2	6.78
4	5.86

Step 4

Graph the inequality. The line will be solid. We shade above the line.



Notice that we show only the first quadrant of the coordinate plane because dollar amounts should be positive. Also, only the points with integer coordinates are possible solutions.

Lesson Summary

- The general procedure for graphing inequalities in two variables is as follows:

Step 1

Rewrite the inequality in slope-intercept form $y = mx + b$.

Step 2

Graph the line of equation $y = mx + b$ by building a table of values.

Draw a dashed line if the equal sign is not included and a solid line if it is included.

Step 3

Shade the half plane above the line if the inequality is greater than.


Shade the half plane under the line if the inequality is less than.






Review Questions






Graph the following inequalities on the coordinate plane.




- $x < 20$
- $y \geq -5$
- $|x| > 10$
- $|y| \leq 7$
- $y \leq 4x + 3$
- $y > -\frac{x}{2} - 6$
- $3x - 4y \geq 12$
- $x + 7y < 5$
- $6x + 5y > 1$
- $y + 5 \leq -4x + 10$
- $x - \frac{1}{2}y \geq 5$
- $30x + 5y < 100$
- An ounce of gold costs \$670 and an ounce of silver costs \$13. Find all possible weights of silver and gold that makes an alloy that costs less than \$600 per ounce.
- A phone company charges 50 cents per minute during the daytime and 10 cents per minute at night. How many daytime minutes and night time minutes would you have to use to pay more than \$20 over a 24 hour period?

Review Answers









CHAPTER

7**Systems of Equations and Inequalities****Chapter Outline**

- 7.1** **LINEAR SYSTEMS BY GRAPHING**
 - 7.2** **SOLVING LINEAR SYSTEMS BY SUBSTITUTION**
 - 7.3** **SOLVING LINEAR SYSTEMS BY ELIMINATION THROUGH ADDITION OR SUBTRACTION**
 - 7.4** **SOLVING SYSTEMS OF EQUATIONS BY MULTIPLICATION**
 - 7.5** **SPECIAL TYPES OF LINEAR SYSTEMS**
 - 7.6** **SYSTEMS OF LINEAR INEQUALITIES**
-

7.1 Linear Systems by Graphing

Learning Objectives

- Determine whether an ordered pair is a solution to a system of equations.
- Solve a system of equations graphically.
- Solve a system of equations graphically with a graphing calculator.
- Solve word problems using systems of equations.

Introduction

In this lesson, we will discover methods to determine if an ordered pair is a solution to a system of two equations. We will then learn to solve the two equations graphically and confirm that the solution is the point where the two lines intersect. Finally, we will look at real-world problems that can be solved using the methods described in this chapter.

Determine Whether an Ordered Pair is a Solution to a System of Equations


A linear system of equations consists of a set of equations that must be solved together. Consider the following system of equations.

$$\begin{aligned}y &= x + 2 \\y &= -2x + 1\end{aligned}$$

Since the two lines are in a system we deal with them together by graphing them on the same coordinate axes. The lines can be graphed using your favorite method. Lets graph by making a table of values for each line.

Line 1 $y = x + 2$

x	y
0	2
1	3



Line 2 $y = -2x + 1$

x	y
0	1
1	-1

A solution for a single equation is any point that lies on the line for that equation. A solution for a system of equations is any point that lies on both lines in the system.

For Example

- Point A is not a solution to the system because it does not lie on either of the lines.
- Point B is not a solution to the system because it lies only on the blue line but not on the red line.
- Point C is a solution to the system because it lies on both lines at the same time.

In particular, this point marks the intersection of the two lines. It solves both equations, so it solves the system. For a system of equations, the geometrical solution is the intersection of the two lines in the system. The algebraic solution is the ordered pair that solves both equations.

You can confirm the solution by plugging it into the system of equations, and confirming that the solution works in each equation.

Example 1

Determine which of the points $(1,3)$, $(0,2)$ or $(2,7)$ is a solution to the following system of equations.

$$y = 4x - 1$$

$$y = 2x + 3$$

Solution

To check if a coordinate point is a solution to the system of equations, we plug each of the x and y values into the equations to see if they work.

Point $(1,3)$

$$\begin{aligned}y &= 4x - 1 \\3^? &= ? 4(1) - 1 \\3 &= 3\end{aligned}$$

The solution checks.

$$\begin{aligned}y &= 2x + 3 \\3^? &= ? 2(1) + 3 \\3 &\neq 5\end{aligned}$$

The solution does not check.

Point $(1, 3)$ is on line $y = 4x - 1$ but it is not on line $y = 2x + 3$ so it is not a solution to the system.

Point $(0, 2)$

$$\begin{aligned}y &= 4x - 1 \\2^? &= ? 4(0) - 1 \\2 &\neq -1\end{aligned}$$

The solution does not check.

Point $(0, 2)$ is not on line $y = 4x - 1$ so it is not a solution to the system. Note that it is not necessary to check the second equation because the point needs to be on both lines for it to be a solution to the system.

Point $(2, 7)$

$$\begin{aligned}y &= 4x - 1 \\7^? &= ? 4(2) - 1 \\7 &= 7\end{aligned}$$

The solution checks.

$$\begin{aligned}y &= 2x + 3 \\7^? &= ? 2(2) + 3 \\7 &= 7\end{aligned}$$

The solution checks.

Point $(2, 7)$ is a solution to the system since it lies on both lines.

Answer The solution to the system is point $(2, 7)$.

Determine the Solution to a Linear System by Graphing

The solution to a linear system of equations is the point which lies on both lines. In other words, the solution is the point where the two lines intersect.

We can solve a system of equations by graphing the lines on the same coordinate plane and reading the intersection point from the graph.

This method most often offers only approximate solutions. It is exact only when the x and y values of the solution are integers. However, this method is a great at offering a visual representation of the system of equations and demonstrates that the solution to a system of equations is the intersection of the two lines in the system.

Example 2

(The equations are in slope-intercept form)

Solve the following system of equations by graphing.

$$y = 3x - 5$$


$$y = -2x + 5$$

Solution

Graph both lines on the same coordinate axis using any method you like.

In this case, let's make a table of values for each line.

Line 1 $y = 3x - 5$



Line 2 $y = -2x + 5$

x	y
1	3
2	1

Answer The solution to the system is given by the intersection point of the two lines. The graph shows that the lines intersect at point $(2, 1)$. So the solution is $x = 2, y = 1$ or $(2, 1)$.

Example 3

(The equations are in standard form)


Solve the following system of equations by graphing

$$\begin{aligned} 2x + 3y &= 6 \\ 4x - y &= -2 \end{aligned}$$

Solution

Graph both lines on the same coordinate axis using your method of choice.

Here we will graph the lines by finding the x - and y -intercepts of each of the lines.



Line 1 $2x + 3y = 6$

x -intercept set $y = 0 \Rightarrow 2x = 6 \Rightarrow x = 3$ which results in point $(3, 0)$.

y -intercept set $x = 0 \Rightarrow 3y = 6 \Rightarrow y = 2$ which results in point $(0, 2)$.

Line 2 $-4x + y = 2$

x -intercept: set $y = 0 \Rightarrow -4x = 2 \Rightarrow x = -\frac{1}{2}$ which results in point $(-\frac{1}{2}, 0)$.


y -intercept: set $x = 0 \Rightarrow y = 2$ which results in point $(0, 2)$

Answer The graph shows that the lines intersect at point $(0, 2)$. Therefore, the solution to the system of equations is $x = 0, y = 2$.

Example 4:

Solve the following system by graphing.

$$\begin{aligned} y &= 3 \\ x + y &= 2 \end{aligned}$$



Line 1 $y = 3$ is a horizontal line passing through point $(0, 3)$.

Line 2 $x + y = 2$

x -intercept: $(2, 0)$

y -intercept: $(0, 2)$

Answer The graph shows that the solution to this system is $(-1, 3)$ $x = -1, y = 3$.

These examples are great at demonstrating that the solution to a system of linear equations means the point at which the lines intersect. This is, in fact, the greatest strength of the graphing method because it offers a very visual representation of system of equations and its solution. You can see, however, that determining a solution from a graph would require very careful graphing of the lines, and is really only practical when you are certain that the solution gives integer values for x and y . In most cases, this method can only offer approximate solutions to systems of equations. For exact solutions other methods are necessary.

Solving a System of Equations Using the Graphing Calculator

A graphing calculator can be used to find or check solutions to a system of equations. In this section, you learned that to solve a system graphically, you must graph the two lines on the same coordinate axes and find the point of intersection. You can use a graphing calculator to graph the lines as an alternative to graphing the equations by hand.

Example 6

Solve the following system of equations using a graphing calculator.

$$\begin{aligned}x - 3y &= 4 \\2x + 5y &= 8\end{aligned}$$

In order to input the equations into the calculator, they must be written in slope-intercept form (i.e., $y = mx + b$ form), or at least you must isolate y .

$$\begin{aligned}x - 3y &= 4 & y &= \frac{1}{3}x - \frac{4}{3} \\ \Rightarrow & & & \\ 2x + 5y &= 8 & y &= \frac{-2}{5}x - \frac{8}{5}\end{aligned}$$


Press the **[y=]** button on the graphing calculator and enter the two functions as:

$$Y_1 = \frac{x}{3} - \frac{4}{3}$$

$$T_2 = -\frac{2x}{3} - \frac{8}{5}$$

Now press **[GRAPH]**. The window below is set to $-5 \leq x \leq 10$ and $-5 \leq y \leq 5$.

The first screen below shows the screen of a TI-83 family graphing calculator with these lines graphed.



There are a few different ways to find the intersection point.

Option 1 Use **[TRACE]** and move the cursor with the arrows until it is on top of the intersection point. The values of the coordinate point will be on the bottom of the screen. The second screen above shows the values to be $X = 4.0957447$ and $Y = .03191489$.

Use the **[ZOOM]** function to zoom into the intersection point and find a more accurate result. The third screen above shows the system of equations after zooming in several times. A more accurate solution appears to be $X = 4$ and $Y = 0$.

Option 2 Look at the table of values by pressing **[2nd]** **[GRAPH]**. The first screen below shows a table of values for this system of equations. Scroll down until the values for x and y are the same. In this case this occurs at $X = 4$ and $Y = 0$.

Use the **[TBLSET]** function to change the starting value for your table of values so that it is close to the intersection point and you don't have to scroll too long. You can also improve the accuracy of the solution by taking smaller values of Table 1.



Option 3 Using the **[2nd]** **[TRACE]** function gives the screen in the second screen above.

Scroll down and select intersect.

The calculator will display the graph with the question **[FIRSTCURVE]?** Move the cursor along the first curve until it is close to the intersection and press **[ENTER]**.

The calculator now shows **[SECONDCURVE]?**

Move the cursor to the second line (if necessary) and press **[ENTER]**.

The calculator displays **[GUESS]?**

Press **[ENTER]** and the calculator displays the solution at the bottom of the screen (see the third screen above).

The point of intersection is $X = 4$ and $Y = 0$.

Notes:

- When you use the "intersect" function, the calculator asks you to select **[FIRSTCURVE]?** and **[SECONDCURVE]?** in case you have more than two graphs on the screen. Likewise, the **[GUESS]?** is requested in case the curves have more than one intersection. With lines you only get one point of intersection, but later in your mathematics studies you will work with curves that have multiple points of intersection.
- Option 3 is the only option on the graphing calculator that gives an exact solution. Using trace and table give you approximate solutions.


Solve Real-World Problems Using Graphs of Linear Systems

Consider the following problem

Peter and Nadia like to race each other. Peter can run at a speed of 5 feet per second and Nadia can run at a speed of 6 feet per second. To be a good sport Nadia likes to give Peter a head start of 20 feet. How long does Nadia take to catch up with Peter? At what distance from the start does Nadia catch up with Peter?

Draw a sketch

At time, $t = 0$:



Formulas

Lets define two variables in this problem.

t = the time from when Nadia starts running

d = the distance of the runners from the starting point.

Since we have two runners we need to write equations for each of them. This will be the **system of equations** for this problem.

Here we use the formula distance = speed \times time

Nadias equation $d = 6t$


Peters equation $d = 5t + 20$

(Remember that Peter was already 20 feet from the starting point when Nadia started running.)

Lets graph these two equations on the same coordinate graph.

Time should be on the horizontal axis since it is the independent variable.


Distance should be on the vertical axis since it is the dependent variable.




We can use any method for graphing the lines. In this case, we will use the slope-intercept method since it makes more sense physically.

To graph the line that describes Nadias run, start by graphing the y -intercept $(0, 0)$. If you do not see that this is the y -intercept, try plugging in the test-value of $x = 0$.


The slope tells us that Nadia runs 6 feet every one second so another point on the line is $(1, 6)$. Connecting these points gives us Nadias line.



To graph the line that describes Peters run, again start with the y -intercept. In this case, this is the point $(0,20)$. The slope tells us that Peter runs 5 feet every one second so another point on the line is $(1,25)$. Connecting these points gives us Peters line.



In order to find when and where Nadia and Peter meet, we will graph both lines on the same graph and extend the lines until they cross. The crossing point is the solution to this problem.



The graph shows that Nadia and Peter meet 20 seconds after Nadia starts running and 120 feet from the starting point.

Review Questions

Determine which ordered pair satisfies the system of linear equations.

1. $y = 3x - 2$
 $y = -x$
 - (a) $(1, 4)$
 - (b) $(2, 9)$
 - (c) $(\frac{1}{2}, -\frac{1}{2})$
2. $y = 2x - 3$
 $y = x + 5$
 - (a) $(8, 13)$
 - (b) $(-7, 6)$
 - (c) $(0, 4)$
3. $2x + y = 8$
 $5x + 2y = 10$
 - (a) $(-9, 1)$
 - (b) $(-6, 20)$
 - (c) $(14, 2)$
4. $3x + 2y = 6$
 $y = \frac{x}{2} - 3$
 - (a) $(3, -\frac{3}{2})$
 - (b) $(-4, 3)$
 - (c) $(\frac{1}{2}, 4)$

Solve the following systems using the graphing method.

5. $y = x + 3$
 $y = -x + 3$
6. $y = 3x - 6$
 $y = -x + 6$
7. $2x = 4$
 $y = -3$
8. $y = -x + 5$
 $-x + y = 1$
9. $x + 2y = 8$
 $5x + 2y = 0$
10. $3x + 2y = 12$
 $4x - y = 5$
11. $5x + 2y = -4$
 $x - y = 2$
12. $2x + 4 = 3y$
 $x - 2y + 4 = 0$
13. $y = \frac{x}{2} - 3$
 $2x - 5y = 5$
14. $y = 4$
 $x = 8 - 3y$
15. Solve the following problems by using the graphing method.
16. Marys car is 10 years old and has a problem. The repair man indicates that it will cost her \$1200 to repair her car. She can purchase a different, more efficient car for \$4500. Her present car averages about \$2000 per year for gas while the new car would average about \$1500 per year. Find the number of years for when the total cost of repair would equal the total cost of replacement.

17. Juan is considering two cell phone plans. The first company charges \$120 for the phone and \$30 per month for the calling plan that Juan wants. The second company charges \$40 for the same phone, but charges \$45 per month for the calling plan that Juan wants. After how many months would the total cost of the two plans be the same?
18. A tortoise and hare decide to race 30 feet. The hare, being much faster, decided to give the tortoise a head start of 20 feet. The tortoise runs at 0.5 feet/sec and the hare runs at 5.5 feet per second. How long will it be until the hare catches the tortoise?

Review Answers

1. (c)
2. (a)
3. (b)
4. (a)
5. (0, 3)
6. (3, 3)
7. (2, -3)
8. (2, 3)
9. (-2, 5)
10. (2, 3)
11. (0, -2)
12. (4, 4)
13. (20, 7)
14. (-4, 4)
15. 6.6 years
16. 5.33 months
17. 4.0 seconds

7.2 Solving Linear Systems by Substitution

Learning Objectives

- Solve systems of equations with two variables by substituting for either variable.
- Manipulate **standard form** equations to isolate a single variable.
- Solve real-world problems using systems of equations.
- Solve mixture problems using systems of equations.

Introduction

In this lesson, we will learn to solve a system of two equations using the method of substitution.

Solving Linear Systems Using Substitution of Variable Expressions

Lets look again at the problem involving Peter and Nadia racing.

Peter and Nadia like to race each other. Peter can run at a speed of 5 feet per second and Nadia can run at a speed of 6 feet per second. To be a good sport Nadia likes to give Peter a head start of 20 feet. How long does Nadia take to catch up with Peter? At what distance from the start does Nadia catch up with Peter?

In that example, we came up with two equations.

$$\begin{array}{ll} \text{Nadia's equation} & d = 6t \\ \text{Peter's equation} & d = 5t + 20 \end{array}$$

We have seen that each relationship produces its own line on a graph, but that to solve the system we find the point at which the lines intersect (Lesson 1). At that point the values for d and t satisfy **both** relationships.

In this simple example, this means that the d in Nadias equation is the same as the d in Peters. We can set the two equations equal to each other to solve for t .

$$\begin{array}{ll} 6t = 5t + 20 & \text{Subtract } 5t \text{ from both sides.} \\ t = 20 & \text{Substitute this value for } t \text{ into Nadia's equation.} \\ d = 6 \cdot 20 = 120 & \end{array}$$

Even if the equations are not so obvious, we can use simple algebraic manipulation to find an expression for one variable in terms of the other. We can rearrange Peters equation to isolate t .

$$\begin{array}{ll} d = 5t + 20 & \text{Subtract 20 from both sides.} \\ d - 20 = 5t & \text{Divide by 5.} \\ \frac{d - 20}{5} = t & \end{array}$$

We can now *substitute* this expression for t into Nadias equation ($d = 6t$) to solve it.

$$\begin{array}{ll}
 d = 6\left(\frac{d-20}{5}\right) & \text{Multiply both sides by 5.} \\
 5d = 6(d-20) & \text{Distribute the 6.} \\
 5d = 6d - 120 & \text{Subtract } 6d \text{ from both sides.} \\
 -d = -120 & \text{Divide by } -1. \\
 d = 120 & \text{Substitute value for } d \text{ into our expression for } t. \\
 t = \frac{120-20}{5} = \frac{100}{5} = 20 &
 \end{array}$$

We find that Nadia and Peter meet 20 seconds after they start racing, at a distance of 120 yards away.

The method we just used is called the **Substitution Method**. In this lesson, you will learn several techniques for isolating variables in a system of equations, and for using the expression you get for solving systems of equations that describe situations like this one.

Example 1

Let us look at an example where the equations are written in **standard form**.

Solve the system

$$\begin{array}{l}
 2x + 3y = 6 \\
 -4x + y = 2
 \end{array}$$

Again, we start by looking to isolate one variable in either equation. If you look at the second equation, you should see that the coefficient of y is 1. It makes sense to use this equation to solve for y .

Solve the second equation for the y variable:

$$\begin{array}{ll}
 -4x + y = 2 & \text{Add } 4x \text{ to both sides.} \\
 y = 2 + 4x &
 \end{array}$$

Substitute this expression into the second equation.

$$\begin{array}{ll}
 2x + 3(2 + 4x) = 6 & \text{Distribute the 3.} \\
 2x + 6 + 12x = 6 & \text{Collect like terms.} \\
 14x + 6 = 6 & \text{Subtract 6 from both sides.} \\
 14x = 0 & \\
 x = 0 &
 \end{array}$$

Substitute back into our expression for y .

$$y = 2 + 4 \cdot 0 = 2$$

As you can see, we end up with the same solution ($x = 0, y = 2$) that we found when we graphed these functions (Lesson 7.1). As long as you are careful with the algebra, the substitution method can be a very efficient way to solve systems.

Next, consider a more complicated example. In the following example the solution gives fractional answers for both x and y , and so would be very difficult to solve by graphing alone!

Example 2

Solve the system

$$\begin{aligned}2x + 3y &= 3 \\ 2x - 3y &= -1\end{aligned}$$

Again, we start by looking to isolate one variable in either equation. Right now it doesn't matter which equation we use or which variable we solve for.

Solve the first equation for x

$$\begin{aligned}2x + 3y &= 3 && \text{Subtract } 3y \text{ from both sides.} \\ 2x &= 3 - 3y && \text{Divide both sides by 2.} \\ x &= \frac{3 - 3y}{2}\end{aligned}$$

Substitute this expression into the second equation.

$$\begin{aligned}2 \cdot \frac{1}{2}(3 - 3y) - 3y &= -1 && \text{Cancel the fraction and rewrite terms.} \\ 3 - 3y - 3y &= -1 && \text{Collect like terms.} \\ 3 - 6y &= -1 && \text{Subtract 3 from both sides.} \\ -6y &= -4 && \text{Divide by } -6. \\ y &= \frac{2}{3}\end{aligned}$$

Substitute into the expression and solve for x .

$$\begin{aligned}x &= \frac{1}{2} \left(3 - 3 \left(\frac{2}{3} \right) \right) \\ x &= \frac{1}{2}\end{aligned}$$

So our solution is, $x = \frac{1}{2}, y = \frac{2}{3}$. You can see why the graphical solution $(\frac{1}{2}, \frac{2}{3})$ might be difficult to read accurately.

Solving Real-World Problems Using Linear Systems

There are many situations where we can use simultaneous equations to help solve real-world problems. We may be considering a purchase. For example, trying to decide whether it is cheaper to buy an item online where you pay shipping or at the store where you do not. Or you may wish to join a CD music club, but do not know if you would


really save any money by buying a new CD every month in that way. One example with which we are all familiar is considering phone contracts. Lets look at an example of that now.

Example 3

Anne is trying to choose between two phone plans. The first plan, with Vendafone costs \$20 per month, with calls costing an additional 25 cents per minute. The second company, Sellnet, charges \$40 per month, but calls cost only 8 cents per minute. Which should she choose?

Annes choice will depend upon how many minutes of calls she expects to use each month. We start by writing two equations for the cost in dollars in terms of the minutes used. Since the number of minutes is the independent variable, it will be our x . Cost is dependent on minutes. The *cost per month* is the *dependent* variable and will be assigned y .

For Vendafone	$y = 0.25x + 20$
For Sellnet	$y = 0.08x + 40$



By writing the equations in slope-intercept form ($y = mx + b$) you can visualize the situation in a simple sketched graph, shown right. The line for Vendafone has an intercept of 20 and a slope of 0.25. The Sellnet line has an intercept of 40 and a slope of 0.08 (which is roughly a third of the Vendafone line). In order to help Anne decide which to choose, we will determine where the two lines cross, by solving the two equations as a system. Since equation one gives us an expression for $y(0.25x + 20)$, we can substitute this expression directly into equation two.

$0.25x + 20 = 0.08x + 40$	Subtract 20 from both sides.
$0.25x = 0.08x + 20$	Subtract $0.08x$ from both sides.
$0.17x = 20$	Divide both sides by 0.17.
$x = 117.65$ minutes	Rounded to two decimal places.

We can now use our sketch, plus this information to provide an answer:

If Anne will use 117 minutes or less every month, she should choose Vendafone. If she plans on using 118 or more minutes, she should choose Sellnet.

Mixture Problems

Systems of equations crop up frequently when considering chemicals in solutions, and can even be seen in things like mixing nuts and raisins or examining the change in your pocket! Lets look at some examples of these.

Example 4

Nadia empties her purse and finds that it contains only nickels (worth 5 cents each) and dimes (worth 10 cents each). If she has a total of 7 coins and they have a combined value of 55 cents, how many of each coin does she have?

Since we have two types of coins, let's call the number of nickels x and the number of dimes will be our y . We are given two key pieces of information to make our equations, the number of coins and their value.

Number of coins equation	$x + y = 7$	(number of nickels) + (number of dimes)
The value equation	$5x + 10y = 55$	Since nickels are worth five cents and dimes ten cents

We can quickly rearrange the first equation to isolate x .



Image courtesy of Kevin@flickr.com/creativecommons

$$\begin{aligned}
 x &= 7 - y \\
 5(7 - y) + 10y &= 55 \\
 35 - 5y + 10y &= 55 \\
 35 + 5y &= 55 \\
 5y &= 20 \\
 y &= 4 \\
 +4 &= 7 \\
 x &= 3
 \end{aligned}$$

Now substitute into equation two.
 Distribute the 5.
 Collect like terms.
 Subtract 35 from both sides.
 Divide by 5.
 Substitute back into equation one.
 Subtract 4 from both sides.

Solution

Nadia has 3 nickels and 4 dimes.

Sometimes the question asks you to determine (from concentrations) how much of a particular substance to use. The substance in question could be something like coins as above, or it could be a chemical in solution, or even heat. In such a case, you need to know the amount of whatever substance is in each part. There are several common

situations where to get one equation you simply add two given quantities, but to get the second equation you need to use a **product**. Three examples are below.

TABLE 7.1:

Type of Mixture	First Equation	Second Equation
Coins (items with \$ value)	Total number of items ($n_1 n_2$)	Total value (item value \times no. of items)
Chemical solutions	Total solution volume ($V_1 + V_2$)	Amount of solute (vol \times concentration)
Density of two substances	Total amount or volume of mix	Total mass (volume \times density)

For example, when considering mixing chemical solutions, we will most likely need to consider the total amount of solute in the individual parts and in the final mixture. A solute is simply the chemical that is dissolved in a solution. An example of a solute is salt when added to water to make a brine. Even if the chemical is more exotic, we are still interested in the **total amount** of that chemical in each part. To find this, simply multiply the amount of the mixture by the **fractional concentration**. To illustrate, let's look at an example where you are given amounts relative to the whole.

Example 5

A chemist needs to prepare 500 ml of copper-sulfate solution with a 15% concentration. In order to do this, he wishes to use a high concentration solution (60%) and dilute it with a low concentration solution (5%). How much of each solution should he use?

To set this problem up, we first need to define our variables. Our unknowns are the amount of concentrated solution (x) and the amount of dilute solution (y). We will also convert the percentages (60%, 15% and 5%) into decimals (0.6, 0.15 and 0.05). The two pieces of critical information we need is the final volume (500 ml) and the final amount of solute (15% of 500 ml = 75 ml). Our equations will look like this.

$$\begin{array}{ll} \text{Volume equation} & x + y = 500 \\ \text{Solute equation} & 0.6x + 0.05y = 75 \end{array}$$

You should see that to isolate a variable for substitution it would be easier to start with equation one.



Flickr.com/creativecommons

$$\begin{aligned}
 x + y &= 500 \\
 x &= 500 - y \\
 0.6(500 - y) + 0.05y &= 75 \\
 300 - 0.6y + 0.05y &= 75 \\
 300 - 0.55y &= 75 \\
 -0.55y &= -225 \\
 y &= 409 \text{ ml} \\
 x &= 500 - 409 = 91 \text{ ml}
 \end{aligned}$$


Subtract y from both sides.
 Now substitute into equation two.
 Distribute the 6.
 Collect like terms.
 Subtract 300 from both sides.
 Divide both sides by -0.55 .
 Substitute back into equation for x .

Solution


The chemist should mix 91 ml of the 60% solution with 409 ml of the 5% solution.

Review Questions


- Solve the system:
 $x + 2y = 9$
 $3x + 5y = 20$
- solve the system.
 $x - 3y = 10$
 $2x + y = 13$
- Of the two non-right angles in a right angled triangle, one measures twice that of the other. What are the angles?



- The sum of two numbers is 70. They differ by 11. What are the numbers?
- A rectangular field is enclosed by a fence on three sides and a wall on the fourth side. The total length of the fence is 320 yards. If the field has a total perimeter of 400 yards, what are the dimensions of the field?



- A ray cuts a line forming two angles. The difference between the two angles is 18° . What does each angle measure?



7. I have \$15 and wish to buy five pounds of mixed nuts for a party. Peanuts cost \$2.20 per pound. Cashews cost \$4.70 per pound. How many pounds of each should I buy?
8. A chemistry experiment calls for one liter of sulfuric acid at a 15% concentration, but the supply room only stocks sulfuric acid in concentrations of 10% and in 35%. How many liters of each should be mixed to give the acid needed for the experiment?
9. Bachellet wants to know the density of her bracelet, which is a mix of gold and silver. Density is total mass divided by total volume. The density of gold is 19.3 g/cc and the density of silver is 10.5 g/cc . The jeweler told her that the volume of silver used was 10 cc and the volume of gold used was 20 cc . Find the combined density of her bracelet.

Review Answers

1. $x = -5, y = 7$
2. $x = 7, y = -1$
3. $x = 30^\circ, y = 60^\circ$
4. 29.5 and 40.5
5. $x = 120$ yards, $y = 80$ yards
6. $x = 81^\circ, y = 99^\circ$
7. 3.4 pounds of peanuts, 1.6 pounds of cashews
8. 0.8 liters of 10%, 0.2 liters of 35%
9. 16.4 g/cc

7.3 Solving Linear Systems by Elimination through Addition or Subtraction

Learning Objectives

- Solve a linear system of equations using elimination by addition.
- Solve a linear system of equations using elimination by subtraction.
- Solve real-world problems using linear systems by elimination.

Introduction

In this lesson, we will look at using simple addition and subtraction to simplify our system of equations to a single equation involving a single variable. Because we go from two unknowns (x and y) to a single unknown (either x or y) this method is often referred to as **solving by elimination**. We eliminate one variable in order to make our equations solvable! To illustrate this idea, let's look at the simple example of buying apples and bananas.

Example 1

If one apple plus one banana costs \$1.25 and one apple plus two bananas costs \$2.00, how much does it cost for one banana? One apple?

It shouldn't take too long to discover that each banana costs \$0.75. You can see this by looking at the difference between the two situations. Algebraically, using a and b as the cost for apples and bananas, we get the following equations.

$$\begin{aligned}a + b &= 1.25 \\ a + 2b &= 2.00\end{aligned}$$

If you look at the difference between the two equations you see that the difference in items purchased is one banana, and the difference in money paid is 75 cents. So one banana costs 75 cents.

$$(a + 2b) - (a + b) = 2.00 - 1.25 \Rightarrow b = 0.75$$

To find out how much one apple costs, we subtract \$0.75 from the cost of one apple and one banana. So an apple costs 50 cents.

$$a + 0.75 = 1.25 \Rightarrow a = 1.25 - 0.75 \Rightarrow a = 0.50$$

To solve systems using addition and subtraction, we will be using exactly this idea. By looking at the sum or difference of the two equations, we can determine a value for one of the unknowns.

Solving Linear Systems Using Addition of Equations

Often considered the easiest and most powerful method of solving systems of equations, the addition (or elimination) method requires us to combine two equations in such a way that the resulting equation has only one variable. We can

then use simple linear algebra methods of solving for that variable. If required, we can always substitute the value we get for that variable back in either one of the original equations to solve for the remaining unknown variable.

Example 2

Solve the system by addition:

$$\begin{aligned}3x + 2y &= 11 \\5x - 2y &= 13\end{aligned}$$

We will add everything on the left of the equals sign from both equations, and this will be equal to the sum of everything on the right.

$$(3x + 2y) + (5x - 2y) = 11 + 13 \Rightarrow 8x = 24 \Rightarrow x = 3$$

A simpler way to visualize this is to keep the equations as they appear above, and to add in columns. However, just like adding units tens and hundreds, you **MUST** keep x 's and y 's in their own columns. You may also wish to use terms like $0y$ as a placeholder!

$$\begin{array}{r}3x + 2y = 11 \\+ (3x - 2y) = 13 \\ \hline8x + 0y = 24\end{array}$$

Again we get $8x = 24$ or $x = 3$.

To find a value for y we simply substitute our value for x back in.

Substitute $x = 3$ into the second equation.

$$\begin{aligned}5 \cdot 3 - 2y &= 13 \\-2y &= -2 \\y &= 1\end{aligned}$$

Since $5 \times 3 = 15$, we subtract 15 from both sides.

Divide by 2 to get the value for y .

The first example has a solution at $x = 3$ and $y = 1$. You should see that the method of addition works when the coefficients of one of the variables are opposites. In this case it is the coefficients of y that are opposites, being $+2$ in the first equation and -2 in the second.

There are other, similar, methods we can use when the coefficients are not opposites, but for now let's look at another example that can be solved with the method of addition.

Example 3



Dagwood21/www.flickr.com/creativecommons

Andrew is paddling his canoe down a fast moving river. Paddling downstream he travels at 7 miles per hour, relative to the river bank. Paddling upstream, he moves slower, traveling at 1.5 miles per hour. If he paddles equally hard in both directions, calculate, in miles per hour, the speed of the river and the speed Andrew would travel in calm water.

Step One First, we convert our problem into equations. We have two unknowns to solve for, so we will call the speed that Andrew paddles at x , and the speed of the river y . When traveling downstream, Andrew's speed is boosted by the river current, so his total speed is the canoe speed plus the speed of the river ($x + y$). Upstream, his speed is hindered by the speed of the river. His speed upstream is $(x - y)$.

Downstream Equation

$$x + y = 7$$

Upstream Equation

$$x - y = 1.5$$

Step Two Next, we are going to eliminate one of the variables. If you look at the two equations, you can see that the coefficient of y is $+1$ in the first equation and -1 in the second. Clearly $(+1) + (-1) = 0$, so this is the variable we will eliminate. To do this we simply add equation 1 to equation 2. We must be careful to collect like terms, and that everything on the left of the equals sign stays on the left, and everything on the right stays on the right:

$$(x + y) + (x - y) = 7 + 1.5 \Rightarrow 2x = 8.5 \Rightarrow x = 4.25$$

Or, using the column method we used in example one.

$$\begin{array}{r} x + y = 7 \\ + (x - y) = 1.5 \\ \hline 2x + 0y = 8.5 \end{array}$$

Again you see we get $2x = 8.5$, or $x = 4.25$. To find a corresponding value for y , we plug our value for x into either equation and isolate our unknown. In this example, we'll plug it into the first equation.

Substitute $x = 4.25$ into the second equation:

$$\begin{aligned} 4.25 + y &= 7 \\ y &= 2.75 \end{aligned}$$

Subtract 4.25 from both sides.

Solution

Andrew paddles at 4.25 miles per hour. The river moves at 2.75 miles per hour.

Solving Linear Systems Using Subtraction of Equations

Another, very similar method for solving systems is subtraction. In this instance, you are looking to have identical coefficients for x or y (including the sign) and then subtract one equation from the other. If you look at Example one you can see that the coefficient for x in both equations is $+1$. You could have also used the method of subtraction.

$$(x + y) - (x - y) = 200 - 80 \Rightarrow 2y = 120 \Rightarrow y = 60$$

or

$$\begin{array}{r} x + y = 200 \\ + (x - y) = -80 \\ \hline 0x + 2y = 120 \end{array}$$

So again we get $y = 60$, from which we can determine x . The method of subtraction looks equally straightforward, and it is so long as you remember the following:

1. Always put the equation you are subtracting in parentheses, and distribute the negative.
2. Don't forget to subtract the numbers on the right hand side.
3. Always remember that subtracting a negative is the same as adding a positive.

Example 4

Peter examines the coins in the fountain at the mall. He counts 107 coins, all of which are either pennies or nickels. The total value of the coins is \$3.47. How many of each coin did he see?

We have two types of coins. Lets call the number of pennies x and the number of nickels y . The total value of pennies is just x , since they are worth one cent each. The total value of nickels is $5y$. We are given two key pieces of information to make our equations. The number of coins and their value.

Number of Coins Equation	$x + y = 107$	(number of pennies) + (number of nickels)
The Value Equation:	$x + 5y = 347$	pennies are worth 1c, nickels are worth 5c.

We will jump straight to the subtraction of the two equations.

$$\begin{array}{r} x + y = 107 \\ + (x + 5y) = -347 \\ \hline 4y = -240 \end{array}$$

Let's substitute this value back into the first equation.

$$x + 60 = 107$$

$$x = 47$$

Subtract 60 from both sides.

So Peter saw 47 pennies (worth 47 cents) and 60 nickels (worth \$3.00) for a total of \$3.47.

We have now learned three techniques for solving systems of equations.

1. Graphing
2. Substitution
3. Elimination

You should be starting to gain an understanding of which method to use when given a particular problem. For example, **graphing** is a good technique for seeing what the equations are doing, and when one service is less expensive than another. Graphing alone may not be ideal when an exact numerical solution is needed.

Similarly, **substitution** is a good technique when one of the coefficients in your equation is $+1$ or -1 .

Addition or **subtraction** is ideal when the coefficient of one of the variables matches the coefficient of the same variable in the other equation. In the next lesson, we will learn the last technique for solving systems of equations exactly, when none of the coefficients match and the coefficient is not one.

Review Questions

1. Solve the system:
 $3x + 4y = 2.5$
 $5x - 4y = 25.5$
2. Solve the system
 $5x + 7y = -31$
 $5x - 9y = 17$
3. Solve the system
 $3y - 4x = -33$
 $5x - 3y = 40.5$
4. Nadia and Peter visit the candy store. Nadia buys three candy bars and four fruit roll-ups for \$2.84. Peter also buys three candy bars, but can only afford one additional fruit roll-up. His purchase costs \$1.79. What is the cost of each candy bar and each fruit roll-up?
5. A small plane flies from Los Angeles to Denver with a tail wind (the wind blows in the same direction as the plane) and an air-traffic controller reads its ground-speed (speed measured relative to the ground) at 275 miles per hour. Another, identical plane, moving in the opposite direction has a ground-speed of 227 miles per hour. Assuming both planes are flying with identical air-speeds, calculate the speed of the wind.
6. An airport taxi firm charges a pick-up fee, plus an additional per-mile fee for any rides taken. If a 12 miles journey costs \$14.29 and a 17 miles journey costs \$19.91, calculate:
 - (a) the pick-up fee
 - (b) the per-mile rate
 - (c) the cost of a seven mile trip
7. Calls from a call-box are charged per minute at one rate for the first five minutes, then a different rate for each additional minute. If a seven minute call costs \$4.25 and a 12 minute call costs \$5.50, find each rate.
8. A plumber and a builder were employed to fit a new bath, each working a different number of hours. The plumber earns \$35 per hour, and the builder earns \$28 per hour. Together they were paid \$330.75, but the plumber earned \$106.75 more than the builder. How many hours did each work?

9. Paul has a part time job selling computers at a local electronics store. He earns a fixed hourly wage, but can earn a bonus by selling warranties for the computers he sells. He works 20 hours per week. In his first week, he sold eight warranties and earned \$220. In his second week, he managed to sell 13 warranties and earned \$280. What is Pauls hourly rate, and how much extra does he get for selling each warranty?

Review Answers

1. $x = 3.5, y = -2$
2. $x = -2, y = -3$
3. $x = 7.5, y = -1$
4. Candy bars cost 48 cents each and fruit roll-ups cost 35 cents each.
5. The wind speed is 24 mph
6. (a) \$.80
(b) \$1.12
(c) \$8.64
7. 75 cents per minute for the first 5 mins, 25 cents per minute additional
8. The plumber works 6.25 hours, the builder works 4 hours
9. Paul earns a base of \$7.00 per hour

7.4 Solving Systems of Equations by Multiplication

Learning objectives

- Solve a linear system by multiplying one equation.
- Solve a linear system of equations by multiplying both equations.
- Compare methods for solving linear systems.
- Solve real-world problems using linear systems by any method.

Introduction

We have now learned three techniques for solving systems of equations.

- Graphing, Substitution and Elimination (through addition and subtraction).

Each one of these methods has both strengths and weaknesses.

- **Graphing** is a good technique for seeing what the equations are doing, and when one service is less expensive than another, but graphing alone to find a solution can be imprecise and may not be good enough when an exact numerical solution is needed.
- **Substitution** is a good technique when one of the coefficients in an equation is $+1$ or -1 , but can lead to more complicated formulas when there are no unity coefficients.
- **Addition or Subtraction** is ideal when the coefficients of either x or y match in both equations, but so far we have not been able to use it when coefficients do not match.

In this lesson, we will again look at the method of elimination that we learned in Lesson 7.3. However, the equations we will be working with will be more complicated and one can not simply add or subtract to eliminate one variable. Instead, we will first have to multiply equations to ensure that the coefficients of one of the variables are matched in the two equations.

Quick Review: Multiplying Equations

Consider the following questions

1. If 10 apples cost \$5, how much would 30 apples cost?
2. If 3 bananas plus 2 carrots cost \$4, how much would 6 bananas plus 4 carrots cost?

You can look at the first equation, and it should be obvious that each apple costs \$0.50. 30 apples should cost \$15.00.

Looking at the second equation, it is not clear what the individual price is for either bananas or carrots. Yet we know that the answer to question two is \$8.00. How?

If we look again at question one, we see that we can write the equation $10a = 5$ (a being the cost of one apple).

To find the cost of 30, we can either solve for a then multiply by 30, or we can multiply both sides of the equation by three.

$$30a = 15$$

$$a = \frac{1}{2} \text{ or } 0.5$$

Now look at the second question. We could write the equation $3b + 2c = 4$.

We see that we need to solve for $(6b + 4c)$ which is simply two times the quantity $(3b + 2c)$!

Algebraically, we are simply multiplying the entire equation by two.

$$2(3b + 2c) = 2 \cdot 4 \qquad \text{Distribute and multiply.}$$

$$6b + 4c = 8$$

So when we multiply an equation, all we are doing is multiplying every term in the equation by a fixed amount.

Solving a Linear System by Multiplying One Equation

We can multiply every term in an equation by a fixed number (a **scalar**), it is clear that we could use the addition method on a whole new set of linear systems. We can manipulate the equations in a system to ensure that the coefficients of one of the variables match. In the simplest case, the coefficient as a variable in one equation will be a multiple of the coefficient in the other equation.

Example 1

Solve the system.

$$7x + 4y = 17$$

$$5x - 2y = 11$$

It is quite simple to see that by multiplying the second equation by two the coefficients of y will be $+4$ and -4 , allowing us to complete the solution by addition.

Take two times equation two and add it to equation one. Then divide both sides by 17 to find x .

$$10x - 4y = 22$$

$$+ (7x + 4y) = 17$$

$$17x = 34$$

$$x = 2$$

Now simply substitute this value for x back into equation one.

$$7 \cdot 2 + 4y = 17$$

$$4y = 3$$

$$y = 0.75$$

Since $7 \times 2 = 14$, subtract 14 from both sides.
Divide by 4.

Example 2

Anne is rowing her boat along a river. Rowing downstream, it takes her two minutes to cover 400 yards. Rowing upstream, it takes her eight minutes to travel the same 400 yards. If she was rowing equally hard in both directions, calculate, in yards per minute, the speed of the river and the speed Anne would travel in calm water.

Step One First we convert our problem into equations. We need to know that *distance traveled* is equal to *speed* \times *time*. We have two unknowns, so we will call the speed of the river x , and the speed that Anne rows at y . When traveling downstream her total speed is the boat speed plus the speed of the river ($x + y$). Upstream her speed is hindered by the speed of the river. Her speed upstream is $(x - y)$.

$$\begin{array}{ll} \text{Downstream Equation} & 2(x + y) = 400 \\ \text{Upstream Equation} & 8(x - y) = 400 \end{array}$$

Distributing gives us the following system.

$$\begin{array}{l} 2x + 2y = 400 \\ 8x - 8y = 400 \end{array}$$

Right now, we cannot use the method of elimination as none of the coefficients match. But, if we were to multiply the top equation by four, then the coefficients of y would be $+8$ and -8 . Lets do that.

$$\begin{array}{r} 8x - 8y = 1,600 \\ + (8x - 8y) = 400 \\ \hline 16x = 2,000 \end{array}$$

Now we divide by 16 to obtain $x = 125$.

Substitute this value back into the first equation.

$$\begin{array}{ll} 2(125 + y) = 400 & \text{Divide both sides by 2.} \\ 125 + y = 200 & \text{Subtract 125 from both sides.} \\ y = 75 & \end{array}$$

Solution

Anne rows at 125 yards per minute, and the river flows at 75 yards per minute.

Solve a Linear System by Multiplying Both Equations

It is a straightforward jump to see what would happen if we have no matching coefficients and no coefficients that are simple multiples of others. Just think about the following fraction sum.

$$\frac{1}{2} + \frac{1}{3} = \frac{3}{6} + \frac{2}{6} = \frac{5}{6}$$

This is an example of finding a **lowest common denominator**. In a similar way, we can always find a lowest common multiple of two numbers (the **lowest common multiple** of 2 and 3 is 6). This way we can always find a way to multiply equations such that two coefficients match.

Example 3

Andrew and Anne both use the I-Haul truck rental company to move their belongings from home to the dorm rooms on the University of Chicago campus. I-Haul has a charge per day and an additional charge per mile. Andrew travels from San Diego, California, a distance of 2,060 miles in five days. Anne travels 880 miles from Norfolk, Virginia, and it takes her three days. If Anne pays \$840 and Andrew pays \$1845, what does I-Haul charge

a) per day?

b) per mile traveled?

First, we will set-up our equations. Again we have two unknowns, the **daily rate** (we will call this x), and the **rate per mile** (lets call this y).

$$\begin{array}{r} \text{Anne's equation} \\ \text{Andrew's Equation} \end{array} \qquad \begin{array}{r} 3x + 880y = 840 \\ 5x + 2060y = 1845 \end{array}$$

We cannot simply multiply a single equation by an integer number in order to arrive at matching coefficients. But if we look at the coefficients of x (as they are easier to deal with than the coefficients of y) we see that they both have a common multiple of 15 (in fact 15 is the **lowest common multiple**). So this time we need to multiply both equations:

Multiply Anne's equation by five:

$$15x + 4400y = 4200$$

Multiply Andrew's equation by three:

$$15x + 6180y = 5535$$

Subtract:

$$\begin{array}{r} 15x + 4400y = 4200 \\ - (15x + 6180y) = 5535 \\ \hline -1780y = -1335 \end{array}$$

Divide both sides by -1780

$$y = 0.75$$

Substitute this back into Anne's equation.

$$3x + 880(0.75) = 840$$

$$3x = 180$$

$$x = 60$$

Since $880 \times 0.75 = 660$, subtract 660 from both sides.

Divide both sides by 3.

Solution

I-Haul charges \$60 per day plus \$0.75 per mile.

Comparing Methods for Solving Linear Systems

Now that we have covered the major methods for solving linear equations, let's review. For simplicity, we will look at the four methods (we will consider **addition** and **subtraction** one method) in table form. This should help you decide which method would be better for a given situation.

TABLE 7.2:

Method:	Best used when you	Advantages:	Comment:
Graphing	don't need an accurate answer.	Often easier to see number and quality of intersections on a graph. With a graphing calculator, it can be the fastest method since you don't have to do any computation.	Can lead to imprecise answers with non-integer solutions.
Substitution	have an explicit equation for one variable (e.g. $y = 14x + 2$)	Works on all systems. Reduces the system to one variable, making it easier to solve.	You are not often given explicit functions in systems problems, thus it can lead to more complicated formulas
Elimination by Addition or Subtraction	have matching coefficients for one variable in both equations.	Easy to combine equations to eliminate one variable. Quick to solve.	It is not very likely that a given system will have matching coefficients.
Elimination by Multiplication and then Addition and Subtraction	do not have any variables defined explicitly or any matching coefficients.	Works on all systems. Makes it possible to combine equations to eliminate one variable.	Often more algebraic manipulation is needed to prepare the equations.

The table above is only a guide. You may like to use the graphical method for every system in order to better understand what is happening, or you may prefer to use the multiplication method even when a substitution would work equally well.

Example 4


Two angles are **complementary** when the sum of their angles is 90° . Angles A and B are complementary angles, and twice the measure of angle A is 9° more than three times the measure of angle B . Find the measure of each angle.

First, we write out our two equations. We will use x to be the measure of Angle A and y to be the measure of Angle B . We get the following system

$$\begin{aligned}x + y &= 90 \\2x &= 3y + 9\end{aligned}$$

The first method we will use to solve this system is the graphical method. For this we need to convert the two equations to $y = mx + b$ form

$$\begin{array}{lll}x + y = 90 & \Rightarrow & y = -x + 90 \\2x = 3y + 9 & \Rightarrow & y = \frac{2}{3}x - 3\end{array}$$



The first line has a slope of -1 and a y -intercept of 90 .

The second has a slope of $\frac{2}{3}$ and a y -intercept of -3 .

In the graph, it appears that the lines cross at around $x = 55$, $y = 35$ but it is difficult to tell exactly! Graphing by hand is not the best method if you need to know the answer with more accuracy!

Next, we will try to solve by substitution. Lets look again at the system:

$$\begin{aligned}x + y &= 90 \\2x &= 3y + 9\end{aligned}$$

We have already seen that we can solve for y with either equation in trying to solve the system graphically.

Solve the first equation for y .

$$y = 90 - x$$

Substitute into the second equation

$$\begin{aligned}2x &= 3(90 - x) + 9 \\2x &= 270 - 3x + 9 \\5x &= 270 + 9 = 279 \\x &= 55.8^\circ\end{aligned}$$

Distribute the 3.
Add $3x$ to both sides.
Divide by 5.

Substitute back into our expression for y .

$$y = 90 - 55.8 = 34.2^\circ$$

Solution

Angle A measures 55.8° . Angle B measures 34.2°

Finally, we will examine the method of elimination by multiplication.

Rearrange equation one to standard form

$$x + y = 90 \Rightarrow 2x + 2y = 180$$

Multiply equation two by two.

$$2x = 3y + 9 \Rightarrow 2x - 3y = 9$$

Subtract.

$$\begin{array}{r} 2x + 2y = 180 \\ - (2x - 3y) = -9 \\ \hline 5y = 171 \end{array}$$

Divide both sides by 5

$$y = 34.2$$

Substitute this value into the very first equation.

$$\begin{aligned} x + 34.2 &= 90 \\ x &= 55.8^\circ \end{aligned}$$

Subtract 34.2 from both sides.

Solution

Angle A measures 55.8° . Angle B measures 34.2° .

Even though this system looked ideal for substitution, the method of multiplication worked well, also. Once the algebraic manipulation was performed on the equations, it was a quick solution. You will need to decide yourself which method to use in each case you see from now on. Try to master all techniques, and recognize which technique will be most efficient for each system you are asked to solve.

Review Questions

1. Solve the following systems using multiplication.

(a) $5x - 10y = 15$

$3x - 2y = 3$

(b) $5x - y = 10$

$3x - 2y = -1$

(c) $5x + 7y = 15$

$7x - 3y = 5$

(d) $9x + 5y = 9$

$12x + 8y = 12.8$

(e) $4x - 3y = 1$

$3x - 4y = 4$

(f) $7x - 3y = -3$

$6x + 4y = 3$

2. Solve the following systems using any method.

(a) $x = 3y$

$x - 2y = -3$

(b) $y = 3x + 2$

$y = -2x + 7$

(c) $5x - 5y = 5$

$5x + 5y = 35$

(d) $y = -3x - 3$

$3x - 2y + 12 = 0$

(e) $3x - 4y = 3$

$4y + 5x = 10$

(f) $9x - 2y = -4$

$2x - 6y = 1$

- Supplementary angles are two angles whose sum is 180° . Angles A and B are supplementary angles. The measure of Angle A is 18° less than twice the measure of Angle B . Find the measure of each angle.
- A farmer has fertilizer in 5% and 15% solutions. How much of each type should he mix to obtain 100 liters of fertilizer in a 12% solution?
- A 150 yards pipe is cut to provide drainage for two fields. If the length of one piece is three yards less than twice the length of the second piece, what are the lengths of the two pieces?
- Mr. Stein invested a total of \$100,000 in two companies for a year. Company A's stock showed a 13% annual gain, while Company B showed a 3% loss for the year. Mr. Stein made an 8% return on his investment over the year. How much money did he invest in each company?
- A baker sells plain cakes for \$7 or decorated cakes for \$11. On a busy Saturday the baker started with 120 cakes, and sold all but three. His takings for the day were \$991. How many plain cakes did he sell that day, and how many were decorated before they were sold?
- Twice John's age plus five times Claire's age is 204. Nine times John's age minus three times Claire's age is also 204. How old are John and Claire?

Review Answers

- $x = 0, y = -1.5$
 - $x = 3, y = 5$
 - $x = 1.25, y = 1.25$
 - $x = \frac{2}{3}, y = \frac{3}{5}$

- (e) $x = -\frac{8}{7}, y = -\frac{13}{7}$
- (f) $x = -\frac{3}{46}, y = \frac{39}{46}$
- 2. (a) $x = -9, y = -3$
- (b) $x = 1, y = 5$
- (c) $x = 4, y = 3$
- (d) $x = -2, y = 3$
- (e) $x = \frac{13}{8}, y = \frac{15}{32}$
- (f) $x = -\frac{13}{25}, y = -\frac{17}{50}$
- 3. $A = 114^\circ, B = 66^\circ$
- 4. 30 liters of 5%, 70 liters of 15%
- 5. 51 yards and 99 yards
- 6. \$68,750 in Company A, \$31,250 in Company B
- 7. 74 plain, 43 decorated
- 8. John is 32, Claire is 28

7.5 Special Types of Linear Systems

Learning Objectives


- Identify and understand what is meant by an **inconsistent linear system**.
- Identify and understand what is meant by a **consistent linear system**.
- Identify and understand what is meant by a **dependent linear system**.

Introduction

As we saw in Section 7.1, a system of linear equations is a set of linear equations which must be solved together. The lines in the system can be graphed together on the same coordinate graph and the solution to the system is the point at which the two lines intersect.

Determining the Type of System Graphically


If we graph two lines on the same coordinate plane, three different situations may occur.



Case 1: The two lines intersect at a single point; hence the lines are not parallel.

If these lines were to represent a system of equations, the system would have exactly one solution, where the lines cross.


A system with exactly one solution is called a **consistent system**.



Case 2: The two lines do not intersect. The two lines are parallel.

If the lines represent a system of equations, then the system has no solutions.

A system with no solutions is called an **inconsistent system**.



Case 3: The two lines are identical. They intersect at all points on the line.

If this were a system of equations it would have an **infinite number** of solutions. Reason being, the two equations are really the same.

Such a system is called a **dependent system**.

To identify a system as **consistent**, **inconsistent**, or **dependent**, we can graph the two lines on the same graph and match the system with one of the three cases we discussed.

Another option is to write each line in slope-intercept form and compare the slopes and y -intercepts of the two lines. To do this we must remember that:

- Lines that intersect have different slopes.
- Lines that are parallel have the same slope but different y -intercepts.
- Lines that have the same slope and the same y -intercepts are identical.

Example 1

Determine whether the following system has exactly one solution, no solutions, or an infinite number of solutions.

$$\begin{aligned}y &= 3x + 2 \\y &= -2x + 1\end{aligned}$$

Solution

The equations are already in slope-intercept form. The slope of the first equation is 3 and the slope of the second equation is -2 . Since the slopes are different, the lines must intersect at a single point. Therefore, the system has exactly one solution. This is a **consistent system**.

Example 2

Determine whether the following system has exactly one solution, no solutions, or an infinite number of solutions.

$$\begin{aligned}2x - 5y &= 2 \\4x + y &= 5\end{aligned}$$

Solution

We must rewrite the equations so they are in slope-intercept form.

$$\begin{array}{rcl} 2x - 5y = 2 & & -5y = -2x + 2 & & y = \frac{2}{5}x - \frac{2}{5} \\ & \Rightarrow & & \Rightarrow & \\ 4x + y = 5 & & y = -4x + 5 & & y = -4x + 5 \end{array}$$

The slopes of the two equations are different. Therefore, the lines must cross at a single point, and the system has exactly one solution. This is a **consistent system**.

Example 3

Determine whether the following system has exactly one solution, no solutions, or an infinite number of solutions.

$$\begin{array}{l} 3x = 5 - 4y \\ 6x + 8y = 7 \end{array}$$

Solution

We must rewrite the equations so they are in slope-intercept form.

$$\begin{array}{rcl} 3x = 5 - 4y & & 4y = -3x + 5 & & y = \frac{-3}{4}x - \frac{5}{4} \\ & \Rightarrow & & \Rightarrow & \\ 6x + 8y = 7 & & 8y = -6x + 7 & & y = \frac{-3}{4}x + \frac{7}{8} \end{array}$$

The slopes of the two equations are the same but the y -intercepts are different, therefore the lines never cross and the system has no solutions. This is an **inconsistent system**.

Example 4

Determine whether the following system has exactly one solution, no solutions, or an infinite number of solutions.

$$\begin{array}{l} x + y = 3 \\ 3x + 3y = 9 \end{array}$$

Solution

We must rewrite the equations so they are in slope-intercept form.

$$\begin{array}{rcl} x + y = 3 & & y = -x + 3 & & y = -x + 3 \\ & \Rightarrow & & \Rightarrow & \\ 3x + 3y = 9 & & 3y = -3x + 9 & & y = -x + 3 \end{array}$$

The lines are identical. Therefore the system has an infinite number of solutions. It is a **dependent system**.

Determining the Type of System Algebraically

A third option for identifying systems as **consistent**, inconsistent or dependent is to solve the system algebraically using any method and use the result as a guide.

Example 5

Solve the following system of equations. Identify the system as consistent, inconsistent or dependent.

$$\begin{aligned} 10x - 3y &= 3 \\ 2x + y &= 9 \end{aligned}$$

Solution

Lets solve this system using the substitution method.

Solve the second equation for the y variable.

$$2x + y = 9 \Rightarrow y = -2x + 9$$

Substitute for y in the first equation.

$$\begin{aligned} 10x - 3y &= 3 \\ 10x - 3(-2x + 9) &= 3 \\ 10x + 6x - 27 &= 3 \\ 16x &= 30 \\ x &= \frac{15}{8} \end{aligned}$$

Substitute the value of x back into the second equation and solve for y.

$$2x + y = 9 \Rightarrow y = -2x + 9 \Rightarrow y = -2 \cdot \frac{15}{8} + 9 \Rightarrow y = \frac{21}{4}$$

Answer The solution to the system is $(\frac{15}{8}, \frac{21}{4})$. The system is consistent since it has only one solution.

Another method to determine if the system of equations is an inconsistent, consistent or dependent system is to solve them algebraically using the elimination or substitution method.

Example 6

Solve the following system of equations. Identify the system as consistent, inconsistent or dependent.

$$\begin{aligned} 3x - 2y &= 4 \\ 9x - 6y &= 1 \end{aligned}$$

Solution

Lets solve this system by the method of multiplication.

Multiply the first equation by 3.

$$\begin{array}{rcl}
 3(3x - 2y = 4) & & 9x - 6y = 12 \\
 & \Rightarrow & \\
 9x - 6y = 1 & & 9x - 6y = 1
 \end{array}$$

Add the two equations.

$$\begin{array}{r}
 9x - 6y = 12 \\
 9x - 6y = 1 \\
 \hline
 0 = 13 \quad \text{This Statement is not true}
 \end{array}$$

Answer If, by trying to obtain a solution to a system, we arrive at a statement that is not true, then the system is **inconsistent**.

Example 7

Solve the following system of equations. Identify the system as consistent, inconsistent or dependent.

$$\begin{array}{r}
 4x + y = 3 \\
 12x + 3y = 9
 \end{array}$$

Solution

Lets solve this system by substitution.

Solve the first equation for y.

$$4x + y = 3 \Rightarrow y = -4x + 3$$

Substitute this expression for y in the second equation.

$$\begin{array}{r}
 12x + 3y = 9 \\
 12x + 3(-4x + 3) = 9 \\
 12x - 12x + 9 = 9 \\
 9 = 9
 \end{array}$$

This is always a **true** statement.

Answer If, by trying to obtain a solution to a system, we arrive at a statement that is always true, then the system is dependent.

A second glance at the system in this example reveals that the second equation is three times the first equation so the two lines are identical. The system has an infinite number of solutions because they are really the same equation and trace out the same line.

Lets clarify this statement. An infinite number of solutions does not mean that any ordered pair (x, y) satisfies the system of equations. Only ordered pairs that solve the equation in the system are also solutions to the system.

For example, $(1, 2)$ is not a solution to the system because when we plug it into the equations it does not check out.

$$4x + y = 3$$

$$4(1) + 2 \neq 3$$

To find which ordered pair satisfies this system, we can pick any value for x and find the corresponding value for y .

$$\text{For } x = 1, 4(1) + y = 3 \Rightarrow y = -1$$

$$\text{For } x = 2, 4(2) + y = 3 \Rightarrow y = -3$$

Lets summarize our finding for determining the type of system algebraically.

- A **consistent system** will always give exactly one solution.
- An **inconsistent system** will always give a FALSE statement (for example $9 = 0$).
- A **dependent system** will always give a TRUE statement (such as $9 = 9$ or $0 = 0$).

Applications

In this section, we will look at a few application problems and see how consistent, inconsistent and dependent systems might arise in practice.

Example 8

A movie rental store CineStar offers customers two choices. Customers can pay a yearly membership of \$45 and then rent each movie for \$2 or they can choose not to pay the membership fee and rent each movie for \$3.50. How many movies would you have to rent before membership becomes the cheaper option?

Solution

Lets translate this problem into algebra. Since there are two different options to consider, we will write two different equations and form a system.

The choices are membership option and no membership option.

Our variables are


The number of movies you rent, lets call this x .

The total cost for renting movies, lets call this y .

TABLE 7.3:

	flat fee	rental fee	total
membership	\$45	$2x$	$y = 45 + 2x$
no membership	\$0	$3.50x$	$y = 3.5x$

The flat fee is the dollar amount you pay per year and the rental fee is the dollar amount you pay when you rent movies. For the membership option, the rental fee is $2x$ since you would pay \$2 for each movie you rented. For the no membership option, the rental fee is $3.50x$ since you would pay \$3.50 for each movie you rented.



Our system of equations is

$$y = 45 + 2x$$

$$y = 3.50x$$

The graph of our system of equations is shown to the right.

This system can be solved easily with the method of substitution since each equation is already solved for y . Substitute the second equation into the first one

$$y = 45 + 2x$$

$$\Rightarrow 3.50x = 45 + 2x \Rightarrow 1.50x = 45 \Rightarrow x = 30 \text{ movies}$$

$$y = 3.50x$$

Answer You would have to rent 30 movies per year before the membership becomes the better option.

This example shows a real situation where a consistent system of equations is useful in finding a solution. Remember that for a consistent system, the lines that make up the system intersect at single point. In other words, the lines are not parallel or the slopes are different.

In this case, the slopes of the lines represent the price of a rental per movie. The lines cross because the price of rental per movie is different for the two options in the problem

Lets examine a situation where the system is inconsistent. From the previous explanation, we can conclude that the lines will not intersect if the slopes are the same (but the y -intercept is different). Lets change the previous problem so that this is the case.

Example 9

Two movie rental stores are in competition. Movie House charges an annual membership of \$30 and charges \$3 per movie rental. Flicks for Cheap charges an annual membership of \$15 and charges \$3 per movie rental. After how many movie rentals would Movie House become the better option?

Solution

It should already be clear to see that Movie House will never become the better option, since its membership is more expensive and it charges the same amount per movie as Flicks for Cheap.

The lines that describe each option have different y -intercepts, namely 30 for Movie House and 15 for Flicks for Cheap. They have the same slope, three dollars per movie. This means that the lines are parallel and the system is inconsistent.

Lets see how this works algebraically:

Our variables are:

The number of movies you rent, lets call this x .

The total cost for renting movies, lets call this y .

TABLE 7.4:

	flat fee	rental fee	total
Movie House	\$30	$3x$	$y = 30 + 3x$
Flicks for Cheap	\$15	$3x$	$y = 15 + 3x$

The system of equations that describes this problem is

$$y = 30 + 3x$$

$$y = 15 + 3x$$

Lets solve this system by substituting the second equation into the first equation.

$$y = 30 + 3x$$

$$\Rightarrow 15 + 3x = 30 + 3x \Rightarrow 15 = 30$$

$$y = 15 + 3x$$

This statement is always false.

Answer This means that the system is inconsistent.

Example 10

Peter buys two apples and three bananas for \$4. Nadia buys four apples and six bananas for \$8 from the same store. How much does one banana and one apple costs?

Solution

We must write two equations, one for Peters purchase and one for Nadias purchase.

Lets define our variables as

a is the cost of one apple

b is the cost of one banana

TABLE 7.5:

	Cost of Apples	Cost of Bananas	Total Cost
Peter	$2a$	$3b$	$2a + 3b = 4$
Nadia	$4a$	$6b$	$4a + 6b = 8$

The system of equations that describes this problem is:

$$2a + 3b = 4$$

$$4a + 6b = 8$$

Lets solve this system by multiplying the first equation by -2 and adding the two equations.

$$\begin{array}{rcl} -2(2a + 3b = 4) & & -4a - 6b = -8 \\ 4a + 6b = 8 & \Rightarrow & 4a + 6b = 8 \\ \hline & & 0 + 0 = 0 \end{array}$$

This statement is always true. This means that the system is **dependent**.

Looking at the problem again, we see that we were given exactly the same information in both statements. If Peter buys two apples and three bananas for \$4 it makes sense that if Nadia buys twice as many apples (four apples) and twice as many bananas (six bananas) she will pay twice the price (\$8). Since the second equation does not give any new information, it is not possible to find out the price of each piece of fruit.

Answer The two equations describe the same line. This means that the system is dependent.

Review Questions

- Express each equation in slope-intercept form. Without graphing, state whether the system of equations is consistent, inconsistent or dependent.

(a) $3x - 4y = 13$

$y = -3x - 7$

(b) $\frac{3x}{5} + y = 3$

$1.2x + 2y = 6$

(c) $3x - 4y = 13$

$y = -3x - 7$

(d) $3x - 3y = 3$

$x - y = 1$

(e) $0.5x - y = 30$

$0.5x - y = -30$

(f) $4x - 2y = -2$

$3x + 2y = -12$

- Find the solution of each system of equations using the method of your choice. Please state whether the system is inconsistent or dependent.

(a) $3x + 2y = 4$

$-2x + 2y = 24$

(b) $5x - 2y = 3$

$2x - 3y = 10$

(c) $3x - 4y = 13$

$y = -3x - y$

(d) $5x - 4y = 1$

$-10x + 8y = -30$

- (e) $4x + 5y = 0$
 $3x = 6y + 4.5$
- (f) $-2y + 4x = 8$
 $y - 2x = -4$
- (g) $x - \frac{y}{2} = \frac{3}{2}$
 $3x + y = 6$
- (h) $0.05x + 0.25y = 6$
 $x + y = 24$
- (i) $x + \frac{2y}{3} = 6$
 $3x + 2y = 2$
- A movie house charges \$4.50 for children and \$8.00 for adults. On a certain day, 1200 people enter the movie house and \$8,375 is collected. How many children and how many adults attended?
 - Andrew placed two orders with an internet clothing store. The first order was for thirteen ties and four pairs of suspenders, and totaled \$487. The second order was for six ties and two pairs of suspenders, and totaled \$232. The bill does not list the per-item price but all ties have the same price and all suspenders have the same price. What is the cost of one tie and of one pair of suspenders?
 - An airplane took four hours to fly 2400 miles in the direction of the jet-stream. The return trip against the jet-stream took five hours. What were the airplanes speed in still air and the jet-stream's speed?
 - Nadia told Peter that she went to the farmers market and she bought two apples and one banana and that it cost her \$2.50. She thought that Peter might like some fruit so she went back to the seller and bought four more apples and two more bananas. Peter thanked Nadia, but told her that he did not like bananas, so will only pay her for four apples. Nadia told him that the second time she paid \$6.00 for fruit. Please help Peter figure out how much to pay Nadia paid for four apples.

Review Answers

- consistent
 - dependent
 - consistent
 - dependent
 - inconsistent
 - consistent
- $x = -4, y = 8$
 - $x = -1, y = -4$
 - $x = -1, y = -4$
 - inconsistent
 - $x = -2.5, y = 2$
 - dependent
 - $x = \frac{9}{5}, y = \frac{3}{5}$
 - $x = 0, y = 24$
 - dependent
- 350 children, 850 Adults
- Ties = \$23, suspenders = \$47
- Airplane speed = 540 mph, jet – stream speed = 60 mph
- This represents an inconsistent system. Someone is trying to overcharge! It is not possible to determine the price of apples alone.

7.6 Systems of Linear Inequalities

Learning Objectives

- Graph linear inequalities in two variables.
- Solve systems of linear inequalities.
- Solve optimization problems.

Introduction

In the last chapter, you learned how to graph a linear inequality in two variables. To do that you graphed the equation of the straight line on the coordinate plane. The line was solid for signs where the equal sign is included. The line was dashed for $<$ or $>$ where signs the equal sign is not included. Then you shaded above the line (if $y >$ or $y \geq$) or below the line (if $y <$ or $y \leq$).

In this section, we will learn how to graph two or more linear inequalities on the same coordinate plane. The inequalities are graphed separately on the same graph and the solution for the system is the common shaded region between all the inequalities in the system. One linear inequality in two variables divides the plane into two **half-planes**. A **system** of two or more linear inequalities can divide the plane into more complex shapes. Lets start by solving a system of two inequalities.

Graph a System of Two Linear Inequalities

Example 1

Solve the following system.

$$\begin{aligned} 2x + 3y &\leq 18 \\ x - 4y &\leq 12 \end{aligned}$$

Solution

Solving systems of linear inequalities means graphing and finding the intersections. So we graph each inequality, and then find the intersection regions of the solution.


Lets rewrite each equation in slope-intercept form. This form is useful for graphing but also in deciding which region of the coordinate plane to shade. Our system becomes

$$\begin{aligned} 3y &\leq -2x + 18 & y &\leq \frac{-2}{3}x + 6 \\ -4y &\leq -x + 12 & y &\geq \frac{x}{4} - 3 \end{aligned} \Rightarrow$$


Notice that the inequality sign in the second equation changed because we divided by a negative number.

For this first example, we will graph each inequality separately and then combine the results.


We graph the equation of the line in the first inequality and draw the following graph.




The line is solid because the equal sign is included in the inequality. Since the inequality is less than or equal to, we shade below the line.




We graph the second equation in the inequality and obtain the following graph.



The line is solid again because the equal sign is included in the inequality. We now shade above because y is **greater** than or equal.



When we combine the graphs, we see that the blue and red shaded regions overlap. This overlap is where both inequalities work. Thus the purple region denotes the solution of the system.



The kind of solution displayed in this example is called **unbounded**, because it continues forever in at least one direction (in this case, forever upward and to the left).


Example 2

There are also situations where a system of inequalities has no solution. For example, let's solve this system.


$$y \leq 2x - 4$$

$$y > 2x + 6$$

Solution: We start by graphing the first line. The line will be solid because the equal sign is included in the inequality. We must shade downwards because y is less than.



Next we graph the second line on the same coordinate axis. This line will be dashed because the equal sign is not included in the inequality. We must shade upward because y is greater than.




This graph shows no overlapping between the two shaded regions. We know that the lines will never intersect because they are parallel. The slope equals two for both lines. The regions will never overlap even if we extend the lines further.

This is an example of a system of inequalities with no solution.

For a system of inequalities, we can still obtain a solution even if the lines are parallel. Let's change the system of inequalities in Example 2 so the inequality signs for the two expressions are reversed.

$$y \geq 2x - 4$$

$$y < 2x + 6$$



The procedure for solving this system is almost identical to the previous one, except we shade upward for the first inequality and we shade downward for the second inequality. Here is the result.

In this case, the shaded regions do overlap and the system of inequalities has the solution denoted by the purple region.

Graph a System of More Than Two Linear Inequalities

In the previous section, we saw how to find the solution to a system of two linear inequalities. The solutions for these kinds of systems are always unbounded. In other words, the region where the shadings overlap continues infinitely in at least one direction. We can obtain **bounded** solutions by solving systems that contain more than two inequalities. In such cases the solution region will be bounded on four sides.

Lets examine such a solution by solving the following example.

Example 3

Find the solution to the following system of inequalities.


$$\begin{aligned} 3x - y &< 4 \\ 4y + 9x &< 8 \\ x &\geq 0 \\ y &\geq 0 \end{aligned}$$

Solution


Lets start by writing our equation in slope-intercept form.

$$\begin{aligned} y &> 3x - 4 \\ y &< -\frac{9}{4}x + 2 \\ x &\geq 0 \\ y &\geq 0 \end{aligned}$$


Now we can graph each line and shade appropriately. First we graph $y > 3x - 4$.



Next we graph $y < -\frac{9}{4}x + 2$



Finally we graph $x \geq 0$ and $y \geq 0$, and the intersecting region is shown in the following figure.



The solution is **bounded** because there are lines on all sides of the solution region. In other words the solution region is a bounded geometric figure, in this case a triangle.

Write a System of Linear Inequalities

There are many interesting application problems that involve the use of system of linear inequalities. However, before we fully solve application problems, let's see how we can translate some simple word problems into algebraic equations.

For example, you go to your favorite restaurant and you want to be served by your best friend who happens to work there. However, your friend works in a certain region of the restaurant. The restaurant is also known for its great views but you have to sit in a certain area of the restaurant that offers these views. Solving a system of linear inequalities will allow you to find the area in the restaurant where you can sit to get the best views and be served by your friend.

Typically, systems of linear inequalities deal with problems where you are trying to find the best possible situation given a set of constraints.

Example 4

Write a system of linear inequalities that represents the following conditions.


The sum of twice a number x and three times another number y is greater than 6, and y is less than three times x .

Solution

Let's take each statement in turn and write it algebraically:


1. The sum of twice a number x and three times another number y is greater than 6.

This can be written as



2. y is less than three times x .

This can be written as



The system of inequalities arising from these statements is

$$\begin{aligned} 2x + 3y &> 6 \\ y &< 3x \end{aligned}$$

This system of inequalities can be solved using the method outlined earlier in this section. We will not solve this system because we want to concentrate on learning how to write a system of inequalities from a word problem.

Solve Real-World Problems Using Systems of Linear Inequalities

As we mentioned before, there are many interesting application problems that require the use of systems of linear inequalities. Most of these application problems fall in a category called **linear programming** problems.

Linear programming is the process of taking various linear inequalities relating to some situation, and finding the best possible value under those conditions. A typical example would be taking the limitations of materials and labor, then determining the best production levels for maximal profits under those conditions. These kinds of problems are used every day in the organization and allocation of resources. These real life systems can have dozens or hundreds of variables, or more. In this section, we will only work with the simple two-variable linear case.

The general process is to:

- Graph the inequalities (called **constraints**) to form a bounded area on the x,y -plane (called **the feasibility region**).
- Figure out the coordinates of the corners (or vertices) of this feasibility region by solving the systems of equations that give the solutions to each of the intersection points.
- Test these corner points in the formula (called the **optimization equation**) for which you're trying to find the **maximum** or **minimum** value.


Example 5

Find the maximum and minimum value of $z = 2x + 5y$ given the constraints.

$$\begin{aligned} 2x - y &\leq 12 \\ 4x + 3y &\geq 0 \\ x - y &\leq 6 \end{aligned}$$

Solution

Step 1: Find the solution to this system of linear inequalities by graphing and shading appropriately. To graph we must rewrite the equations in slope-intercept form.



$$\begin{aligned} y &\geq 2x - 12 \\ y &\geq -\frac{4}{3}x \\ y &\geq x - 6 \end{aligned}$$

These three linear inequalities are called the **constraints**.

The solution is the shaded region in the graph. This is called the **feasibility region**. That means all possible solutions occur in that region. However in order to find the optimal solution we must go to the next steps.

Step 2

Next, we want to find the corner points. In order to find them exactly, we must form three systems of linear equations and solve them algebraically.

System 1:

$$y = 2x - 12$$

$$y = -\frac{4}{3}x$$

Substitute the first equation into the second equation:

$$-\frac{4}{3}x = 2x - 12 \Rightarrow -4x = 6x - 36 \Rightarrow -10x = -36 \Rightarrow x = 3.6$$

$$y = 2x - 12 \Rightarrow y = 2(3.6) - 12 \Rightarrow y = -4.8$$

The intersection point of lines is $(3.6, -4.8)$

System 2:

$$y = 2x - 12$$

$$y = x - 6$$

Substitute the first equation into the second equation.

$$x - 6 = 2x - 12 \Rightarrow 6 = x \Rightarrow x = 6$$

$$y = x - 6 \Rightarrow y = 6 - 6 \Rightarrow y = 0$$

The intersection point of lines is $(6, 0)$.

System 3:

$$y = -\frac{4}{3}x$$

$$y = x - 6$$

Substitute the first equation into the second equation.

$$x - 6 = -\frac{4}{3}x \Rightarrow 3x - 18 = -4x \Rightarrow 7x = 18 \Rightarrow x = 2.57$$

$$y = x - 6 \Rightarrow y = 2.57 - 6 \Rightarrow y = -3.43$$

The intersection point of lines is $(2.57, -3.43)$.

So the corner points are $(3.6, -4.8)$, $(6, 0)$ and $(2.57, -3.43)$.

Step 3

Somebody really smart proved that, for linear systems like this, the maximum and minimum values of the optimization equation will always be on the corners of the feasibility region. So, to find the solution to this exercise, we need to plug these three points into $z = 2x + 5y$.

$(3.6, -4.8)$	$z = 2(3.6) + 5(-4.8) = -16.8$
$(6, 0)$	$z = 2(6) + 5(0) = 12$
$(2.57, -3.43)$	$z = 2(2.57) + 5(-3.43) = -12.01$

The highest value of 12 occurs at point $(6, 0)$ and the lowest value of -16.8 occurs at $(3.6, -4.8)$.

In the previous example, we learned how to apply the method of linear programming out of context of an application problem. In the next example, we will look at a real-life application.

Example 6

You have \$10,000 to invest, and three different funds from which to choose. The municipal bond fund has a 5% return, the local bank's CDs have a 7% return, and a high-risk account has an expected 10% return. To minimize risk, you decide not to invest any more than \$1,000 in the high-risk account. For tax reasons, you need to invest at least three times as much in the municipal bonds as in the bank CDs. Assuming the year-end yields are as expected, what are the optimal investment amounts?

Solution

Lets define some *variables*.

x is the amount of money invested in the municipal bond at 5% return.

y is the amount of money invested in the banks CD at 7% return.

$10000 - x - y$ is the amount of money invested in the high-risk account at 10% return.

z is the total interest returned from all the investments or $z = .05x + .07y + .1(10000 - x - y)$ or $z = 1000 - 0.05x - 0.03y$. This is the amount that we are trying to maximize. Our goal is to find the values of x and y that maximizes the value of z .

Now, lets write inequalities for the *constraints*.

You decide not to invest more than \$1000 in the high-risk account.

$$10000 - x - y \leq 1000$$

You need to invest at least three times as much in the municipal bonds as in the bank CDs.

$$3y \leq x$$

Also we write expressions for the fact that we invest more than zero dollars in each account.

$$\begin{aligned}x &\geq 0 \\y &\geq 0 \\10000 - x - y &\geq 0\end{aligned}$$

To summarize, we must maximize the expression $z = 1000 - .05x - .03y$.

Using the constraints,

$$10000 - x - y \leq 1000$$

$$3y \leq x$$

$$x \geq 0$$

$$y \geq 0$$

$$10000 - x - y \geq 0$$

$$y \geq 9000 - x$$

$$y \leq \frac{x}{3}$$

$$x \geq 0$$


$$y \geq 0$$

$$y \leq 10000 - x$$

Let's rewrite each in slope-intercept form.

Step 1 Find the solution region to the set of inequalities by graphing each line and shading appropriately.

The following figure shows the overlapping region.



The purple region is the feasibility region where all the possible solutions can occur.

Step 2 Next, we need to find the corner points of the shaded solution region. Notice that there are four intersection points. To find them we must pair up the relevant equations and solve the resulting system.

System 1:

$$\begin{aligned}y &= -\frac{x}{3} \\y &= 10000 - x\end{aligned}$$

Substitute the first equation into the second equation.

$$\begin{aligned}\frac{x}{3} &= 10000 - x \Rightarrow x = 30000 - 3x \Rightarrow x = 7500 \\ y &= \frac{x}{3} \Rightarrow y = \frac{7500}{3} \Rightarrow y = 2500\end{aligned}$$

The intersection point is (7500, 2500).

System 2:

$$\begin{aligned}y &= -\frac{x}{3} \\ y &= 9000 - x\end{aligned}$$

Substitute the first equation into the second equation.

$$\frac{x}{3} = 9000 - x \Rightarrow x = 27000 - 3x \Rightarrow 4x = 27000 \Rightarrow x = 6750$$

$$\frac{x}{3} \Rightarrow y = \frac{6750}{3} \Rightarrow y = 2250$$

The intersection point is (6750, 2250).

System 3:

$$\begin{aligned}y &= 0 \\ y &= 10000 - x\end{aligned}$$

The intersection point is (10000, 0).

System 4:

$$\begin{aligned}y &= 0 \\ y &= 9000 - x\end{aligned}$$

The intersection point is (9000, 0).

Step 3: In order to find the maximum value for z , we need to plug all intersection points into z and take the largest number.

(7500, 2500)	$z = 1000 - 0.05(7500) - 0.03(2500) = 550$
(6750, 2250)	$z = 1000 - 0.05(6750) - 0.03(2250) = 595$
(10000, 0)	$z = 1000 - 0.05(10000) - 0.03(0) = 500$
(9000, 0)	$z = 1000 - 0.05(9000) - 0.03(0) = 550$

Answer

The maximum return on the investment of \$595 occurs at point (6750, 2250). This means that

\$6,750 is invested in the municipal bonds.

\$2,250 is invested in the bank CDs.

\$1,000 is invested in the high-risk account.

Review Questions






Find the solution region of the following systems of inequalities


1. $x - y < -6$
 $2y \geq 3x + 17$
2. $4y - 5x < 8$
 $-5x \geq 16 - 8y$
3. $5x - y \geq 5$
 $2y - x \geq -10$
4. $5x + 2y \geq -25$
 $3x - 2y \leq 17$
 $x - 6y \geq 27$
5. $2x - 3y \leq 21$
 $x + 4y \leq 6$
 $3x + y \geq -4$
6. $12x - 7y < 120$
 $7x - 8y \geq 36$
 $5x + y \geq 12$

Solve the following linear programming problems:

7. Given the following constraints find the maximum and minimum values of $z = -x + 5y$
 $x + 3y \leq 0$
 $x - y \geq 0$
 $3x - 7y \leq 16$
8. In Andrews Furniture Shop, he assembles both bookcases and TV cabinets. Each type of furniture takes him about the same time to assemble. He figures he has time to make at most 18 pieces of furniture by this Saturday.
The materials for each bookcase cost him \$20 and the materials for each TV stand costs him \$45 . He has \$600 to spend on materials. Andrew makes a profit of \$60 on each bookcase and a profit of \$100 for each TV stand.
Find how many of each piece of furniture Andrew should make so that he maximizes his profit.

Review Answers





7. Maximum of $z = 0$ at point $(0, 0)$, minimum of $z = -16$ at point $(-4, -4)$
8. Maximum profit of \$1,440 by making 9 bookcases and 9 TV stands.

CHAPTER **8** Exponents and Polynomials

Chapter Outline

- 8.1 EXPONENT PROPERTIES INVOLVING PRODUCTS**
 - 8.2 EXPONENT PROPERTIES INVOLVING QUOTIENTS**
 - 8.3 ZERO, NEGATIVE, AND FRACTIONAL EXPONENTS**
 - 8.4 SCIENTIFIC NOTATION**
 - 8.5 ADDITION AND SUBTRACTION OF POLYNOMIALS**
 - 8.6 MULTIPLICATION OF POLYNOMIALS**
 - 8.7 SPECIAL PRODUCTS OF POLYNOMIALS**
 - 8.8 DIVISION OF POLYNOMIALS**
-

8.1 Exponent Properties Involving Products


Learning Objectives

- Use the product of a power property.
- Use the power of a product property.
- Simplify expressions involving product properties of exponents.


Introduction

In this chapter, we will discuss exponents and exponential functions. In Lessons 8.1, 8.2 and 8.3, we will be learning about the rules governing exponents. We will start with what the word exponent means.

Consider the area of the square shown right. We know that the area is given by:



But we also know that for any rectangle, Area = (width) \cdot (height), so we can see that:



Similarly, the volume of the cube is given by:

$$\text{Volume} = \text{width} \cdot \text{depth} \cdot \text{height} = x \cdot x \cdot x$$

But we also know that the volume of the cube is given by Volume = x^3 so clearly

$$x^3 = x \cdot x \cdot x$$

You probably know that the **power** (the small number to the top right of the x) tells you how many x 's to multiply together. In these examples the x is called the **base** and the **power** (or **exponent**) tells us how many **factors** of the **base** there are in the full expression.

$$x^2 = \underbrace{x \cdot x}_{2 \text{ factors of } x}$$

$$x^3 = \underbrace{x \cdot x \cdot x}_{3 \text{ factors of } x}$$

$$x^7 = \underbrace{x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot x}_{7 \text{ factors of } x}$$

$$x^n = \underbrace{x \cdot x \cdot \dots \cdot x}_{n \text{ factors of } x}$$

Example 1

Write in exponential form.

(a) $2 \cdot 2$

(b) $(-3)(-3)(-3)$

(c) $y \cdot y \cdot y \cdot y \cdot y$

(d) $(3a)(3a)(3a)(3a)$

Solution

(a) $2 \cdot 2 = 2^2$ because we have 2 factors of 2

(b) $(-3)(-3)(-3) = (-3)^3$ because we have 3 factors of (-3)

(c) $y \cdot y \cdot y \cdot y \cdot y = y^5$ because we have 5 factors of y

(d) $(3a)(3a)(3a)(3a) = (3a)^4$ because we have 4 factors of $3a$

When we deal with numbers, we usually just simplify. We'd rather deal with 16 than with 2^4 . However, with variables, we need the exponents, because we'd rather deal with x^7 than with $x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot x$.

Lets simplify Example 1 by evaluating the numbers.

Example 2

Simplify.

(a) $2 \cdot 2$

(b) $(-3)(-3)(-3)$

(c) $y \cdot y \cdot y \cdot y \cdot y$

(d) $(3a)(3a)(3a)(3a)$

Solution

(a) $2 \cdot 2 = 2^2 = 4$

(b) $(-3)(-3)(-3) = (-3)^3 = -27$

(c) $y \cdot y \cdot y \cdot y \cdot y = y^5$

(d) $(3a)(3a)(3a)(3a) = (3a)^4 = 3^4 \cdot a^4 = 81a^4$

Note: You must be careful when taking powers of negative numbers. Remember these rules.

(negative number) (positive number) = negative number

(negative number) (negative number) = positive number

For **even powers of negative numbers**, the answer is always positive. Since we have an even number of factors, we make pairs of negative numbers and all the negatives cancel out.

$$(-2)^6 = (-2)(-2)(-2)(-2)(-2)(-2) = \underbrace{(-2)(-2)}_{+4} \cdot \underbrace{(-2)(-2)}_{+4} \cdot \underbrace{(-2)(-2)}_{+4} = +64$$

For **odd powers of negative numbers**, the answer is always negative. Since we have an odd number of factors, we can make pairs of negative numbers to get positive numbers but there is always an unpaired negative factor, so the answer is negative:

$$\text{Ex: } (-2)^5 = (-2)(-2)(-2)(-2)(-2) = \underbrace{(-2)(-2)}_{+4} \cdot \underbrace{(-2)(-2)}_{+4} \cdot \underbrace{(-2)}_{-2} = -32$$

Use the Product of Powers Property

What happens when we multiply one power of x by another? See what happens when we multiply x *to the power 5* by x *cubed*. To illustrate better we will use the full factored form for each:

$$\underbrace{(x \cdot x \cdot x \cdot x \cdot x)}_{x^5} \cdot \underbrace{(x \cdot x \cdot x)}_{x^3} = \underbrace{(x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot x)}_{x^8}$$

So $x^5 \cdot x^3 = x^8$. You may already see the pattern to multiplying powers, but let's confirm it with another example. We will multiply x *squared* by x *to the power 4*:

$$\underbrace{(x \cdot x)}_{x^2} \cdot \underbrace{(x \cdot x \cdot x \cdot x)}_{x^4} = \underbrace{(x \cdot x \cdot x \cdot x \cdot x \cdot x)}_{x^6}$$

So $x^2 \cdot x^4 = x^6$. Look carefully at the powers and how many factors there are in each calculation. 5 factors of x times 3 factors of x equals $(5 + 3) = 8$ factors of x . 2 factors of x times 4 factors of x equals $(2 + 4) = 6$ factors of x .

You should see that when we take the product of two powers of x , the number of factors of x in the answer is the sum of factors in the terms you are multiplying. In other words the exponent of x in the answer is the sum of the exponents in the product.

Product rule for exponents: $x^n \cdot x^m = x^{n+m}$

Example 3

Multiply $x^4 \cdot x^5$.

Solution

$$x^4 \cdot x^5 = x^{4+5} = x^9$$

When multiplying exponents of the same base, it is a simple case of adding the exponents. It is important that when you use the product rule you avoid easy-to-make mistakes. Consider the following.

Example 4

Multiply $2^2 \cdot 2^3$.

Solution

$$2^2 \cdot 2^3 = 2^5 = 32$$

Note that when you use the product rule you **DO NOT MULTIPLY BASES**. In other words, you must avoid the common error of writing ~~$2^2 \cdot 2^3 = 4^5$~~ . Try it with your calculator and check which is right!

Example 5

Multiply $2^2 \cdot 3^3$.

Solution

$$2^2 \cdot 3^3 = 4 \cdot 27 = 108$$

In this case, the bases are different. The product rule for powers **ONLY APPLIES TO TERMS THAT HAVE THE SAME BASE**. Common mistakes with problems like this include ~~$2^2 \cdot 3^3 = 6^5$~~ .

Use the Power of a Product Property

We will now look at what happens when we raise a whole expression to a power. Lets take *to the power 4* and *cube it*. Again we will use the full factored form for each.

$$(x^4)^3 = x^4 \cdot x^4 \cdot x^4 \qquad 3 \text{ factors of } x \text{ to the power 4.}$$

$$\underbrace{(x \cdot x \cdot x \cdot x)}_{x^4} \cdot \underbrace{(x \cdot x \cdot x \cdot x)}_{x^4} \cdot \underbrace{(x \cdot x \cdot x \cdot x)}_{x^4} = \underbrace{(x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot x)}_{x^{12}}$$

So $(x^4)^3 = x^{12}$. It is clear that when we raise a power of x to a new power, the powers multiply.

When we take an expression and raise it to a power, we are multiplying the existing powers of x by the power above the parenthesis.

Power rule for exponents: $(x^n)^m = x^{n \cdot m}$

Power of a product

If we have a product inside the parenthesis and a power on the parenthesis, then the power goes on each element inside. So that, for example, $(x^2y)^4 = (x^2)^4 \cdot (y)^4 = x^8y^4$. Watch how it works the long way.

$$\underbrace{(x \cdot x \cdot y)}_{x^2y} \cdot \underbrace{(x \cdot x \cdot y)}_{x^2y} \cdot \underbrace{(x \cdot x \cdot y)}_{x^2y} \cdot \underbrace{(x \cdot x \cdot y)}_{x^2y} = \underbrace{(x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot x \cdot y \cdot y \cdot y \cdot y)}_{x^8y^4}$$

Power rule for exponents: $(x^n)^m = x^{nm}$ and $(x^n y^m)^p = x^{np} y^{mp}$

WATCH OUT! This does NOT work if you have a sum or difference inside the parenthesis. For example, $(x + y)^2 \neq x^2 + y^2$. This is a commonly made mistake. It is easily avoidable if you remember what an exponent means $(x + y)^2 = (x + y)(x + y)$. We will learn how to simplify this expression in a later chapter.

Lets apply the rules we just learned to a few examples.

When we have numbers, we just evaluate and most of the time it is not really important to use the product rule and the power rule.

Example 6

Simplify the following expressions.

(a) $3^4 \cdot 3^7$

(b) $2^6 \cdot 2$

(c) $(4^2)^3$

Solution

In each of the examples, we want to evaluate the numbers.

(a) Use the product rule first: $3^5 \cdot 3^7 = 3^{12}$

Then evaluate the result: $3^{12} = 531,441$

OR

We can evaluate each part separately and then multiply them. $3^5 \cdot 3^7 = 243 \cdot 2,187 = 531,441$.

Use the product rule first. $2^6 \cdot 2 = 2^7$

Then evaluate the result. $2^7 = 128$

OR

We can evaluate each part separately and then multiply them. $2^6 \cdot 2 = 64 \cdot 2 = 128$

(c) Use the power rule first. $(4^2)^3 = 4^6$

Then evaluate the result. $4^6 = 4096$

OR

We evaluate inside the parenthesis first. $(4^2)^3 = (16)^3$

Then apply the power outside the parenthesis. $(16)^3 = 4096$

When we have just one variable in the expression then we just apply the rules.

Example 7

Simplify the following expressions.

(a) $x^2 \cdot x^7$

(b) $(y^3)^5$

Solution

(a) Use the product rule. $x^2 \cdot x^7 = x^{2+7} = x^9$

(b) Use the power rule. $(y^3)^5 = y^{3 \cdot 5} = y^{15}$

When we have a mix of numbers and variables, we apply the rules to the numbers or to each variable separately.

Example 8

Simplify the following expressions.

(a) $(3x^2y^3) \cdot (4xy^2)$

(b) $(4xyz) \cdot (x^2y^3) \cdot (2yz^4)$

(c) $(2a^3b^3)^2$

Solution

(a) We group like terms together.

$$(3x^2y^3) \cdot (4xy^2) = (3 \cdot 4) \cdot (x^2 \cdot x) \cdot (y^3 \cdot y^2)$$

We multiply the numbers and apply the product rule on each grouping.

$$12x^3y^5$$

(b) We group like terms together.

$$(4xyz) \cdot (x^2y^3) \cdot (2yz^4) = (4 \cdot 2) \cdot (x \cdot x^2) \cdot (y \cdot y^3 \cdot y) \cdot (z \cdot z^4)$$

We multiply the numbers and apply the product rule on each grouping.

$$8x^3y^5z^5$$

(c) We apply the power rule for each separate term in the parenthesis.

$$(2a^3b^3)^2 = 2^2 \cdot (a^3)^2 \cdot (b^3)^2$$

We evaluate the numbers and apply the power rule for each term.

$$4a^6b^6$$

In problems that we need to apply the product and power rules together, we must keep in mind the order of operation. Exponent operations take precedence over multiplication.

Example 9

Simplify the following expressions.

(a) $(x^2)^2 \cdot x^3$

(b) $(2x^2y) \cdot (3xy^2)^3$

(c) $(4a^2b^3)^2 \cdot (2ab^4)^3$

Solution

(a) $(x^2)^2 \cdot x^3$

We apply the power rule first on the first parenthesis.

$$(x^2)^2 \cdot x^3 = x^4 \cdot x^3$$

Then apply the product rule to combine the two terms.

$$x^4 \cdot x^3 = x^7$$

(b) $(2x^2y) \cdot (3xy^2)^3$

We must apply the power rule on the second parenthesis first.

$$(2x^2y) \cdot (3xy^2)^3 = (2x^2y) \cdot (27x^3y^6)$$

Then we can apply the product rule to combine the two parentheses.

$$(2x^2y) \cdot (27x^3y^6) = 54x^5y^7$$

(c) $(4a^2b^3)^2 \cdot (2ab^4)^3$

We apply the power rule on each of the parentheses separately.

$$(4a^2b^3)^2 \cdot (2ab^4)^3 = (16a^4b^6) \cdot (8a^3b^{12})$$

Then we can apply the product rule to combine the two parentheses.

$$(16a^4b^6) \cdot (8a^3b^{12}) = 128a^7b^{18}$$

Review Questions

Write in exponential notation.

1. $4 \cdot 4 \cdot 4 \cdot 4 \cdot 4$
2. $3x \cdot 3x \cdot 3x$
3. $(-2a)(-2a)(-2a)(-2a)$
4. $6 \cdot 6 \cdot 6 \cdot x \cdot x \cdot y \cdot y \cdot y \cdot y$

Find each number:

5. 5^4
6. $(-2)^6$
7. $(0.1)^5$
8. $(-0.6)^3$

Multiply and simplify.

9. $6^3 \cdot 6^6$
10. $2^2 \cdot 2^4 \cdot 2^6$
11. $3^2 \cdot 4^3$
12. $x^2 \cdot x^4$
13. $(-2y^4)(-3y)$
14. $(4a^2)(-3a)(-5a^4)$

Simplify.

15. $(a^3)^4$
16. $(xy)^2$
17. $(3a^2b^3)^4$
18. $(-2xy^4z^2)^5$
19. $(-8x)^3(5x)^2$
20. $(4a^2)(-2a^3)^4$
21. $(12xy)(12xy)^2$
22. $(2xy^2)(-x^2y)^2(3x^2y^2)$

Review Answers

1. 4^5
2. $(3x)^3$

3. $(-2a)^4$
4. $6^3x^2y^4$
5. 625
6. 64
7. 0.00001
8. -0.216
9. 10077696
10. 4096
11. 576
12. x^6
13. $6y^5$
14. $60a^7$
15. a^{12}
16. x^2y^2
17. $81a^8b^{12}$
18. $-32x^5y^{20}z^{10}$
19. $12800x^5$
20. $64a^{14}$
21. $1728x^3y^3$
22. $6x^7y^6$

8.2 Exponent Properties Involving Quotients

Learning Objectives

- Use the quotient of powers property.
- Use the power of a quotient property.
- Simplify expressions involving quotient properties of exponents.

Use the Quotient of Powers Property

You saw in the last section that we can use exponent rules to simplify products of numbers and variables. In this section, you will learn that there are similar rules you can use to simplify quotients. Lets take an example of a quotient, x^7 divided by x^4 .

$$\frac{x^7}{x^4} = \frac{\cancel{x} \cdot \cancel{x} \cdot \cancel{x} \cdot \cancel{x} \cdot x \cdot x \cdot x}{\cancel{x} \cdot \cancel{x} \cdot \cancel{x} \cdot \cancel{x}} = \frac{x \cdot x \cdot x}{1} = x^3$$

You should see that when we divide two powers of x , the number of factors of x in the solution is the difference between the factors in the numerator of the fraction, and the factors in the denominator. In other words, when dividing expressions with the same base, keep the base and subtract the exponent in the denominator from the exponent in the numerator.

Quotient Rule for Exponents: $\frac{x^n}{x^m} = x^{n-m}$

When we have problems with different bases, we apply the quotient rule separately for each base.

$$\frac{x^5y^3}{x^3y^2} = \frac{\cancel{x} \cdot \cancel{x} \cdot \cancel{x} \cdot x \cdot x}{\cancel{x} \cdot \cancel{x} \cdot \cancel{x}} \cdot \frac{y \cdot y \cdot y}{y \cdot y} = \frac{x \cdot x}{1} \cdot \frac{y}{1} = x^2y \quad \text{OR} \quad \frac{x^5y^3}{x^3y^2} = x^{5-3} \cdot y^{3-2} = x^2y$$

Example 1

Simplify each of the following expressions using the quotient rule.

(a) $\frac{x^{10}}{x^5}$

(b) $\frac{a^6}{a}$

(c) $\frac{a^5b^4}{a^3b^2}$

Solution

Apply the quotient rule.

(a) $\frac{x^{10}}{x^5} = x^{10-5} = x^5$

(b) $\frac{a^6}{a} = a^{6-1} = a^5$

(c) $\frac{a^5b^4}{a^3b^2} = a^{5-3} \cdot b^{4-2} = a^2b^2$

Now lets see what happens if the exponent on the denominator is bigger than the exponent in the numerator.

Example 2Divide $x^4 \div x^7$

Apply the quotient rule.

$$\frac{x^4}{x^7} = x^{4-7} = x^{-3}$$

A negative exponent!?! What does that mean?

Lets do the division longhand by writing each term in factored form.

$$\frac{x^4}{x^6} = \frac{\cancel{x} \cdot \cancel{x} \cdot \cancel{x} \cdot \cancel{x}}{\cancel{x} \cdot \cancel{x} \cdot \cancel{x} \cdot \cancel{x} \cdot x \cdot x} = \frac{1}{x \cdot x} = \frac{1}{x^2}$$

We see that when the exponent in the denominator is bigger than the exponent in the numerator, we still subtract the powers. This time we subtract the smaller power from the bigger power and we leave the x 's in the denominator.

When you simplify quotients, to get answers with positive exponents you subtract the smaller exponent from the bigger exponent and leave the variable where the bigger power was.

- We also discovered what a negative power means $x^{-3} = \frac{1}{x^3}$. We'll learn more on this in the next section!

Example 3

Simplify the following expressions, leaving all powers positive.

(a) $\frac{x^2}{x^6}$

(b) $\frac{a^2b^6}{a^5b}$

Solution

(a) Subtract the exponent in the numerator from the exponent in the denominator and leave the x s in the denominator.

$$\frac{x^2}{x^6} = \frac{1}{x^{6-2}} = \frac{1}{x^4}$$

(b) Apply the rule on each variable separately.

$$\frac{a^2b^6}{a^5b} = \frac{1}{a^{5-2}} \cdot \frac{b^{6-1}}{1} = \frac{b^5}{a^3}$$

The Power of a Quotient Property

When we apply a power to a quotient, we can learn another special rule. Here is an example.

$$\left(\frac{x^3}{y^2}\right)^4 = \left(\frac{x^3}{y^2}\right) \cdot \left(\frac{x^3}{y^2}\right) \cdot \left(\frac{x^3}{y^2}\right) \cdot \left(\frac{x^3}{y^2}\right) = \frac{(x \cdot x \cdot x) \cdot (x \cdot x \cdot x) \cdot (x \cdot x \cdot x) \cdot (x \cdot x \cdot x)}{(y \cdot y) \cdot (y \cdot y) \cdot (y \cdot y) \cdot (y \cdot y)} = \frac{x^{12}}{y^8}$$

Notice that the power on the outside of the parenthesis multiplies with the power of the x in the numerator and the power of the y in the denominator. This is called the power of a quotient rule.

Power Rule for Quotients $\left(\frac{x^n}{y^m}\right)^p = \frac{x^{n \cdot p}}{y^{m \cdot p}}$

Simplifying Expressions Involving Quotient Properties of Exponents

Lets apply the rules we just learned to a few examples.

- When we have numbers with exponents and not variables with exponents, we evaluate.

Example 4

Simplify the following expressions.

(a) $\frac{4^5}{4^2}$

(b) $\frac{5^3}{5^7}$

(c) $\left(\frac{3^4}{5^2}\right)^2$

Solution

In each of the examples, we want to evaluate the numbers.

(a) Use the quotient rule first.

$$\frac{4^5}{4^2} = 4^{5-2} = 4^3$$

Then evaluate the result.

$$4^3 = 64$$

OR

We can evaluate each part separately and then divide.

$$\frac{1024}{16} = 64$$

(b) Use the quotient rule first.

$$\frac{5^3}{5^7} = \frac{1}{5^{7-3}} = \frac{1}{5^4}$$

Then evaluate the result.

$$\frac{1}{5^4} = \frac{1}{625}$$

OR

We can evaluate each part separately and then reduce.

$$\frac{5^3}{5^7} = \frac{125}{78125} = \frac{1}{625}$$

It makes more sense to apply the quotient rule first for examples (a) and (b). In this way the numbers we are evaluating are smaller because they are simplified first before applying the power.

(c) Use the power rule for quotients first.

$$\left(\frac{3^4}{5^2}\right)^2 = \frac{3^8}{5^4}$$

Then evaluate the result.

$$\frac{3^8}{5^4} = \frac{6561}{625}$$

OR

We evaluate inside the parenthesis first.

$$\left(\frac{3^4}{5^2}\right)^2 = \left(\frac{81}{25}\right)^2$$

Then apply the power outside the parenthesis.

$$\left(\frac{81}{25}\right)^2 = \frac{6561}{625}$$

When we have just one variable in the expression, then we apply the rules straightforwardly.

Example 5: Simplify the following expressions:

(a) $\frac{x^{12}}{x^5}$

(b) $\left(\frac{x^4}{x}\right)^5$

Solution:

(a) Use the quotient rule.

$$\frac{x^{12}}{x^5} = x^{12-5} = x^7$$

(b) Use the power rule for quotients first.

$$\left(\frac{x^4}{x}\right)^5 = \frac{x^{20}}{x^5}$$

Then apply the quotient rule

$$\frac{x^{20}}{x^5} = x^{15}$$

OR

Use the quotient rule inside the parenthesis first.

$$\left(\frac{x^4}{x}\right)^5 = (x^3)^5$$

Then apply the power rule.

$$(x^3)^5 = x^{15}$$

When we have a mix of numbers and variables, we apply the rules to each number or each variable separately.

Example 6

Simplify the following expressions.

(a) $\frac{6x^2y^3}{2xy^2}$

(b) $\left(\frac{2a^3b^3}{8a^7b}\right)^2$

Solution

(a) We group like terms together.

$$\frac{6x^2y^3}{2xy^2} = \frac{6}{2} \cdot \frac{x^2}{x} \cdot \frac{y^3}{y^2}$$

We reduce the numbers and apply the quotient rule on each grouping.

$$3xy >$$

(b) We apply the quotient rule inside the parenthesis first.

$$\left(\frac{2a^3b^3}{8a^7b}\right)^2 = \left(\frac{b^2}{4a^4}\right)^2$$

Apply the power rule for quotients.

$$\left(\frac{b^2}{4a^4}\right)^2 = \frac{b^4}{16a^8}$$

In problems that we need to apply several rules together, we must keep in mind the order of operations.

Example 7

Simplify the following expressions.

(a) $(x^2)^2 \cdot \frac{x^6}{x^4}$

$$(b) \left(\frac{16a^2}{4b^5} \right)^3 \cdot \frac{b^2}{a^{16}}$$

Solution

(a) We apply the power rule first on the first parenthesis.

$$(x^2)^2 \cdot \frac{x^6}{x^4} = x^4 \cdot \frac{x^6}{x^4}$$

Then apply the quotient rule to simplify the fraction.

$$x^4 \cdot \frac{x^6}{x^4} = x^4 \cdot x^2$$

Apply the product rule to simplify.

$$x^4 \cdot x^2 = x^6$$

(b) Simplify inside the first parenthesis by reducing the numbers.

$$\left(\frac{4a^2}{b^5} \right)^3 \cdot \frac{b^2}{a^{16}}$$

Then we can apply the power rule on the first parenthesis.

$$\left(\frac{4a^2}{b^5} \right)^3 \cdot \frac{b^2}{a^{16}} = \frac{64a^6}{b^{15}} \cdot \frac{b^2}{a^{16}}$$

Group like terms together.

$$\frac{64a^6}{b^{15}} \cdot \frac{b^2}{a^{16}} = 64 \cdot \frac{a^6}{a^{16}} \cdot \frac{b^2}{b^{15}}$$

Apply the quotient rule on each fraction.

$$64 \cdot \frac{a^6}{a^{16}} \cdot \frac{b^2}{b^{15}} = \frac{64}{a^{10}b^{13}}$$

Review Questions

Evaluate the following expressions.

1. $\frac{5^6}{5^2}$
2. $\frac{6^7}{6^3}$

3. $\frac{3^4}{3^{10}}$

4. $\left(\frac{2^2}{3^3}\right)^3$

Simplify the following expressions.

5. $\frac{a^3}{a^2}$

6. $\frac{x^5}{x^9}$

7. $\left(\frac{a^3b^4}{a^2b}\right)^3$

8. $\frac{x^6y^2}{x^2y^5}$

9. $\frac{6a^3}{2a^2}$

10. $\frac{15x^5}{5x}$

11. $\left(\frac{18a^4}{15a^{10}}\right)^4$

12. $\frac{25yx^6}{20y^5x^2}$

13. $\left(\frac{x^6y^2}{x^4y^4}\right)^3$

14. $\left(\frac{6a^2}{4b^4}\right)^2 \cdot \frac{5b}{3a}$

15. $\frac{(3ab)^2(4a^3b^4)^3}{(6a^2b)^4}$

16. $\frac{(2a^2bc^2)(6abc^3)}{4ab^2c}$

Review Answers

1. 5^4

2. $6^4 = 1296$

3. $\frac{1}{3^6} = \frac{1}{729}$

4. $\frac{2^6}{3^9} = \frac{64}{19683}$

5. a

6. $\frac{1}{x^4}$

7. a^3b^9

8. $\frac{x^4}{y^3}$

9. $3a$

10. $3x^4$

11. $\frac{1296}{625a^4}$

12. $\frac{5x^4}{4y^4}$

13. $\frac{x^6}{y^6}$

14. $\frac{15a^3}{4b^7}$

15. $\frac{4a^3b^{10}}{9}$

16. $3a^2c^4$

8.3 Zero, Negative, and Fractional Exponents

Learning Objectives

- Simplify expressions with zero exponents.
- Simplify expressions with negative exponents.
- Simplify expression with fractional exponents.
- Evaluate exponential expressions.

Introduction

There are many interesting concepts that arise when contemplating the product and quotient rule for exponents. You may have already been wondering about different values for the exponents. For example, so far we have only considered positive, whole numbers for the exponent. So called **natural numbers** (or **counting numbers**) are easy to consider, but even with the everyday things around us we think about questions such as is it possible to have a negative amount of money? or what would one and a half pairs of shoes look like? In this lesson, we consider what happens when the exponent is not a natural number. We will start with What happens when the exponent is zero?

Simplify Expressions with Exponents of Zero

Let us look again at the quotient rule for exponents (that $\frac{x^n}{x^m} = x^{n-m}$) and consider what happens when $n = m$. Lets take the example of x^4 divided by x^4 .

$$\frac{x^4}{x^4} = x^{(4-4)} = x^0$$

Now we arrived at the quotient rule by considering how the factors of x cancel in such a fraction. Lets do that again with our example of x^4 divided by x^4 .

$$\frac{x^4}{x^4} = \frac{\cancel{x} \cdot \cancel{x} \cdot \cancel{x} \cdot \cancel{x}}{\cancel{x} \cdot \cancel{x} \cdot \cancel{x} \cdot \cancel{x}} = 1$$

So $x^0 = 1$.

This works for any value of the exponent, not just 4.

$$\frac{x^n}{x^n} = x^{n-n} = x^0$$

Since there is the same number of factors in the numerator as in the denominator, they cancel each other out and we obtain $x^0 = 1$. The zero exponent rule says that any number raised to the power zero is one.

Zero Rule for Exponents: $x^0 = 1$, $x \neq 0$

Simplify Expressions With Negative Exponents

Again we will look at the quotient rule for exponents (that $\frac{x^n}{x^m} = x^{n-m}$) and this time consider what happens when $m > n$. Lets take the example of x^4 divided by x^6 .

$$\frac{x^4}{x^6} = x^{(4-6)} = x^{-2} \text{ for } x \neq 0.$$

By the quotient rule our exponent for x is -2 . But what does a negative exponent really mean? Lets do the same calculation long-hand by dividing the factors of x^4 by the factors of x^6 .

$$\frac{x^4}{x^6} = \frac{\cancel{x} \cdot \cancel{x} \cdot \cancel{x} \cdot \cancel{x}}{\cancel{x} \cdot \cancel{x} \cdot \cancel{x} \cdot \cancel{x} \cdot x \cdot x} = \frac{1}{x \cdot x} = \frac{1}{x^2}$$

So we see that x to the power -2 is the same as one divided by x to the power $+2$. Here is the negative power rule for exponents.

Negative Power Rule for Exponents $\frac{1}{x^n} = x^{-n}$ $x \neq 0$

You will also see negative powers applied to products and fractions. For example, here it is applied to a product.

$$\begin{aligned} (x^3y)^{-2} &= x^{-6}y^{-2} && \text{using the power rule} \\ x^{-6}y^{-2} &= \frac{1}{x^6} \cdot \frac{1}{y^2} = \frac{1}{x^6y^2} && \text{using the negative power rule separately on each variable} \end{aligned}$$

Here is an example of a negative power applied to a quotient.

$$\begin{aligned} \left(\frac{a}{b}\right)^{-3} &= \frac{a^{-3}}{b^{-3}} && \text{using the power rule for quotients} \\ \frac{a^{-3}}{b^{-3}} &= \frac{a^{-3}}{1} \cdot \frac{1}{b^{-3}} = \frac{1}{a^3} \cdot \frac{b^3}{1} && \text{using the negative power rule on each variable separately} \\ \frac{1}{a^3} \cdot \frac{b^3}{1} &= \frac{b^3}{a^3} && \text{simplifying the division of fractions} \\ \frac{b^3}{a^3} &= \left(\frac{b}{a}\right)^3 && \text{using the power rule for quotients in reverse.} \end{aligned}$$

The last step is not necessary but it helps define another rule that will save us time. A fraction to a negative power is flipped.

Negative Power Rule for Fractions $\left(\frac{x}{y}\right)^{-n} = \left(\frac{y}{x}\right)^n$, $x \neq 0, y \neq 0$

In some instances, it is more useful to write expressions without fractions and that makes use of negative powers.

Example 1

Write the following expressions without fractions.

- (a) $\frac{1}{x}$
- (b) $\frac{2}{x^2}$
- (c) $\frac{x^2}{y^3}$
- (d) $\frac{3}{xy}$

Solution

We apply the negative rule for exponents $\frac{1}{x^n} = x^{-n}$ on all the terms in the denominator of the fractions.

(a) $\frac{1}{x} = x^{-1}$

(b) $\frac{2}{x^2} = 2x^{-2}$

(c) $\frac{x^2}{y^3} = x^2y^{-3}$

(d) $\frac{3}{xy} = 3x^{-1}y^{-1}$

Sometimes, it is more useful to write expressions without negative exponents.

Example 2

Write the following expressions without negative exponents.

(a) $3x^{-3}$

(b) $a^2b^{-3}c^{-1}$

(c) $4x^{-1}y^3$

(d) $\frac{2x^{-2}}{y^{-3}}$

Solution

We apply the negative rule for exponents $\frac{1}{x^n} = x^{-n}$ on all the terms that have negative exponents.

(a) $3x^{-3} = \frac{3}{x^3}$

(b) $a^2b^{-3}c^{-1} = \frac{a^2}{b^3c}$

(c) $4x^{-1}y^3 = \frac{4y^3}{x}$

(d) $\frac{2x^{-2}}{y^{-3}} = \frac{2y^3}{x^2}$

Example 3

Simplify the following expressions and write them without fractions.

(a) $\frac{4a^2b^3}{2a^5b}$

(b) $\left(\frac{x}{3y^2}\right)^3 \cdot \frac{x^2y}{4}$

Solution

(a) Reduce the numbers and apply quotient rule on each variable separately.

$$\frac{4a^2b^3}{6a^5b} = 2 \cdot a^{2-5} \cdot b^{3-1} = 2a^{-3}b^2$$

(b) Apply the power rule for quotients first.

$$\left(\frac{2x}{y^2}\right)^3 \cdot \frac{x^2y}{4} = \frac{8x^3}{y^6} \cdot \frac{x^2y}{4}$$

Then simplify the numbers, use product rule on the x 's and the quotient rule on the y 's.

$$\frac{8x^3}{y^6} \cdot \frac{x^2y}{4} = 2 \cdot x^{3+2} \cdot y^{1-6} = 2x^5y^{-5}$$

Example 4

Simplify the following expressions and write the answers without negative powers.

(a) $\left(\frac{ab^{-2}}{b^3}\right)^2$

(b) $\frac{x^{-3}y^2}{x^2y^{-2}}$

Solution

(a) Apply the quotient rule inside the parenthesis.

$$\left(\frac{ab^{-2}}{b^3}\right)^2 = (ab^{-5})^2$$

Apply the power rule.

$$(ab^{-5})^2 = a^2b^{-10} = \frac{a^2}{b^{10}}$$

(b) Apply the quotient rule on each variable separately.

$$\frac{x^{-3}y^2}{x^2y^{-2}} = x^{-3-2}y^{2-(-2)} = x^{-5}y^4 = \frac{y^4}{x^5}$$

Simplify Expressions With Fractional Exponents

The exponent rules you learned in the last three sections apply to all powers. So far we have only looked at positive and negative integers. The rules work exactly the same if the powers are fractions or irrational numbers. Fractional exponents are used to express the taking of roots and radicals of something (square roots, cube roots, etc.). Here is an example.

$$\sqrt{a} = a^{1/2} \text{ and } \sqrt[3]{a} = a^{1/3} \text{ and } \sqrt[5]{a^2} = (a^2)^{\frac{1}{5}} = a^{\frac{2}{5}} = a^{2/5}$$

$$\text{Roots as Fractional Exponents } \sqrt[m]{a^n} = a^{n/m}$$

We will examine roots and radicals in detail in a later chapter. In this section, we will examine how exponent rules apply to fractional exponents.

Evaluate Exponential Expressions

When evaluating expressions we must keep in mind the order of operations. You must remember **PEMDAS**.

Evaluate inside the **P**arenthesis.

Evaluate **E**xponents.

Perform **M**ultiplication and **D**ivision operations from left to right.

Perform **A**ddition and **S**ubtraction operations from left to right.

Example 6

Evaluate the following expressions to a single number.

(a) 5^0

(b) 7^2

(c) $\left(\frac{2}{3}\right)^3$

(d) 3^{-3}

Solution(a) $5^0 = 1$ Remember that a number raised to the power 0 is always 1.

(b) $7^2 = 7 \cdot 7 = 49$

(c) $\left(\frac{2}{3}\right)^3 = \frac{2^3}{3^3} = \frac{8}{27}$

(d) $3^{-3} = \frac{1}{3^3} = \frac{1}{27}$

Example 7*Evaluate the following expressions to a single number.*

(a) $3 \cdot 5^5 - 10 \cdot 5 + 1$

(b) $\frac{2 \cdot 4^2 - 3 \cdot 5^2}{3^2}$

(c) $\left(\frac{3^3}{2^2}\right)^{-2} \cdot \frac{3}{4}$

Solution

(a) Evaluate the exponent.

$$3 \cdot 5^2 - 10 \cdot 5 + 1 = 3 \cdot 25 - 10 \cdot 5 + 1$$

Perform multiplications from left to right.

$$3 \cdot 25 - 10 \cdot 5 + 1 = 75 - 50 + 1$$

Perform additions and subtractions from left to right.

$$75 - 50 + 1 = 26$$

(b) Treat the expressions in the numerator and denominator of the fraction like they are in parenthesis.

$$\frac{(2 \cdot 4^2 - 3 \cdot 5^2)}{(3^2 - 2^2)} = \frac{(2 \cdot 16 - 3 \cdot 25)}{(9 - 4)} = \frac{(32 - 75)}{5} = \frac{-43}{5}$$

(c) $\left(\frac{3^3}{2^2}\right)^{-2} \cdot \frac{3}{4} = \left(\frac{2^2}{3^3}\right)^2 \cdot \frac{3}{4} = \frac{2^4}{3^6} \cdot \frac{3}{4} = \frac{2^4}{3^6} \cdot \frac{3}{2^2} = \frac{2^2}{3^5} = \frac{4}{243}$

Example 8*Evaluate the following expressions for $x = 2, y = -1, z = 3$.*

(a) $2x^2 - 3y^3 + 4z$

(b) $(x^2 - y^2)^2$

(c) $\left(\frac{3x^2y^5}{4z}\right)^{-2}$

Solution

(a) $2x^2 - 3y + 4z = 2 \cdot 2^2 - 3 \cdot (-1)^3 + 4 \cdot 3 = 2 \cdot 4 - 3 \cdot (-1) + 4 \cdot 3 = 8 + 3 + 12 = 23$

(b) $(x^2 - y^2)^2 = (2^2 - (-1)^2)^2 = (4 - 1)^2 = 3^2 = 9$

(c) $\left(\frac{3x^2 - y^5}{4z}\right)^{-2} = \left(\frac{3 \cdot 2^2 \cdot (-1)^5}{4 \cdot 3}\right)^{-2} = \left(\frac{3 \cdot 4 \cdot (-1)}{12}\right)^{-2} = \left(\frac{-12}{12}\right)^{-2} = \left(\frac{-1}{1}\right)^{-2} = \left(\frac{1}{-1}\right)^2 = (-1)^2 = 1$

Review Questions

Simplify the following expressions, be sure that there aren't any negative exponents in the answer.

1. $x^{-1} \cdot y^2$
2. x^{-4}
3. $\frac{x^{-3}}{x^{-7}}$
4. $\frac{x^{-3}y^{-5}}{z^{-7}}$
5. $(x^{\frac{1}{2}}y^{-\frac{2}{3}})(x^2y^{\frac{1}{3}})$
6. $\left(\frac{a}{b}\right)^{-2}$
7. $(3a^{-2}b^2c^3)^3$
8. $x^{-3} \cdot x^3$

Simplify the following expressions so that there aren't any fractions in the answer.

9. $\frac{a^{-3}(a^5)}{a^{-6}}$
10. $\frac{5x^6y^2}{x^8y}$
11. $\frac{(4ab^6)^3}{(ab)^5}$
12. $\left(\frac{3x}{y^{1/3}}\right)^3$
13. $\frac{3x^2y^{3/2}}{xy^{1/2}}$
14. $\frac{(3x^3)(4x^4)}{(2y)^2}$
15. $\frac{a^{-2}b^{-3}}{c^{-1}}$
16. $\frac{x^{1/2}y^{5/2}}{x^{3/2}y^{3/2}}$

Evaluate the following expressions to a single number.

17. 3^{-2}
18. $(6.2)^0$
19. $8^{-4} \cdot 8^6$
20. $(16^{\frac{1}{2}})^3$
21. $x^24x^3y^44y^2$ if $x = 2$ and $y = -1$
22. $a^4(b^2)^3 + 2ab$ if $a = -2$ and $b = 1$
23. $5x^2 - 2y^3 + 3z$ if $x = 3$, $y = 2$, and $z = 4$
24. $\left(\frac{a^2}{b^3}\right)^{-2}$ if $a = 5$ and $b = 3$

Review Answers

1. $\frac{y^2}{x}$
2. $\frac{1}{x^4}$
3. x^4

4. $\frac{z^7}{x^3y^5}$
5. $\frac{x^{5/2}}{y^{1/3}}$
6. $\left(\frac{b}{a}\right)^2$ or $\frac{b^2}{a^2}$
7. $\frac{27b^6c^9}{a^6}$
8. 1
9. a^8
10. $5x^{-2}y$
11. $64a^{-2}b^{\frac{1}{3}}$
12. $27x^2y^{-1}$
13. $3xy$
14. $6x^7y^{-2}$
15. $a^{-2}b^{-3}c$
16. $x^{-1}y$
17. 0.111
18. 1
19. 64
20. 64
21. 512
22. 12
23. 41
24. 1.1664

8.4 Scientific Notation

Learning Objectives

- Write numbers in scientific notation.
- Evaluate expressions in scientific notation.
- Evaluate expressions in scientific notation using a graphing calculator.

Introduction Powers of 10

Consider the number six hundred and forty three thousand, two hundred and ninety seven. We write it as 643,297 and each digit's position has a value assigned to it. You may have seen a table like this before.

hundred-thousands	ten-thousands	thousands	hundreds	tens	units (ones)
6	4	3	2	9	7

We have seen that when we write an exponent above a number it means that we have to multiply a certain number of factors of that number together. We have also seen that a zero exponent always gives us one, and negative exponents make fractional answers. Look carefully at the table above. Do you notice that all the column headings are powers of ten? Here they are listed.

$$100,000 = 10^5$$

$$10,000 = 10^4$$

$$1,000 = 10^3$$

$$100 = 10^2$$

$$10 = 10^1$$

Even the units column is really just a power of ten. **Unit** means 1 and $1 = 10^0$.

If we divide 643,297 by 100,000 we get 6.43297. If we multiply this by 100,000 we get back to our original number. But we have just seen that 100,000 is the same as 10^5 , so if we multiply 6.43297 by 10^5 we should also get our original answer. In other words

$$6.43297 \times 10^5 = 643,297$$

So we have found a new way of writing numbers! What do you think happens when we continue the powers of ten? Past the units column down to zero we get into decimals, here the exponent becomes negative.

Writing Numbers Greater Than One in Scientific Notation

Scientific notation numbers are always written in the following form.


$$a \times 10^b$$

Where $1 \leq a < 10$ and b , the exponent, is an integer. This notation is especially useful for numbers that are either very small or very large. When we use scientific notation to write numbers, the exponent on the 10 determines the position of the decimal point.


Look at the following examples.

- $1.07 \times 10^4 = 10,700$
- $1.07 \times 10^3 = 1,070$
- $1.07 \times 10^2 = 107$
- $1.07 \times 10^1 = 10.7$
- $1.07 \times 10^0 = 1.07$
- $1.07 \times 10^{-1} = 0.107$
- $1.07 \times 10^{-2} = 0.0107$
- $1.07 \times 10^{-3} = 0.00107$
- $1.07 \times 10^{-4} = 0.000107$

Look at the first term of the list and examine the position of the decimal point in both expressions.



So the exponent on the ten acts to move the decimal point over to the right. An exponent of 4 moves it 4 places and an exponent of 3 would move it 3 places.



Example 1

Write the following numbers in scientific notation.

- (a) 63
- (b) 9,654
- (c) 653,937,000

(d) 1,000,000,006

(a) $63 = 6.3 \times 10 = 6.3 \times 10^1$

(b) $9,654 = 9.654 \times 1,000 = 9.654 \times 10^3$

(c) $653,937,000 = 6.53937000 \times 100,000,000 = 6.53937 \times 10^8$

(d) $1,000,000,006 = 1.000000006 \times 1,000,000,000 = 1.000000006 \times 10^9$

Example 2*The Sun is approximately 93million miles for Earth. Write this distance in scientific notation.*

This time we will simply write out the number long-hand (with a decimal point) and count decimal places.

Solution

$$\underbrace{93,000,000.0}_{7 \text{ decimal places}} = 9.3 \times 10^7 \text{ miles}$$

A Note on Significant Figures

We often combine scientific notation with rounding numbers. If you look at Example 2, the distance you are given has been rounded. It is unlikely that the distance is **exactly** 93 million miles! Looking back at the numbers in Example 1, if we round the final two answers to 2 significant figures (2 s.f.) they become:

1(c) 6.5×10^8

1(d) 1.0×10^9

Note that the zero after the decimal point has been left in for Example 1(d) to indicate that the result has been rounded. It is important to know when it is OK to round and when it is not.

Writing Numbers Less Than One in Scientific Notation

We have seen how we can use scientific notation to express large numbers, but it is equally good at expressing extremely small numbers. Consider the following example.

Example 3*The time taken for a light beam to cross a football pitch is 0.0000004 seconds. Express this time in scientific notation.*

We will proceed in a similar way as before.

$$0.0000004 = 4 \times 0.0000001 = 4 \times \frac{1}{10,000,000} = 4 \times \frac{1}{10^7} = 4 \times 10^{-7}$$

So...

$$4 \times 10^{-7} = 0.0000004$$

↑ decimal position
↘ 7 decimal places difference

Just as a positive exponent on the ten moves the decimal point that many places to the right, a negative exponent moves the decimal place that many places to the left.

Example 4

Express the following numbers in scientific notation.

- (a) 0.003
 (b) 0.000056
 (c) 0.00005007
 (d) 0.00000000000954

Lets use the method of counting how many places we would move the decimal point before it is after the first non-zero number. This will give us the value for our negative exponent.

- (a) $\underbrace{0.003}_{3 \text{ decimal places}} = 3 \times 10^{-3}$
 (b) $\underbrace{0.000056}_{5 \text{ decimal places}} = 5.6 \times 10^{-5}$
 (c) $\underbrace{0.00005007}_{5 \text{ decimal places}} = 5.007 \times 10^{-5}$
 (d) $\underbrace{0.00000000000954}_{12 \text{ decimal places}} = 9.54 \times 10^{-12}$

Evaluating Expressions in Scientific Notation

When we are faced with products and quotients involving scientific notation, we need to remember the rules for exponents that we learned earlier. It is relatively straightforward to work with scientific notation problems if you remember to deal with all the powers of 10 together. The following examples illustrate this.

Example 5

Evaluate the following expressions and write your answer in scientific notation.

- (a) $(3.2 \times 10^6) \cdot (8.7 \times 10^{11})$
 (b) $(5.2 \times 10^{-4}) \cdot (3.8 \times 10^{-19})$
 (c) $(1.7 \times 10^6) \cdot (2.7 \times 10^{-11})$

The key to evaluating expressions involving scientific notation is to keep the powers of 10 together and deal with them separately. Remember that when we use scientific notation, the leading number **must be between 1 and 10**. We need to move the decimal point over one place to the left. See how this adds 1 to the exponent on the 10.

(a)

$$\begin{aligned} (3.2 \times 10^6) \cdot (8.7 \times 10^{11}) &= \underbrace{3.2 \times 8.7}_{27.84} \times \underbrace{10^6 \times 10^{11}}_{10^{17}} \\ (3.2 \times 10^6) \cdot (8.7 \times 10^{11}) &= 2.784 \times 10^1 \times 10^{17} \end{aligned}$$

Solution

$$(3.2 \times 10^6) \cdot (8.7 \times 10^{11}) = 2.784 \times 10^{18}$$

(b)

$$\begin{aligned}
 (5.2 \times 10^{-4}) \cdot (3.8 \times 10^{-19}) &= \underbrace{5.2 \times 3.8}_{19.76} \times \underbrace{10^{-4} \times 10^{-19}}_{10^{-23}} \\
 &= 1.976 \times 10^1 \times 10^{-23}
 \end{aligned}$$

Solution

$$(5.2 \times 10^{-4}) \cdot (3.8 \times 10^{-19}) = 1.976 \times 10^{-22}$$

(c)

$$(1.7 \times 10^6) \cdot (2.7 \times 10^{-11}) = \underbrace{1.7 \times 2.7}_{4.59} \times \underbrace{10^6 \times 10^{-11}}_{10^{-5}}$$

Solution

$$(1.7 \times 10^6) \cdot (2.7 \times 10^{-11}) = 4.59 \times 10^{-5}$$

Example 6

Evaluate the following expressions. Round to 3 significant figures and write your answer in scientific notation.

(a) $(3.2 \times 10^6) \div (8.7 \times 10^{11})$

(b) $(5.2 \times 10^{-4}) \div (3.8 \times 10^{-19})$

(c) $(1.7 \times 10^6) \div (2.7 \times 10^{-11})$

It will be easier if we convert to fractions and THEN separate out the powers of 10.

(a)

$$\begin{aligned}
 (3.2 \times 10^6) \div (8.7 \times 10^{11}) &= \frac{3.2 \times 10^6}{8.7 \times 10^{11}} \\
 &= \frac{3.2}{8.7} \times \frac{10^6}{10^{11}} \\
 &= 0.368 \times 10^{(6-11)} \\
 &= 3.68 \times 10^{-1} \times 10^{-5}
 \end{aligned}$$

Next we separate the powers of 10.

Evaluate each fraction (round to 3 s.f.):

Remember how to write scientific notation!

Solution

$$(3.2 \times 10^6) \div (8.7 \times 10^{11}) = 3.86 \times 10^{-6} \text{ (rounded to 3 significant figures)}$$

(b)

$$\begin{aligned}
 (5.2 \times 10^{-4}) \div (3.8 \times 10^{-19}) &= \frac{5.2 \times 10^{-4}}{3.8 \times 10^{-19}} \\
 &= \frac{5.2}{3.8} \times \frac{10^{-4}}{10^{-19}} \\
 &= 1.37 \times 10^{((-4) - (-19))} \\
 &= 1.37 \times 10^{15}
 \end{aligned}$$

Separate the powers of 10.

Evaluate each fraction (round to 3 s.f.).

Solution

$$(5.2 \times 10^{-4}) \div (3.8 \times 10^{-19}) = 1.37 \times 10^{15} \text{ (rounded to 3 significant figures)}$$

(c)

$$\begin{aligned} (1.7 \times 10^6) \div (2.7 \times 10^{-11}) &= \frac{1.7 \times 10^6}{2.7 \times 10^{-11}} \\ &= \frac{1.7}{2.7} \times \frac{10^6}{10^{-11}} \\ &= 0.630 \times 10^{6-(-11)} \\ &= 6.30 \times 10^{-1} \times 10^{17} \end{aligned}$$

Next we separate the powers of 10.

Evaluate each fraction (round to 3 s.f.).

Remember how to write scientific notation!

Solution

$$(1.7 \times 10^6) \div (2.7 \times 10^{-11}) = 6.30 \times 10^{16} \text{ (rounded to 3 significant figures)}$$


Note that the final zero has been left in to indicate that the result has been rounded.

Evaluate Expressions in Scientific Notation Using a Graphing Calculator

All scientific and graphing calculators have the ability to use scientific notation. It is extremely useful to know how to use this function.

To insert a number in scientific notation, use the **[EE]** button. This is **[2nd] [,]** on some TI models.

For example to enter 2.6×10^5 enter 2.6**[EE]** 5.




When you hit **[ENTER]** the calculator displays 2.6E5 if its set in **Scientific** mode OR it displays 260000 if its set in **Normal** mode.

(To change the mode, press the Mode key)

Example 7

Evaluate $(1.7 \times 10^6) \div (2.7 \times 10^{-11})$ using a graphing calculator.

[ENTER] 1.7 EE 6 \div 2.7 EE -11 and press **[ENTER]**




The calculator displays $6.296296296E16$ whether it is in Normal mode or Scientific mode. This is the case because the number is so big that it does not fit inside the screen in Normal mode.

Solution

$$(1.7 \times 10^6) \div (2.7 \times 10^{-11}) = 6.\bar{3} \times 10^{16}$$

Example 8

Evaluate $(2.3 \times 10^6) \times (4.9 \times 10^{-10})$ using a graphing calculator.



[ENTER] 2.3 EE 6 \times 4.9 EE -10 and press [ENTER]

The calculator displays .001127 in Normal mode or $1.127E - 3$ in Scientific mode.


Solution

$$(2.3 \times 10^6) \times (4.9 \times 10^{-10}) = 1.127 \times 10^{-3}$$

Example 9

Evaluate $(4.5 \times 10^{14})^3$ using a graphing calculator.

[ENTER] (4.5EE14)³ and press [ENTER].



The calculator displays 9.1125E43

Solution

$$(4.5 \times 10^{14})^3 = 9.1125 \times 10^{43}$$

Solve Real-World Problems Using Scientific Notation

Example 10

The mass of a single lithium atom is approximately one percent of one millionth of one billionth of one billionth of one kilogram. Express this mass in scientific notation.

We know that percent means we divide by 100, and so our calculation for the mass (in kg) is

$$\frac{1}{100} \times \frac{1}{1,000,000} \times \frac{1}{1,000,000,000} \times \frac{1}{1,000,000,000} = 10^{-2} \times 10^{-6} \times 10^{-9} \times 10^{-9} \times 10^{-9}$$

Next, we use the product of powers rule we learned earlier in the chapter.

$$10^{-2} \times 10^{-6} \times 10^{-9} \times 10^{-9} = 10^{((-2)+(-6)+(-9)+(-9))} = 10^{-26} \text{ kg.}$$

Solution

The mass of one lithium atom is approximately 1×10^{-26} kg .

Example 11

You could fit about 3million E. coli bacteria on the head of a pin. If the size of the pin head in question is 1.2×10^{-5} m², calculate the area taken up by one E. coli bacterium. Express your answer in scientific notation.

Since we need our answer in scientific notation it makes sense to convert 3 million to that format first:

$$3,000,000 = 3 \times 10^6$$

Next, we need an expression involving our unknown. The area taken by one bacterium. Call this A.

$$3 \times 10^6 \cdot A = 1.2 \times 10^{-5}$$

Since 3 million of them make up the area of the pin-head.

Isolate A:

$$A = \frac{1}{3 \times 10^6} \cdot 1.2 \times 10^{-5}$$

Rearranging the terms gives

$$A = \frac{1.2}{3} \cdot \frac{1}{10^6} \times 10^{-5}$$

Then using the definition of a negative exponent

$$A = \frac{1.2}{3} \times 10^{-6} \times 10^{-5}$$

Evaluate combine exponents using the product rule.

$$A = 0.4 \times 10^{-11}$$

We cannot, however, leave our answer like this.

Solution

The area of one bacterium $A = 4.0 \times 10^{-12} \text{ m}^2$

Notice that we had to move the decimal point over one place to the right, subtracting 1 from the exponent on the 10.

Review Questions

- Write the numerical value of the following.
 - 3.102×10^2
 - 7.4×10^4
 - 1.75×10^{-3}
 - 2.9×10^{-5}
 - 9.99×10^{-9}
- Write the following numbers in scientific notation.
 - 120,000
 - 1,765,244
 - 12
 - 0.00281
 - 0.000000027
- The moon is approximately a sphere with radius $r = 1.08 \times 10^3$ miles. Use the formula Surface Area = $4\pi r^2$ to determine the surface area of the moon, in square miles. Express your answer in scientific notation, rounded to 2 significant figures.
- The charge on one electron is approximately 1.60×10^{-19} coulombs. One **Faraday** is equal to the total charge on 6.02×10^{23} electrons. What, in coulombs, is the charge on one Faraday?
- Proxima Centauri, the next closest star to our Sun is approximately 2.5×10^{13} miles away. If light from Proxima Centauri takes 3.7×10^4 hours to reach us from there, calculate the speed of light in miles per hour. Express your answer in scientific notation, rounded to 2 significant figures.

Review Answers

- 310.2
 - 74,000
 - 0.00175
 - 0.000029

- (e) 0.00000000999
2. (a) 1.2×10^5
(b) 1.765224×10^{10}
(c) 1.2×10^1
(d) 2.81×10^{-3}
(e) 2.7×10^{-8}
3. 1.5×10^7 miles²
4. 96,320 or 9.632×10^4
5. 6.8×10^8 miles per hour

8.5 Addition and Subtraction of Polynomials

Learning Objectives

- Write a polynomial expression in standard form.
- Classify polynomial expression by degree
- Add and subtract polynomials
- Problem solving using addition and subtraction of polynomials

Introduction

So far we have seen functions described by straight lines (linear functions) and functions where the variable appeared in the exponent (exponential functions). In this section we will introduce polynomial functions. A **polynomial** is made up of different terms that contain **positive integer** powers of the variables. Here is an example of a polynomial.

$$4x^3 + 2x^2 - 3x + 1$$

Each part of the polynomial that is added or subtracted is called a **term** of the polynomial. The example above is a polynomial with *four terms*.

$$4x^3 + 2x^2 - 3x + 1$$

The numbers appearing in each term in front of the variable are called the **coefficients**. The number appearing all by itself without a variable is called a **constant**.

$$4x^3 + 2x^2 - 3x + 1$$

In this case, the coefficient of x^3 is 4, the coefficient of x^2 is 2, the coefficient of x is -3 and the constant is 1.

Degrees of Polynomials and Standard Form

Each term in the polynomial has a **degree**. This is the power of the variable in that term.

$4x^3$ Has a degree of 3 and is called a **cubic term** or **3rd order term**.

$2x^2$ Has a degree of 2 and is called the **quadratic term** or **2nd order term**.

$-3x$ Has a degree of 1 and is called the **linear term** or **1st order term**.

1 Has a degree of 0 and is called the **constant**.

By definition, **the degree of the polynomial** is the same as the degree of the term with the highest degree. This example is a polynomial of degree 3, which is also called a "cubic" polynomial. (Why do you think it is called a cubic?).

Polynomials can have more than one variable. Here is another example of a polynomial.

$$t^4 - 6s^3t^2 - 12st + 4s^4 - 5$$

This is a polynomial because all exponents on the variables are positive integers. This polynomial has five terms. Lets look at each term more closely. **Note:** *The degree of a term is the sum of the powers on each variable in the term.*

t^4 Has a degree of 4, so its a 4^{th} order term

$-6s^3t^2$ Has a degree of 5, so its a 5^{th} order term.

$-12st$ Has a degree of 2, so its a 2^{nd} order term

$4s^4$ Has a degree of 4, so its a 4^{th} order term

-5 Is a constant, so its degree is 0.

Since the highest degree of a term in this polynomial is 5, then this is polynomial of degree 5 or a 5^{th} order polynomial.

A polynomial that has only one term has a special name. It is called a **monomial** (*mono means one*). A monomial can be a constant, a variable, or a product of a constant and one or more variables. You can see that each term in a polynomial is a monomial. A polynomial is the sum of monomials. Here are some examples of monomials.

b^2
This is a monomial

8
So is this

$-2ab^2$
and this

$\frac{1}{4}x^4$
and this

$-29xy$
and this

Example 1

For the following polynomials, identify the coefficient on each term, the degree of each term and the degree of the polynomial.

a) $x^5 - 3x^3 + 4x^2 - 5x + 7$

b) $x^4 - 3x^3y^2 + 8x - 12$

Solution

a) $x^5 - 3x^3 + 4x^2 - 5x + 7$

The coefficients of each term in order are 1, -3 , 4 , -5 and the constant is 7.

The degrees of each term are 5, 3, 2, 1, and 0. Therefore, the degree of the polynomial is 5.

b) $x^4 - 3x^3y^2 + 8x - 12$

The coefficients of each term in order are 1, -3 , 8 and the constant is -12 .

The degrees of each term are 4, 5, 1, and 0. Therefore, the degree of the polynomial is 5.

Example 2

Identify the following expressions as polynomials or non-polynomials.

a) $5x^2 - 2x$

b) $3x^2 - 2x^{-2}$

c) $x\sqrt{x} - 1$

d) $\frac{5}{x^3+1}$

e) $4x^{1/3}$

f) $4xy^2 - 2x^2y - 3 + y^3 - 3x^3$

Solution

(a) $5x^2 - 2x$ This *is* a polynomial.

(b) $3x^2 - 2x^{-2}$ This is *not* a polynomial because it has a negative exponent.

(c) $x\sqrt{x} - 1$ This is *not* a polynomial because it has a square root.

(d) $\frac{5}{x^3+1}$ This is *not* a polynomial because the power of x appears in the denominator.

(e) $4x^{1/3}$ This is *not* a polynomial because it has a fractional exponent.

(f) $4xy^2 - 2x^2y - 3 + y^3 - 3x^3$ This *is* a polynomial.

You saw that each term in a polynomial has a degree. The degree of the highest term is also the degree of the polynomial. Often, we arrange the terms in a polynomial so that the term with the highest degree is first and it is followed by the other terms in order of decreasing power. This is called **standard form**.

The following polynomials are in standard form.

$$4x^4 - 3x^3 + 2x^2 - x + 1$$

$$a^4b^3 - 2a^3b^3 + 3a^4b - 5ab^2 + 2$$

The first term of a polynomial in standard form is called the **leading term** and the coefficient of the leading term is called the **leading coefficient**.

The first polynomial above has a leading term of $4x^4$ and a leading coefficient of 4.

The second polynomial above has a leading term of a^4b^3 and a leading coefficient of 1.

Example 3

Rearrange the terms in the following polynomials so that they are in standard form. Indicate the leading term and leading coefficient of each polynomial.

(a) $7 - 3x^3 + 4x$

(b) $ab - a^3 + 2b$

(c) $-4b + 4 + b^2$

Solution

(a) $7 - 3x^3 + 4x$ is rearranged as $-3x^3 + 4x + 7$. The leading term is $-3x^3$ and the leading coefficient is -3 .

(b) $ab - a^3 + 2b$ is rearranged as $-a^3 + ab + 2b$. The leading term is $-a^3$ and the leading coefficient is -1 .

(c) $-4b + 4 + b^2$ is rearranged as $b^2 - 4b + 4$. The leading term is b^2 and the leading coefficient is 1.

Simplifying Polynomials

A polynomial is simplified if it has no terms that are alike. **Like terms** are terms in the polynomial that have the same variable(s) with the same exponents, but they can have different coefficients.

$2x^2y$ and $5x^2y$ are like terms.

$6x^2y$ and $6xy^2$ are not like terms.

If we have a polynomial that has like terms, we simplify by combining them.

$$x^2 + \overbrace{6xy}^{\nearrow} - \overbrace{4xy}^{\nwarrow} + y^2$$

Like terms

This sample polynomial simplified by combining the like terms $6xy - 4xy = 2xy$. We write the simplified polynomial as

$$x^2 + 2xy + y^2$$

Example 4

Simplify the following polynomials by collecting like terms and combining them.

(a) $2x - 4x^2 + 6 + x^2 - 4 + 4x$

(b) $a^3b^3 - 5ab^4 + 2a^3b - a^3b^3 + 3ab^4 - a^2b$

Solution

(a) $2x - 4x^2 + 6 + x^2 - 4 + 4x$

Rearrange the terms so that like terms are grouped together

$$= (-4x^2 + x^2) + (2x + 4x) + (6 - 4)$$

Combine each set of like terms by adding or subtracting the coefficients

$$= -3x^2 + 6x + 2$$

(b) $a^3b^3 - 5ab^4 + 2a^3b - a^3b^3 + 3ab^4 - a^2b$

Rearrange the terms so that like terms are grouped together:

$$= (a^3b^3 - a^3b^3) + (-5ab^4 + 3ab^4) + 2a^3b - a^2b$$

Combine each set of like terms:

$$= 0 - 2ab^4 + 2a^3b - a^2b$$

$$= -2ab^4 + 2a^3b - a^2b$$

Add and Subtract Polynomials

Polynomial addition

To add two or more polynomials, write their sum and then simplify by combining like terms.

Example 5

Add and simplify the resulting polynomials.

(a) Add $3x^2 - 4x + 7$ and $2x^3 - 4x^2 - 6x + 5$.

(b) Add $x^2 - 2xy + y^2$ and $2y^2 - 4x^2$ and $10xy + y^3$.

Solution:

(a) Add $3x^2 - 4x + 7$ and $2x^3 - 4x^2 - 6x + 5$

$$\begin{aligned} & \text{Group like terms} & & = (3x^2 - 4x + 7) + (2x^3 - 4x^2 - 6x + 5) \\ & \text{Simplify} & & = 2x^3 + (3x^2 - 4x^2) + (-4x - 6x) + (7 + 5) \\ & & & = 2x^3 - x^2 - 10x + 12 \end{aligned}$$

(b) Add $x^2 - 2xy + y^2$ and $2y^2 - 3x^2$ and $10xy + y^3$

$$\begin{aligned} & \text{Group like terms} & & = (x^2 - 2xy + y^2) + (2y^2 - 3x^2) + (10xy + y^3) \\ & \text{Simplify} & & = (x^2 - 3x^2) + (y^2 + 2y^2) + (-2xy + 10xy) + y^3 \\ & & & = 2x^2 + 3y^2 + 8xy + y^3 \end{aligned}$$

Polynomial subtraction

To subtract one polynomial from another, add the opposite of each term of the polynomial you are subtracting.

Example 6

Subtract and simplify the resulting polynomials.

a) Subtract $x^3 - 3x^2 + 8x + 12$ from $4x^2 + 5x - 9$.

b) Subtract $5b^2 - 2a^2$ from $4a^2 - 8ab - 9b^2$.

Solution

a)

$$\begin{aligned} (4x^2 + 5x - 9) - (x^3 - 3x^2 + 8x + 12) &= (4x^2 + 5x - 9) + (-x^3 + 3x^2 - 8x - 12) \\ \text{Group like terms} &= -x^3 - (4x^2 + 3x^2) + (5x - 8x) + (-9 - 12) \\ \text{Simplify} &= -x^3 + 7x^2 - 3x - 21 \end{aligned}$$

b)

$$\begin{aligned} (4a^2 - 8ab - 9b^2) - (5b^2 - 2a^2) &= (4a^2 - 8ab - 9b^2) + (-5b^2 + 2a^2) \\ \text{Group like terms} &= (4a^2 + 2a^2) + (-9b^2 - 5b^2) - 8ab \\ \text{Simplify} &= 6a^2 - 14b^2 - 8ab \end{aligned}$$

Note: An easy way to check your work after adding or subtracting polynomials is to substitute a convenient value in for the variable, and check that your answer and the problem both give the same value. For example, in part (b) of Example 6, if we let $a = 2$ and $b = 3$, then we can check as follows.

Given

$$\begin{aligned}
 &(4a^2 - 8ab - 9b^2) - (5b^2 - 2a^2) \\
 &(4(2)^2 - 8(2)(3) - 9(3)^2) - (5(3)^2 - 2(2)^2) \\
 &(4(4) - 8(2)(3) - 9(9)) - (5(9) - 2(4)) \\
 &(-113) - 37 \\
 &- 150
 \end{aligned}$$

Solution

$$\begin{aligned}
 &6a^2 - 14b^2 - 8ab \\
 &6(2)^2 - 14(3)^2 - 8(2)(3) \\
 &6(4) - 14(9) - 8(2)(3) \\
 &24 - 126 - 48 \\
 &- 150
 \end{aligned}$$

Since both expressions evaluate to the same number when we substitute in arbitrary values for the variables, we can be reasonably sure that our answer is correct. Note, when you use this method, do not choose 0 or 1 for checking since these can lead to common problems.


Problem Solving Using Addition or Subtraction of Polynomials

An application of polynomials is their use in finding areas of a geometric object. In the following examples, we will see how the addition or subtraction of polynomials might be useful in representing different areas.


Example 7

Write a polynomial that represents the area of each figure shown.


a)





b)



c)



d)

**Solutions**

a) This shape is formed by two squares and two rectangles.


The blue square has area: $y \cdot y = y^2$

The yellow square has area: $x \cdot x = x^2$

The pink rectangles each have area: $x \cdot y = xy$

To find the total area of the figure we add all the separate areas.

$$\begin{aligned} \text{Total area} &= y^2 + x^2 + xy + xy \\ &= y^2 + x^2 + 2xy \end{aligned}$$




b) This shape is formed by two squares and one rectangle.

The yellow squares each have an area: $a \cdot a = a^2$.

The orange rectangle has area: $2a \cdot b = 2ab$.

To find the total area of the figure we add all the separate areas.

$$\begin{aligned} \text{Total area} &= a^2 + a^2 + 2ab \\ &= 2a^2 + 2ab \end{aligned}$$




c) To find the area of the green region we find the area of the big square and subtract the area of the little square.

The big square has area $y \cdot y = y^2$.

The little square has area $x \cdot x = x^2$.

Area of the green region $= y^2 - x^2$




d) To find the area of the figure we can find the area of the big rectangle and add the areas of the pink squares.

The pink squares each have area: $a \cdot a = a^2$.

The blue rectangle has area: $3a \cdot a = 3a^2$.

To find the total area of the figure we add all the separate areas.

$$\text{Total area} = a^2 + a^2 + a^2 + 3a^2 = 6a^2$$



Another way to find this area is to find the area of the big square and subtract the areas of the three yellow squares.

The big square has area: $3a \cdot 3a = 9a^2$.

The yellow squares each have areas: $a \cdot a = a^2$.

To find the total area of the figure we subtract:

$$\begin{aligned} \text{Area} &= 9a^2 - (a^2 + a^2 + a^2) \\ &= 9a^2 - 3a^2 \\ &= 6a^2 \end{aligned}$$

Review Questions

Indicate which expressions are polynomials.

1. $x^2 + 3x^{1/2}$
2. $\frac{1}{3}x^2y - 9y^2$
3. $3x^{-3}$
4. $\frac{2}{3}t^2 - \frac{1}{t^2}$

Express each polynomial in standard form. Give the degree of each polynomial.

5. $3 - 2x$
6. $8 - 4x + 3x^3$
7. $-5 + 2x - 5x^2 + 8x^3$
8. $x^2 - 9x^4 + 12$





Add and simplify.

9. $(x + 8) + (-3x - 5)$
10. $(-2x^2 + 4x - 12) + (7x + x^2)$
11. $(2a^2b - 2a + 9) + (5a^2b - 4b + 5)$
12. $(6.9a^2 - 2.3b^2 + 2ab) + (3.1a - 2.5b^2 + b)$

Subtract and simplify.

13. $(-t + 15t^2) - (5t^2 + 2t - 9)$
14. $(-y^2 + 4y - 5) - (5y^2 + 2y + 7)$
15. $(-5m^2 - m) - (3m^2 + 4m - 5)$
16. $(2a^2b - 3ab^2 + 5a^2b^2) - (2a^2b^2 + 4a^2b - 5b^2)$

Find the area of the following figures.



Review Answers

1. No

2. yes
3. no
4. no
5. $-2x + 3$; Degree = 1
6. $3x^3 - 4x + 8$; Degree = 3
7. $8x^3 - 5x^2 + 2x - 5$; Degree = 3
8. $-9x^4 + x^2 + 12$; Degree = 4
9. $-2x + 3$
10. $-x^2 + 11x - 12$
11. $7a^2b - 2a - 4b + 14$
12. $6.9a^2 - 4.8b^2 + 2ab + 3.1a + b$
13. $-3t + 9$
14. $-6y^2 + 2y - 12$
15. $-8m^2 - 5m + 5$
16. $-2a^2b - 3ab^2 + 3a^2b^2 + 5b^2$
17. Area = $2xz - xy$
18. Area = $4ab + ac$
19. $2xy - 2x^2$
20. Area = $3ab$

8.6 Multiplication of Polynomials

Learning Objectives

- Multiply a polynomial by a monomial
- Multiply a polynomial by a binomial
- Solve problems using multiplication of polynomials

Introduction

When multiplying polynomials we must remember the exponent rules that we learned in the last chapter.

The Product Rule $x^n \cdot x^m = x^{n+m}$

This says that if we multiply expressions that have the same base, we just add the exponents and keep the base unchanged.

If the expressions we are multiplying have coefficients and more than one variable, we multiply the coefficients just as we would any number and we apply the product rule on each variable separately.

$$(2x^2y^3) \cdot (3x^2y) = (2 \cdot 3) \cdot (x^2 + 2) \cdot (y^3 + 1) = 6x^4y^4$$

Multiplying a Polynomial by a Monomial

We begin this section by multiplying a monomial by a monomial. As you saw above, we need to multiply the coefficients separately and then apply the exponent rules to each variable separately. Lets try some examples.

Example 1

Multiply the following monomials.

- $(2x^2)(5x^3)$
- $(-3y^4)(2y^2)$
- $(3xy^5)(-6x^4y^2)$
- $(-12a^2b^3c^4)(-3a^2b^2)$

Solution


- $(2x^2)(5x^3) = (2 \cdot 5) \cdot (x^2 \cdot x^3) = 10x^{2+3} = 10x^5$
- $(-3y^4)(2y^2) = (-3 \cdot 2) \cdot (y^4 \cdot y^2) = -6y^{4+2} = -6y^6$
- $(3xy^5)(-6x^4y^2) = 18x^{1+4}y^{5+2} = -18x^5y^7$
- $(-12a^2b^3c^4)(-3a^2b^2) = 36a^{2+2}b^{3+2}c^4 = 36a^4b^5c^4$

To multiply a polynomial by a monomial, we use the **Distributive Property**.

This says that

$$a(b + c) = ab + ac$$

This property is best illustrated by an area problem. We can find the area of the big rectangle in two ways.



One way is to use the formula for the area of a rectangle.

$$\text{Area of the big rectangle} = \text{length} \cdot \text{width}$$

$$\text{Length} = a, \text{Width} = b + c$$

$$\text{Area} = a \cdot (b + c)$$

The area of the big rectangle can also be found by adding the areas of the two smaller rectangles.

$$\text{Area of red rectangle} = ab$$

$$\text{Area of blue rectangle} = ac$$

$$\text{Area of big rectangle} = ab + ac$$

This means that $a(b + c) = ab + ac$. It shows why the Distributive Property works.

This property is useful for working with numbers and also with variables.

For instance, to solve this problem, you would add 2 and 7 to get 9 and then multiply by 5 to get 45. But there is another way to do this.

$$5(2 + 7) = 5 \cdot 2 + 5 \cdot 7$$

It means that each number in the parenthesis is multiplied by 5 separately and then the products are added together.

$$5(2 + 7) = 5 \cdot 2 + 5 \cdot 7 = 10 + 35 = 45$$

In general, if we have a number or variable in front of a parenthesis, this means that each term in the parenthesis is multiplied by the expression in front of the parenthesis. The distributive property works no matter how many terms there are inside the parenthesis.

$$a(b + c + d + e + f + \dots) = ab + ac + ad + ae + af + \dots$$

The ... means and so on.

Lets now apply this property to multiplying polynomials by monomials.

Example 2

Multiply

a) $3(x^2 + 3x - 5)$

b) $4x(3x^2 - 7)$

c) $-7y(4y^2 - 2y + 1)$

Solution

a) $3(x^2 + 3x - 5) = 3(x^2) + 3(3x) - 3(5) = 3x^2 + 9x - 15$

b) $4x(3x^2 - 7) = (4x)(3x^2) + (4x)(-7) = 12x^3 - 28x$

c) $-7y(4y^2 - 2y + 1) = (-7y)(4y^2) + (-7y)(-2y) + (-7y)(1) = -28y^3 + 14y^2 - 7y$

Notice that the use of the Distributive Property simplifies the problems to just multiplying monomials by monomials and adding all the separate parts together.

Example 3

Multiply

a) $2x^3(-3x^4 + 2x^3 - 10x^2 + 7x + 9)$

b) $-7a^2bc^3(5a^2 - 3b^2 - 9c^2)$

Solution

a)

$$\begin{aligned} 2x^3(-3x^4 + 2x^3 - 10x^2 + 7x + 9) &= (2x^3)(-3x^4) + (2x^3)(2x^3) + (2x^3)(-10x^2) + (2x^3)(7x) + (2x^3)(9) \\ &= -6x^7 + 4x^6 - 20x^5 + 14x^4 + 18x^3 \end{aligned}$$

b)

$$\begin{aligned} -7a^2bc^3(5a^2 - 3b^2 - 9c^2) &= (-7a^2bc^3)(5a^2) + (-7a^2bc^3)(-3b^2) + (-7a^2bc^3)(-9c^2) \\ &= -35a^4bc^3 + 21a^2b^3c^3 + 63a^2bc^5 \end{aligned}$$

Multiply a Polynomial by a Binomial

Lets start by multiplying two binomials together. A binomial is a polynomial with two terms, so a product of two binomials will take the form.

$$(a + b)(c + d)$$

The Distributive Property also applies in this situation. Lets think of the first parenthesis as one term. The Distributive Property says that the term in front of the parenthesis multiplies with each term inside the parenthesis separately. Then, we add the results of the products.

$$(a + b)(c + d) = (a + b) \cdot c + (a + b) \cdot d$$

Lets rewrite this answer as $c \cdot (a + b) + d \cdot (a + b)$

We see that we can apply the distributive property on each of the parenthesis in turn.

$$c \cdot (a + b) + d \cdot (a + b) = c \cdot a + c \cdot b + d \cdot a + d \cdot b \text{ (or } ca + cb + da + db)$$

What you should notice is that when multiplying any two polynomials, **every term in one polynomial is multiplied by every term in the other polynomial.**

Lets look at some examples of multiplying polynomials.

Example 4

Multiply and simplify $(2x + 1)(x + 3)$

Solution

We must multiply each term in the first polynomial with each term in the second polynomial.

Lets try to be systematic to make sure that we get all the products.

First, multiply the first term in the first parenthesis by all the terms in the second parenthesis.

$$(2x + 1)(x + 3) = (2x)(x) + (2x)(3) + \dots$$

We are now done with the first term.

Now we multiply the second term in the first parenthesis by all terms in the second parenthesis and add them to the previous terms.

$$(2x + 1)(x + 3) = (2x)(x) + (2x)(3) + (1)(x) + (1)(3)$$

We are done with the multiplication and we can simplify.

$$\begin{aligned} (2x)(x) + (2x)(3) + (1)(x) + (1)(3) &= 2x^2 + 6x + x + 3 \\ &= 2x^2 + 7x + 3 \end{aligned}$$

This way of multiplying polynomials is called **in-line** multiplication or **horizontal** multiplication.

Another method for multiplying polynomials is to use **vertical** multiplication similar to the vertical multiplication you learned with regular numbers. Lets demonstrate this method with the same example.

$$\begin{array}{r} 2x + 1 \\ x + 3 \\ \hline \end{array}$$

$$6x + 3 \leftarrow \text{Multiply each term on top by } 3$$

$$\text{Multiply each term on top by } x \rightarrow 2x^2 + x$$

$$\begin{array}{r} 2x^2 + 7x + 3 \leftarrow \text{Arrange like terms on top of each other and add vertically} \end{array}$$

This method is typically easier to use although it does take more space. Just make sure that like terms are together in vertical columns so you can combine them at the end.

Example 5

Multiply and simplify

(a) $(4x - 5)(x - 20)$

(b) $(3x - 2)(3x + 2)$

(c) $(3x^2 + 2x - 5)(2x - 3)$

(d) $(x^2 - 9)(4x^4 + 5x^2 - 2)$

Solution

a) $(4x - 5)(x - 20)$

Horizontal multiplication

$$\begin{aligned}(4x - 5)(x - 20) &= (4x)(x) + (4x)(-20) + (-5)(x) + (-5)(-20) \\ &= 4x^2 - 80x - 5x + 100 = 4x^2 - 85x + 100\end{aligned}$$

Vertical multiplication

Arrange the polynomials on top of each other with like terms in the same columns.

$$\begin{array}{r}4x - 5 \\ x - 20 \\ \hline-80x + 100 \\ 4x^2 - 5x \\ \hline 4x^2 - 85x + 100\end{array}$$


Both techniques result in the same answer, $4x^2 - 85x + 100$.

For the last question, we'll show the solution with vertical multiplication because it may be a technique you are not used to. Horizontal multiplication will result in the exact same terms and the same answer.

b) $(3x - 2)(3x + 2)$


$$\begin{array}{r}3x - 2 \\ 3x + 2 \\ \hline 6x - 4 \\ 9x^2 - 6x \\ \hline 9x^2 + 0x - 4\end{array}$$

Answer $9x^2 - 4$




Find the volumes of the following figures


c)



d)



Solution



a) We use the formula for the area of a rectangle.


$$\text{Area} = \text{length} \cdot \text{width}$$

For the big rectangle

$$\text{Length} = b + 3, \text{ Width} = b + 2$$

$$\begin{aligned} \text{Area} &= (b + 3)(b + 2) \\ &= b^2 + 2b + 3b + 6 \\ &= b^2 + 5b + 6 \end{aligned}$$

b) Lets find the area of the big rectangle in the second figure and subtract the area of the yellow rectangle.




$$\text{Area of big rectangle} = 20(12) = 240$$

$$\begin{aligned} \text{Area of yellow rectangle} &= (12 - x)(20 - 2x) \\ &= 240 - 24x - 20x + 2x^2 \\ &= 240 - 44x + 2x^2 \\ &= 2x^2 - 44x + 240 \end{aligned}$$

The desired area is the difference between the two.


$$\begin{aligned} \text{Area} &= 240 - (2x^2 - 44x + 240) \\ &= 240 + (-2x^2 + 44x - 240) \\ &= 240 - 2x^2 + 44x - 240 \\ &= -2x^2 + 44x \end{aligned}$$



c) The volume of this shape = (area of the base) · (height).

$$\begin{aligned}\text{Area of the base} &= x(x+2) \\ &= x^2 + 2x\end{aligned}$$

$$\begin{aligned}\text{Height} &= 2x + 1 \\ \text{Volume} &= (x^2 + 2x)(2x + 1) \\ &= 2x^3 + x^2 + 4x^2 + 2x \\ &= 2x^3 + 5x^2 + 2x\end{aligned}$$



d) The volume of this shape = (area of the base) · (height).

$$\begin{aligned}\text{Area of the base} &= (4a - 3)(2a + 1) \\ &= 8a^2 + 4a - 6a - 3 \\ &= 8a^2 - 2a - 3\end{aligned}$$

$$\begin{aligned}\text{Height} &= a + 4 \\ \text{Volume} &= (8a^2 - 2a - 3)(a + 4)\end{aligned}$$

Lets multiply using the vertical method:

$$\begin{array}{r} 8a^2 - 2a - 3 \\ a + 4 \\ \hline 32a^2 - 8a - 12 \\ 8a^3 - 2a^2 - 3a \\ \hline 8a^3 + 30a^2 - 11a - 12 \end{array}$$

Answer Volume = $8a^3 + 30a^2 - 11a - 12$

Review Questions



Multiply the following monomials.

- $(2x)(-7x)$
- $(-5a^2b)(-12a^3b^3)$
- $(3xy^2z^2)(15x^2yz^3)$


Multiply and simplify.


- $2x(4x - 5)$
- $9x^3(3x^2 - 2x + 7)$
- $-3a^2b(9a^2 - 4b^2)$
- $(x - 3)(x + 2)$
- $(a^2 + 2)(3a^2 - 4)$
- $(7x - 2)(9x - 5)$
- $(2x - 1)(2x^2 - x + 3)$
- $(3x + 2)(9x^2 - 6x + 4)$
- $(a^2 + 2a - 3)(a^2 - 3a + 4)$

Find the areas of the following figures.



Find the volumes of the following figures.



**Review Answers**

1. $-14x^2$
2. $60a^5b^4$
3. $45x^3y^3z^5$
4. $8x^2 - 10x$
5. $27x^5 - 18x^4 + 63x^3$
6. $-27a^4b + 12a^2b^3$
7. $x^2 - x - 6$
8. $3a^4 + 2a^2 - 8$
9. $63x^2 - 53x + 10$
10. $4x^3 - 4x^2 + 7x - 3$
11. $27x^3 + 8$
12. $a^4 - a^3 - 5a^2 + 17a - 12$
13. $(2x + 4)(x + 6) = 2x^2 + 16x + 24$
14. $x(3x + 8) = 3x^2 + 8x$
15. $6x^3 + 14x^2 + 8x$
16. $24x^3 - 28x^2 - 12x$

8.7 Special Products of Polynomials

Learning Objectives

- Find the square of a binomial
- Find the product of binomials using sum and difference formula
- Solve problems using special products of polynomials

Introduction

We saw that when we multiply two binomials we need to make sure that each term in the first binomial multiplies with each term in the second binomial. Let's look at another example.

Multiply two linear (i.e. with degree = 1) binomials:

$$(2x + 3)(x + 4)$$

When we multiply, we obtain a quadratic (i.e. with degree = 2) polynomial with four terms.

$$2x^2 + 8x + 3x + 12$$

The middle terms are like terms and we can combine them. We simplify and get:

$$2x^2 + 11x + 12$$

This is a quadratic or 2nd degree **trinomial** (polynomial with three terms).

You can see that every time we multiply two linear binomials with one variable, we will obtain a quadratic polynomial. In this section we will talk about some special products of binomials.

Find the Square of a Binomial

A special binomial product is the **square of a binomial**. Consider the following multiplication.

$$(x + 4)(x + 4)$$

Since we are multiplying the same expression by itself that means that we are squaring the expression. This means that:

$$(x + 4)(x + 4) = (x + 4)^2$$

Let's multiply:

$$(x + 4)(x + 4) = x^2 + 4x + 4x + 16$$

And combine like terms:

$$= x^2 + 8x + 16$$

Notice that the middle terms are the same. Is this a coincidence? In order to find that out, let's square a general linear binomial.

$$\begin{aligned}(a + b)^2 &= (a + b)(a + b) = a^2 + ab + ab + b^2 \\ &= a^2 + 2ab + b^2\end{aligned}$$

It looks like the middle terms are the same again. So far we have squared the sum of binomials. Let's now square a difference of binomials.

$$\begin{aligned}(a - b)^2 &= (a - b)(a - b) = a^2 - ab - ab + b^2 \\ &= a^2 - 2ab + b^2\end{aligned}$$

We notice a pattern when squaring binomials. To square a binomial, add the square of the first term, add or subtract twice the product of the terms, and the square of the second term. You should remember these formulas:

Square of a Binomial

$$(a + b)^2 = a^2 + 2ab + b^2 \text{ and } (a - b)^2 = a^2 - 2ab + b^2$$

Remember! A polynomial that is raised to an exponent means that we multiply the polynomial by itself however many times the exponent indicates. For instance

$$(a + b)^2 = (a + b)(a + b)$$

Don't make the common mistake $(a + b)^2 = a^2 + b^2$. To see why $(a + b)^2 \neq a^2 + b^2$ try substituting numbers for a and b into the equation (for example, $a = 4$ and $b = 3$), and you will see that it is *not* a true statement. The middle term, $2ab$, is needed to make the equation work.

We can apply the formulas for squaring binomials to any number of problems.

Example 1

Square each binomial and simplify.

- (a) $(x + 10)^2$
- (b) $(2x - 3)^2$
- (c) $(x^2 + 4)^2$
- (d) $(5x - 2y)^2$

Solution

Let's use the square of a binomial formula to multiply each expression.

a) $(x + 10)^2$

If we let $a = x$ and $b = 10$, then

$$\begin{aligned}(a^2 + b) &= a^2 + 2a b + b^2 \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ (x + 10)^2 &= (x)^2 + 2(x)(10) + (10)^2 \\ &= x^2 + 20x + 100\end{aligned}$$

b) $(2x - 3)^2$

If we let $a = 2x$ and $b = 3$, then

$$\begin{aligned}(a - b)^2 &= a^2 - 2ab + b^2 \\ (2x - 3)^2 &= (2x)^2 - 2(2x)(3) + (3)^2 \\ &= 4x^2 - 12x + 9\end{aligned}$$

c) $(x + 4)^2$

If we let $a = x^2$ and $b = 4$, then

$$\begin{aligned}(x^2 + 4)^2 &= (x^2)^2 + 2(x^2)(4) + (4)^2 \\ &= x^4 + 8x^2 + 16\end{aligned}$$

d) $(5x - 2y)^2$

If we let $a = 5x$ and $b = 2y$, then

$$\begin{aligned}(5x - 2y)^2 &= (5x)^2 - 2(5x)(2y) + (2y)^2 \\ &= 25x^2 - 20xy + 4y^2\end{aligned}$$

Find the Product of Binomials Using Sum and Difference Patterns

Another special binomial product is the product of a sum and a difference of terms. For example, lets multiply the following binomials.

$$\begin{aligned}(x + 4)(x - 4) &= x^2 - 4x + 4x - 16 \\ &= x^2 - 16\end{aligned}$$

Notice that the middle terms are opposites of each other, so they cancel out when we collect like terms. This is not a coincidence. This always happens when we multiply a sum and difference of the same terms.

$$\begin{aligned}(a + b)(a - b) &= a^2 - ab - b^2 \\ &= a^2 - b^2\end{aligned}$$

When multiplying a sum and difference of the same two terms, the middle terms cancel out. We get the square of the first term minus the square of the second term. You should remember this formula.

Sum and Difference Formula

$$(a + b)(a - b) = a^2 - b^2$$

Lets apply this formula to a few examples.

Example 2

Multiply the following binomials and simplify.

(a) $(x+3)(x-3)$

(b) $(5x+9)(5x-9)$

(c) $(2x^3+7)(2x^3-7)$

(d) $(4x+5y)(4x-5y)$

Solution

(a) Let $a = x$ and $b = 3$, then

$$\begin{array}{cccccc} (a+b)(a-b) & = & a^2 & - & b^2 & \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\ (x+3)(x-3) & = & (x)^2 & - & (3)^2 & \\ & & = & x^2 & - & 9 \end{array}$$

(b) Let $a = 5x$ and $b = 9$, then

$$\begin{array}{cccccc} (a+b)(a-b) & = & a^2 & - & b^2 & \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\ (5x+9)(5x-9) & = & (5x)^2 & - & (9)^2 & \\ & & = & 25x^2 & - & 81 \end{array}$$

(c) Let $a = 2x^3$ and $b = 7$, then


$$\begin{array}{cccccc} (2x^3+7)(2x^3-7) & = & (2x^3)^2 & - & (7)^2 & \\ & & = & 4x^6 & - & 49 \end{array}$$

(d) Let $a = 4x$ and $b = 5y$, then

$$\begin{array}{cccccc} (4x+5y)(4x-5y) & = & (4x)^2 & - & (5y)^2 & \\ & & = & 16x^2 & - & 25y^2 \end{array}$$

Solve Real-World Problems Using Special Products of Polynomials

Lets now see how special products of polynomials apply to geometry problems and to mental arithmetic.



Example 3

Find the area of the following square

Solution

The area of the square = side \times side

$$\begin{aligned}\text{Area} &= (a + b)(a + b) \\ &= a^2 + 2ab + b^2\end{aligned}$$

Notice that this gives a visual explanation of the square of binomials product.

$$\begin{array}{rclcl} \text{Area of the big square} & = & \text{area of the blue square} & + & 2(\text{area of yellow rectangle}) & + & \text{area of red square} \\ (a + b)^2 & = & a^2 & + & 2ab & + & b^2 \end{array}$$

The next example shows how to use the special products in doing fast mental calculations.

Example 4

Use the difference of squares and the binomial square formulas to find the products of the following numbers without using a calculator.

- (a) 43×57
- (b) 112×88
- (c) 45^2
- (d) 481×309

Solution

The key to these mental tricks is to rewrite each number as a sum or difference of numbers you know how to square easily.

(a) Rewrite $43 = (50 - 7)$ and $57 = (50 + 7)$.

$$\text{Then } 43 \times 57 = (50 - 7)(50 + 7) = (50)^2 - (7)^2 = 2500 - 49 = 2,451$$

(b) Rewrite $112 = (100 + 12)$ and $88 = (100 - 12)$

$$\text{Then } 112 \times 88 = (100 + 12)(100 - 12) = (100)^2 - (12)^2 = 10,000 - 144 = 9,856$$

(c) $45^2 = (40 + 5)^2 = (40)^2 + 2(40)(5) + (5)^2 = 1600 + 400 + 25 = 2,025$

(d) Rewrite $481 = (400 + 81)$ and $319 = (400 - 81)$

$$\text{Then, } 481 \times 319 = (400 + 81)(400 - 81) = (400)^2 - (81)^2$$

$(400)^2$ is easy - it equals 160,000

$(81)^2$ is not easy to do mentally. Lets rewrite it as $81 = 80 + 1$

$$(81)^2 = (80 + 1)^2 = (80)^2 + 2(80)(1) + (1)^2 = 6400 + 160 + 1 = 6,561$$

$$\text{Then, } 481 \times 309 = (400)^2 - (81)^2 = 160,000 - 6,561 = 153,439$$

Review Questions


Use the special product for squaring binomials to multiply these expressions.

1. $(x+9)^2$
2. $(3x-7)^2$
3. $(4x^2+y^2)^2$
4. $(8x-3)^2$

Use the special product of a sum and difference to multiply these expressions.

5. $(2x-1)(2x+1)$
6. $(x-12)(x+12)$
7. $(5a-2b)(5a+2b)$
8. $(ab-1)(ab+1)$

Find the area of the orange square in the following figure. It is the lower right shaded box.



Multiply the following numbers using the special products.

10. 45×55
11. 56^2
12. 1002×998
13. 36×44

Review Answers

1. $x^2 + 18x + 81$
2. $9x^2 - 42x + 49$
3. $16x^4 + 8x^2y^2 + y^4$
4. $64x^2 - 48x + 9$
5. $4x^2 - 1$
6. $x^2 - 144$
7. $25a^2 - 4b^2$
8. $a^2b^2 - 1$
9. Area = $(a-b)^2 = a^2 - 2ab + b^2$
10. $(50-5)(50+5) = 2475$
11. $(50+6)^2 = 3136$
12. $(1000+2)(1000-2) = 999,996$
13. $(40-4)(40+4) = 1584$

8.8 Division of Polynomials

Learning Objectives

- Divide a polynomials by a monomial.
- Divide a polynomial by a binomial.
- Rewrite and graph rational functions.

Introduction

A **rational expression** is formed by taking the quotient of two polynomials.

Some examples of rational expressions are

a) $\frac{2x}{x^2-1}$

b) $\frac{4x^2-3x+4}{2x}$

c) $\frac{9x^2+4x-5}{x^2+5x-1}$

d) $\frac{2x^3}{2x+3}$

Just as with rational numbers, the expression on the top is called the **numerator** and the expression on the bottom is called the **denominator**. In special cases we can simplify a rational expression by dividing the numerator by the denominator.

Divide a Polynomial by a Monomial

We start by dividing a polynomial by a monomial. To do this, we divide each term of the polynomial by the monomial. When the numerator has different terms, the term on the bottom of the fraction serves as **common denominator** to all the terms in the numerator.

Example 1

Divide.

a) $\frac{8x^2-4x+16}{2}$

b) $\frac{3x^3-6x-1}{x}$

c) $\frac{-3x^2-18x+6}{9x}$

Solution

$$\begin{aligned}\frac{8x^2-4x+16}{2} &= \frac{8x^2}{2} - \frac{4x}{2} + \frac{16}{2} = 4x^2 - 2x + 8 \\ \frac{3x^3-6x-1}{x} &= \frac{3x^3}{x} + \frac{6x}{x} - \frac{1}{x} = 3x^2 + 6 - \frac{1}{x} \\ \frac{-3x^2-18x+6}{9x} &= \frac{3x^2}{9x} - \frac{18x}{9x} + \frac{6}{9x} = -\frac{x}{3} - 2 + \frac{2}{3x}\end{aligned}$$

A common error is to cancel the denominator with just one term in the numerator.

Consider the quotient $\frac{3x+4}{4}$

Remember that the denominator of 4 is common to both the terms in the numerator. In other words we are dividing both of the terms in the numerator by the number 4.

The correct way to simplify is

$$\frac{3x+4}{4} = \frac{3x}{4} + \frac{4}{4} = \frac{3x}{4} + 1$$

A common mistake is to cross out the number 4 from the numerator and the denominator

$$\frac{\cancel{3x+4}}{\cancel{4}} = 3x$$

This is incorrect because the term $3x$ does not get divided by 4 as it should be.

Example 2

Divide $\frac{5x^3-10x^2+x-25}{-5x^2}$.

Solution

$$\frac{5x^3 - 10x^2 + x - 25}{-5x^2} = \frac{5x^3}{-5x^2} - \frac{10x^2}{-5x^2} + \frac{x}{-5x^2} - \frac{25}{-5x^2}$$

The negative sign in the denominator changes all the signs of the fractions:

$$-\frac{5x^3}{5x^2} + \frac{10x^2}{5x^2} - \frac{x}{5x^2} + \frac{25}{5x^2} = -x + 2 - \frac{1}{5x} + \frac{5}{x^2}$$

Divide a Polynomial by a Binomial

We divide polynomials in a similar way that we perform long division with numbers. We will explain the method by doing an example.

Example 3

Divide $\frac{x^2+4x+5}{x+3}$.

Solution: When we perform division, the expression in the numerator is called the **dividend** and the expression in the denominator is called the **divisor**.

To start the division we rewrite the problem in the following form.

$$x + 3 \overline{) x^2 + 4x + 5}$$

We start by dividing the first term in the dividend by the first term in the divisor $\frac{x^2}{x} = x$.

We place the answer on the line above the x term.

$$\begin{array}{r} x \\ x + 3 \overline{) x^2 + 4x + 5} \end{array}$$

Next, we multiply the x term in the answer by each of the $x + 3$ in the divisor and place the result under the divided matching like terms.

$$\begin{array}{r} x \\ x + 3 \overline{) x^2 + 4x + 5} \end{array}$$

$$x(x + 3) = x^2 + 3x$$

Now subtract $x^2 + 3x$ from $x^2 + 4x + 5$. It is useful to change the signs of the terms of $x^2 + 3x$ to $-x^2 - 3x$ and add like terms vertically.

$$\begin{array}{r} x \\ x + 3 \overline{) x^2 + 4x + 5} \\ \underline{-x^2 - 3x} \\ x \end{array}$$

Now, bring down 5, the next term in the dividend.

$$\begin{array}{r} x \\ x + 3 \overline{) x^2 + 4x + 5} \\ \underline{-x^2 - 3x} \\ x + 5 \end{array}$$

We repeat the procedure.

First divide the first term of $x + 5$ by the first term of the divisor $\left(\frac{x}{x}\right) = 1$.

Place this answer on the line above the constant term of the dividend,

$$\begin{array}{r} x + 1 \\ x + 3 \overline{) x^2 + 4x + 5} \\ \underline{-x^2 - 3x} \\ x + 5 \end{array}$$

Multiply 1 by the divisor $x + 3$ and write the answer below $x + 5$ matching like terms.

$$\begin{array}{r} x + 1 \\ x + 3 \overline{) x^2 + 4x + 5} \\ \underline{-x^2 - 3x} \\ x + 5 \end{array}$$

$$1(x + 3) = x + 3$$

Answer $\frac{4x^2-25x-21}{x-7} = 4x + 3$

Check $(4x + 3)(x - 7) + 0 = 4x^2 - 25x - 21$. The answer checks out.

Review Questions

Divide the following polynomials:

1. $\frac{2x+4}{2}$
2. $\frac{x-4}{x}$
3. $\frac{5x-35}{5x}$
4. $\frac{x^2+2x-5}{x}$
5. $\frac{4x^2+12x-36}{-4x}$
6. $\frac{2x^2+10x+7}{2x^2}$
7. $\frac{x^3-x}{-2x^2}$
8. $\frac{5x^4-9}{3x}$
9. $\frac{x^3-12x^2+3x-4}{12x^2}$
10. $\frac{3-6x+x^3}{-9x^3}$
11. $\frac{x^2+3x+6}{x+1}$
12. $\frac{x^2-9x+6}{x-1}$
13. $\frac{x^2+5x+4}{x+4}$
14. $\frac{x^2-10x+25}{x-5}$
15. $\frac{x^2-20x+12}{x-3}$
16. $\frac{3x^2-x+5}{x-2}$
17. $\frac{9x^2+2x-8}{x+4}$
18. $\frac{3x^2-4}{3x+1}$
19. $\frac{5x^2+2x-9}{2x-1}$
20. $\frac{x^2-6x-12}{5x+4}$

Review Answers

1. $x + 2$
2. $1 - \frac{4}{x}$
3. $1 - \frac{7}{x}$
4. $x + 2 - \frac{5}{x}$
5. $-x - 3 + \frac{9}{x}$
6. $1 + \frac{5}{x} + \frac{7}{2x^2}$
7. $-\frac{x}{2} + \frac{1}{2x}$
8. $\frac{5x^3}{3} - \frac{3}{x}$
9. $\frac{x}{12} - 1 + \frac{1}{4x} - \frac{1}{3x^2}$
10. $-\frac{1}{3x^3} + \frac{2}{3x^2} - \frac{1}{9}$
11. $x + 2 + \frac{4}{x+1}$
12. $x - 8 - \frac{2}{x-1}$
13. $x + 1$
14. $x - 5$
15. $x - 17 - \frac{39}{x-3}$

16. $3x + 5 + \frac{15}{x-2}$

17. $9x - 34 + \frac{128}{x+4}$

18. $x - \frac{1}{3} - \frac{11}{3(3x+1)}$

19. $\frac{5}{2}x + \frac{9}{4} - \frac{27}{4(2x-1)}$

20. $\frac{1}{5}x - \frac{34}{25} - \frac{164}{25(5x+4)}$

CHAPTER

9**Introduction to Factoring****Chapter Outline**

- 9.1 POLYNOMIAL EQUATIONS IN FACTORED FORM**
 - 9.2 FACTORING QUADRATIC EXPRESSIONS**
 - 9.3 FACTORING SPECIAL PRODUCTS**
 - 9.4 FACTORING POLYNOMIALS COMPLETELY**
-


9.1 Polynomial Equations in Factored Form

Learning Objectives

- Use the zero-product property
- Find greatest common monomial factor
- Solve simple polynomial equations by factoring

Introduction

In the last few sections, we learned how to multiply polynomials. We did that by using the Distributive Property. All the terms in one polynomial must be multiplied by all terms in the other polynomial. In this section, you will start learning how to do this process in reverse. The reverse of distribution is called **factoring**.



Lets look at the areas of the rectangles again: Area = length \cdot width. The total area of the figure on the right can be found in two ways.

Method 1 Find the areas of all the small rectangles and add them

$$\text{Blue rectangle} = ab$$

$$\text{Orange rectangle} = ac$$

$$\text{Red rectangle} = ad$$

$$\text{Green rectangle} = ae$$

$$\text{Pink rectangle} = 2a$$

$$\text{Total area} = ab + ac + ad + ae + 2a$$

Method 2 Find the area of the big rectangle all at once

$$\text{Length} = a$$

$$\text{Width} = b + c + d + e + 2$$

$$\text{Area} = a(b + c + d + e + 2)$$

Since the area of the rectangle is the same no matter what method you use then the answers are the same:

$$ab + ac + ad + ae + 2a = a(b + c + d + e + 2)$$

Factoring means that you take the factors that are common to all the terms in a polynomial. Then, multiply them by a parenthesis containing all the terms that are left over when you divide out the common factors.

Use the Zero-Product Property

Polynomials can be written in **expanded form** or in **factored form**. Expanded form means that you have sums and differences of different terms:

$$6x^4 + 7x^3 - 26x^2 + 17x + 30$$

Notice that the degree of the polynomials is four. It is written in standard form because the terms are written in order of decreasing power.

Factored form means that the polynomial is written as a product of different factors. The factors are also polynomials, usually of lower degree. Here is the same polynomial in factored form.

$$\underbrace{x-1}_{1^{\text{st}} \text{ factor}} \underbrace{x+2}_{2^{\text{nd}} \text{ factor}} \underbrace{2x-3}_{3^{\text{rd}} \text{ factor}} \underbrace{3x+5}_{4^{\text{th}} \text{ factor}}$$

Notice that each factor in this polynomial is a binomial. Writing polynomials in factored form is very useful because it helps us solve polynomial equations. Before we talk about how we can solve polynomial equations of degree 2 or higher, let's review how to solve a linear equation (degree 1).

Example 1

Solve the following equations

a) $x - 4 = 0$

b) $3x - 5 = 0$

Solution

Remember that to solve an equation you are trying to find the value of x :

a)

$$x - 4 = 0$$

$$+ 4 = +4$$

$$\underline{\underline{x = 4}}$$

b)

$$\begin{array}{r}
 3x - 5 = 0 \\
 + 5 = +5 \\
 \hline
 3x = 5 \\
 \frac{3x}{3} = \frac{5}{3} \\
 x = \frac{5}{3}
 \end{array}$$

Now we are ready to think about solving equations like $2x^2 + 5x = 42$. Notice we can't isolate x in any way that you have already learned. But, we can subtract 42 on both sides to get $2x^2 + 5x - 42 = 0$. Now, the left hand side of this equation can be factored!

Factoring a polynomial allows us to break up the problem into easier chunks. For example, $2x^2 + 5x - 42 = (x + 6)(2x - 7)$. So now we want to solve: $(x + 6)(2x - 7) = 0$

How would we solve this? If we multiply two numbers together and their product is zero, what can we say about these numbers? The only way a product is zero is if one or both of the terms are zero. This property is called the **Zero-product Property**.

How does that help us solve the polynomial equation? Since the product equals 0, then either of the terms or factors in the product must equal zero. We set each factor equal to zero and we solve.

$$(x + 6) = 0 \qquad \text{OR} \qquad (2x - 7) = 0$$

We can now solve each part individually and we obtain:

$$\begin{array}{r}
 x + 6 = 0 \\
 x = -6
 \end{array}
 \qquad \text{or} \qquad
 \begin{array}{r}
 2x - 7 = 0 \\
 2x = 7 \\
 x = \frac{7}{2}
 \end{array}$$

Notice that the solution is $x = -6$ **OR** $x = 7/2$. The **OR** says that either of these values of x would make the product of the two factors equal to zero. Lets plug the solutions back into the equation and check that this is correct.

$$\begin{array}{l}
 \text{Check } x = -6; \\
 (x + 6)(2x - 7) = \\
 (-6 + 6)(2(6) - 7) = \\
 (0)(5) = 0
 \end{array}$$

Check $x = 7/2$

$$\begin{aligned}(x+6)(2x-7) &= \\ \left(\frac{7}{2}+6\right)\left(2\cdot\frac{7}{2}-7\right) &= \\ \left(\frac{19}{2}\right)(7-7) &= \\ \left(\frac{19}{2}\right)(0) &= 0\end{aligned}$$

Both solutions check out. You should notice that the product equals to zero because each solution makes one of the factors simplify to zero. Factoring a polynomial is very useful because the Zero-product Property allows us to break up the problem into simpler separate steps.

If we are not able to factor a polynomial the problem becomes harder and we must use other methods that you will learn later.

As a last note in this section, keep in mind that the Zero-product Property only works when a product equals to zero. For example, if you multiplied two numbers and the answer was nine you could not say that each of the numbers was nine. In order to use the property, you must have the factored polynomial equal to zero.

Example 2

Solve each of the polynomials

a) $(x-9)(3x+4) = 0$

b) $x(5x-4) = 0$

c) $4x(x+6)(4x-9) = 0$

Solution

Since all polynomials are in factored form, we set each factor equal to zero and solve the simpler equations separately

a) $(x-9)(3x+4) = 0$ can be split up into two linear equations

$$\begin{array}{lll}x-9=0 & \text{or} & 3x+4=0 \\ & & 3x=-4 \\ x=9 & \text{or} & x=-\frac{4}{3}\end{array}$$

b) $x(5x-4) = 0$ can be split up into two linear equations

$$\begin{array}{lll}x=0 & \text{or} & 5x-4=0 \\ & & 5x=4 \\ & & x=\frac{4}{5}\end{array}$$

c) $4x(x+6)(4x-9) = 0$ can be split up into three linear equations.

$$\begin{array}{lcl}
 4x = 0 & & 4x - 9 = 0 \\
 x = \frac{0}{4} & \text{or} & x + 6 = 0 \\
 x = 0 & & x = -6
 \end{array}$$

Find Greatest Common Monomial Factor

Once we get a polynomial in factored form, it is easier to solve the polynomial equation. But first, we need to learn how to factor. There are several factoring methods you will be learning in the next few sections. In most cases, factoring takes several steps to complete because we want to **factor completely**. That means that we factor until we cannot factor anymore.

Lets start with the simplest case, finding the greatest monomial factor. When we want to factor, we always look for common monomials first. Consider the following polynomial, written in expanded form.

$$ax + bx + cx + dx$$

A common factor can be a number, a variable or a combination of numbers and variables that appear in all terms of the polynomial. We are looking for expressions that divide out evenly from each term in the polynomial. Notice that in our example, the factor x appears in all terms. Therefore x is a **common factor**

$$ax + bx + cx + dx$$

Since x is a common factor, we factor it by writing in front of a parenthesis:

$$x (\quad)$$

Inside the parenthesis, we write what is left over when we divide x from each term.

$$x(a + b + c + d)$$

Lets look at more examples.

Example 3

Factor

a) $2x + 8$

b) $15x - 25$

c) $3a + 9b + 6$

Solution

a) We see that the factor 2 divides evenly from both terms.

$$2x + 8 = 2(x) + 2(4)$$

We factor the 2 by writing it in front of a parenthesis.

$$2(\quad)$$

Inside the parenthesis, we write what is left from each term when we divide by 2.

$$2(x + 4) \text{ This is the factored form.}$$

b) We see that the factor of 5 divides evenly from all terms.

$$\text{Rewrite } 15x - 25 = 5(3x) - 5(5)$$

$$\text{Factor 5 to get } 5(3x - 5)$$

c) We see that the factor of 3 divides evenly from all terms.

$$\text{Rewrite } 3a + 9b + 6 = 3(a) + 3(3b) + 3(2)$$

$$\text{Factor 3 to get } 3(a + 3b + 2)$$

Here are examples where different powers of the common factor appear in the polynomial

Example 4

Find the greatest common factor

$$\text{a) } a^3 - 3a^2 + 4a$$

$$\text{b) } 12a^4 - 5a^3 + 7a^2$$

Solution

a) Notice that the factor a appears in all terms of $a^3 - 3a^2 + 4a$ but each term has a different power of a . The common factor is the lowest power that appears in the expression. In this case the factor is a .

$$\text{Lets rewrite } a^3 - 3a^2 + 4a = a(a^2) + a(-3a) + a(4)$$

$$\text{Factor } a \text{ to get } a(a^2 - 3a + 4)$$

b) The factor a appears in all the term and the lowest power is a^2 .

$$\text{We rewrite the expression as } 12a^4 - 5a^3 + 7a^2 = 12a^2 \cdot a^2 - 5a \cdot a^2 + 7 \cdot a^2$$

$$\text{Factor } a^2 \text{ to get } a^2(12a^2 - 5a + 7)$$

Lets look at some examples where there is more than one common factor.

Example 5:

Factor completely

$$\text{a) } 3ax + 9a$$

$$\text{b) } x^3y + xy$$

$$\text{c) } 5x^3y - 15x^2y^2 + 25xy^3$$

Solution

a) Notice that 3 is common to both terms.

$$\text{When we factor 3 we get } 3(ax + 3a)$$

This is not completely factored though because if you look inside the parenthesis, we notice that a is also a common factor.

$$\text{When we factor } a \text{ we get } 3 \cdot a(x + 3)$$

This is the answer because there are no more common factors.

A different option is to factor **all** common factors at once.

Since both 3 and a are common we factor the term $3a$ and get $3a(x + 3)$.

b) Notice that both x and y are common factors.

Lets rewrite the expression $x^3y + xy = xy(x^2) + xy(1)$

When we factor xy we obtain $xy(x^2 + 1)$

c) The common factors are $5xy$.

When we factor $5xy$ we obtain $5xy(x^2 - 3xy + 5y^2)$

Solve Simple Polynomial Equations by Factoring

Now that we know the basics of factoring, we can solve some simple polynomial equations. We already saw how we can use the Zero-product Property to solve polynomials in factored form. Here you will learn how to solve polynomials in expanded form. These are the steps for this process.

Step 1

If necessary, **re-write** the equation in standard form such that:

Polynomial expression = 0

Step 2

Factor the polynomial completely

Step 3

Use the zero-product rule to set **each factor equal to zero**

Step 4

Solve each equation from step 3

Step 5

Check your answers by substituting your solutions into the original equation

Example 6

Solve the following polynomial equations

a) $x^2 - 2x = 0$

b) $2x^2 = 5x$

c) $9x^2y - 6xy = 0$

Solution:

a) $x^2 - 2x = 0$

Rewrite this is not necessary since the equation is in the correct form.

Factor The common factor is x , so this factors as: $x(x - 2) = 0$.

Set each factor equal to zero.

$$x = 0$$

or

$$x - 2 = 0$$

Solve

$$x = 0 \qquad \qquad \qquad \text{or} \qquad \qquad \qquad x = 2$$

Check Substitute each solution back into the original equation.

$$x = 0 \qquad \Rightarrow \qquad (0)^2 - 2(0) = 0 \qquad \text{works out}$$

$$x = 2 \qquad \Rightarrow \qquad (2)^2 - 2(2) = 4 - 4 = 0 \qquad \text{works out}$$

Answer $x = 0, x = 2$

b) $2x^2 = 5x$

Rewrite $2x^2 = 5x \Rightarrow 2x^2 - 5x = 0$.

Factor The common factor is x , so this factors as: $x(2x - 5) = 0$.

Set each factor equal to zero:

$$x = 0 \qquad \qquad \qquad \text{or} \qquad \qquad \qquad 2x - 5 = 0$$

Solve

$$\underline{x = 0} \qquad \qquad \qquad \text{or} \qquad \qquad \qquad 2x = 5$$

$$\qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad x = \frac{5}{2}$$

Check Substitute each solution back into the original equation.

$$x = 0 \Rightarrow 2(0)^2 = 5(0) \Rightarrow 0 = 0 \qquad \text{works out}$$

$$x = \frac{5}{2} \Rightarrow 2 \left(\frac{5}{2} \right)^2 = 5 \cdot \frac{5}{2} \Rightarrow 2 \cdot \frac{25}{4} = \frac{25}{2} \Rightarrow \frac{25}{2} = \frac{25}{2} \qquad \text{works out}$$

Answer $x = 0, x = 5/2$

c) $9x^2y - 6xy = 0$

Rewrite Not necessary

Factor The common factor is $3xy$, so this factors as $3xy(3x - 2) = 0$.

Set each factor equal to zero.

$3 = 0$ is never true, so this part does not give a solution

$$x = 0 \qquad \qquad \qquad \text{or} \qquad \qquad \qquad y = 0 \qquad \qquad \qquad \text{or} \qquad \qquad \qquad 3x - 2 = 0$$

Solve

$$x = 0 \qquad \text{or} \qquad y = 0 \qquad \text{or} \qquad 3x = 2$$

$$\qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad x = \frac{2}{3}$$

Check Substitute each solution back into the original equation.

$$x = 0 \Rightarrow 9(0)y - 6(0)y = 0 - 0 = 0$$

works out

$$y = 0 \Rightarrow 9x^2(0) - 6x = 0 - 0 = 0$$

works out

$$\frac{2}{3} \Rightarrow 9 \cdot \left(\frac{2}{3}\right)^2 y - 6 \cdot \frac{2}{3}y = 9 \cdot \frac{4}{9}y - 4y = 4y - 4y = 0$$

works out

Answer $x = 0, y = 0, x = 2/3$

Review Questions

Factor the common factor in the following polynomials.

- $3x^3 - 21x$
- $5x^6 + 15x^4$
- $4x^3 + 10x^2 - 2x$
- $-10x^6 + 12x^5 - 4x^4$
- $12xy + 24xy^2 + 36xy^3$
- $5a^3 - 7a$
- $45y^{12} + 30y^{10}$
- $16xy^2z + 4x^3y$

Solve the following polynomial equations.

- $x(x + 12) = 0$
- $(2x + 1)(2x - 1) = 0$
- $(x - 5)(2x + 7)(3x - 4) = 0$
- $2x(x + 9)(7x - 20) = 0$
- $18y - 3y^2 = 0$
- $9x^2 = 27x$
- $4a^2 + a = 0$
- $b^2 - 5/3b = 0$

Review Answers

- $3x(x^2 - 7)$
- $5x^4(x^2 + 3)$
- $2x(2x^2 + 5x - 1)$
- $2x^4(-5x^2 + 6x - 2)$
- $12xy(1 + 2y + 3y^2)$
- $a(5a^2 - 7)$
- $15y^{10}(3y^2 + 2)$

8. $4xy(4yz + x^2)$
9. $x = 0, x = -12$
10. $x = -1/2, x = 1/2$
11. $x = 5, x = -7/2, x = 4/3$
12. $x = 0, x = -9, x = 20/7$
13. $y = 0, y = 6$
14. $x = 0, x = 3$
15. $a = 0, a = -1/4$
16. $b = 0, b = 5/3$

9.2 Factoring Quadratic Expressions

Learning Objectives

- Write quadratic equations in standard form.
- Factor quadratic expressions for different coefficient values.
- Factor when $a = -1$.

Write Quadratic Expressions in Standard Form

Quadratic polynomials are polynomials of 2^{nd} degree. The standard form of a quadratic polynomial is written as

$$ax^2 + bx + c$$

Here a, b , and c stand for constant numbers. Factoring these polynomials depends on the values of these constants. In this section, we will learn how to factor quadratic polynomials for different values of a, b , and c . In the last section, we factored common monomials, so you already know how to factor quadratic polynomials where $c = 0$.

For example for the quadratic $ax^2 + bx$, the common factor is x and this expression is factored as $x(ax + b)$. When all the coefficients are not zero these expressions are also called **Quadratic Trinomials**, since they are polynomials with three terms.

Factor when $a = 1$, b is Positive, and c is Positive

Lets first consider the case where $a = 1, b$ is positive and c is positive. The quadratic trinomials will take the following form.

$$x^2 + bx + c$$

You know from multiplying binomials that when you multiply two factors $(x + m)(x + n)$ you obtain a quadratic polynomial. Lets multiply this and see what happens. We use The Distributive Property.

$$(x + m)(x + n) = x^2 + nx + mx + mn$$

To simplify this polynomial we would combine the like terms in the middle by adding them.

$$(x + m)(x + n) = x^2 + (n + m)x + mn$$

To factor we need to do this process in reverse.

We see that

$$x^2 + (n + m)x + mn$$

Is the same form as

$$x^2 + bx + c$$

This means that we need to find two numbers m and n where

$$n + m = b$$

and

$$mn = c$$

To factor $x^2 + bx + c$, the answer is the product of two parentheses.

$$(x + m)(x + n)$$

so that $n + m = b$ and $mn = c$

Lets try some specific examples.

Example 1

Factor $x^2 + 5x + 6$

Solution We are looking for an answer that is a product of two binomials in parentheses.

$$(x + \underline{\quad})(x + \underline{\quad})$$

To fill in the blanks, we want two numbers m and n that multiply to 6 and add to 5. A good strategy is to list the possible ways we can multiply two numbers to give us 6 and then see which of these pairs of numbers add to 5. The number six can be written as the product of.

$$6 = 1 \cdot 6$$

and

$$1 + 6 = 7$$

$$6 = 2 \cdot 3$$

and

$$2 + 3 = 5$$

←

This is the correct choice.

So the answer is $(x + 2)(x + 3)$.

We can check to see if this is correct by multiplying $(x + 2)(x + 3)$.

$$\begin{array}{r} x + 2 \\ x + 3 \\ \hline 3x + 6 \\ x^2 + 2x \\ \hline x^2 + 5x + 9 \end{array}$$

The answer checks out.

Example 2

Factor $x^2 + 7x + 12$

Solution

We are looking for an answer that is a product of two parentheses $(x + \underline{\quad})(x + \underline{\quad})$.

The number 12 can be written as the product of the following numbers.

$$\begin{array}{llll}
 12 = 1 \cdot 12 & \text{and} & 1 + 12 = 13 & \\
 12 = 2 \cdot 6 & \text{and} & 2 + 6 = 8 & \\
 12 = 3 \cdot 4 & \text{and} & 3 + 4 = 7 & \leftarrow \text{ This is the correct choice.}
 \end{array}$$

The answer is $(x + 3)(x + 4)$.

Example 3

Factor $x^2 + 8x + 12$.

Solution

We are looking for an answer that is a product of the two parentheses $(x + \underline{\quad})(x + \underline{\quad})$.

The number 12 can be written as the product of the following numbers.

$$\begin{array}{llll}
 12 = 1 \cdot 12 & \text{and} & 1 + 12 = 13 & \\
 12 = 2 \cdot 6 & \text{and} & 2 + 6 = 8 & \leftarrow \text{ This is the correct choice.} \\
 12 = 3 \cdot 4 & \text{and} & 3 + 4 = 7 &
 \end{array}$$

The answer is $(x + 2)(x + 6)$.

Example 4

Factor $x^2 + 12x + 36$.

Solution

We are looking for an answer that is a product of the two parentheses $(x + \underline{\quad})(x + \underline{\quad})$.

The number 36 can be written as the product of the following numbers.

$$\begin{array}{llll}
 36 = 1 \cdot 36 & \text{and} & 1 + 36 = 37 & \\
 36 = 2 \cdot 18 & \text{and} & 2 + 18 = 20 & \\
 36 = 3 \cdot 12 & \text{and} & 3 + 12 = 15 & \\
 36 = 4 \cdot 9 & \text{and} & 4 + 9 = 13 & \\
 36 = 6 \cdot 6 & \text{and} & 6 + 6 = 12 & \leftarrow \text{ This is the correct choice}
 \end{array}$$

The answer is $(x + 6)(x + 6)$.

Factor when a = 1, b is Negative and c is Positive

Now lets see how this method works if the middle coefficient (b) is negative.

Example 5

Factor $x^2 - 6x + 8$.

Solution

We are looking for an answer that is a product of the two parentheses $(x + \underline{\quad})(x + \underline{\quad})$.

The number 8 can be written as the product of the following numbers.

$8 = 1 \cdot 8$ and $1 + 8 = 9$ Notice that these are two different choices.

But also,

$$\begin{array}{llll} 8 = (-1) \cdot (-8) & \text{and} & -1 + (-8) = -9 & \text{Notice that these are two different choices.} \\ 8 = 2 \cdot 4 & \text{and} & 2 + 4 = 6 & \end{array}$$

But also,

$$8 = (-2) \cdot (-4) \quad \text{and} \quad -2 + (-4) = -6 \quad \leftarrow \quad \text{This is the correct choice.}$$

The answer is $(x - 2)(x - 4)$

We can check to see if this is correct by multiplying $(x - 2)(x - 4)$.

$$\begin{array}{r} x - 2 \\ x - 4 \\ \hline -4x + 8 \\ x^2 - 2x \\ \hline x^2 - 6x + 8 \end{array}$$

The answer checks out.

Example 6

Factor $x^2 - 17x + 16$

Solution

We are looking for an answer that is a product of two parentheses: $(x \pm \underline{\quad})(x \pm \underline{\quad})$.

The number 16 can be written as the product of the following numbers:

$$\begin{array}{llll} 16 = 1 \cdot 16 & \text{and} & 1 + 16 = 17 & \\ 16 = (-1) \cdot (-16) & \text{and} & -1 + (-16) = -17 & \leftarrow \quad \text{This is the correct choice.} \\ 16 = 2 \cdot 8 & \text{and} & 2 + 8 = 10 & \\ 16 = (-2) \cdot (-8) & \text{and} & -2 + (-8) = -10 & \\ 16 = 4 \cdot 4 & \text{and} & 4 + 4 = 8 & \\ 16 = (-4) \cdot (-4) & \text{and} & -4 + (-4) = -8 & \end{array}$$

The answer is $(x - 1)(x - 16)$.

Factor when a = 1 and c is Negative

Now let's see how this method works if the constant term is negative.

Example 7

Factor $x^2 + 2x - 15$

Solution

We are looking for an answer that is a product of two parentheses $(x \pm \underline{\quad})(x \pm \underline{\quad})$.

In this case, we must take the negative sign into account. The number -15 can be written as the product of the following numbers.

$$-15 = -1 \cdot 15 \quad \text{and} \quad -1 + 15 = 14 \quad \text{Notice that these are two different choices.}$$

And also,

$$-15 = 1 \cdot (-15) \quad \text{and} \quad 1 + (-15) = -14 \quad \text{Notice that these are two different choices.}$$

$$\begin{array}{llll} -15 = -3 \cdot 5 & \text{and} & -3 + 5 = 2 & \leftarrow \text{ This is the correct choice.} \\ -15 = 3 \cdot (-5) & \text{and} & 3 + (-5) = -2 & \end{array}$$

The answer is $(x - 3)(x + 5)$.

We can check to see if this is correct by multiplying $(x - 3)(x + 5)$.

$$\begin{array}{r} x - 3 \\ x + 5 \\ \hline 5x - 15 \\ x^2 - 3x \\ \hline x^2 + 2x - 15 \end{array}$$

The answer checks out.

Example 8

Factor $x^2 - 10x - 24$

Solution

We are looking for an answer that is a product of two parentheses $(x \pm \underline{\quad})(x \pm \underline{\quad})$.

The number -24 can be written as the product of the following numbers.

$-24 = -1 \cdot 24$	and	$-1 + 24 = 23$	
$-24 = 1 \cdot (-24)$	and	$1 + (-24) = -23$	
$-24 = -2 \cdot 12$	and	$-2 + 12 = 10$	
$-24 = 2 \cdot (-12)$	and	$2 + (-12) = -10$	← This is the correct choice.
$-24 = -3 \cdot 8$	and	$-3 + 8 = 5$	
$-24 = 3 \cdot (-8)$	and	$3 + (-8) = -5$	
$-24 = -4 \cdot 6$	and	$-4 + 6 = 2$	
$-24 = 4 \cdot (-6)$	and	$4 + (-6) = -2$	

The answer is $(x - 12)(x + 2)$.

Example 9

Factor $x^2 + 34x - 35$

Solution

We are looking for an answer that is a product of two parentheses $(x \pm \underline{\quad})(x \pm \underline{\quad})$

The number -35 can be written as the product of the following numbers:

$-35 = -1 \cdot 35$	and	$-1 + 35 = 34$	← This is the correct choice.
$-35 = 1 \cdot (-35)$	and	$1 + (-35) = -34$	
$-35 = -5 \cdot 7$	and	$-5 + 7 = 2$	
$-35 = 5 \cdot (-7)$	and	$5 + (-7) = -2$	

The answer is $(x - 1)(x + 35)$.

Factor when a = - 1

When $a = -1$, the best strategy is to factor the common factor of -1 from all the terms in the quadratic polynomial. Then, you can apply the methods you have learned so far in this section to find the missing factors.

Example 10

Factor $x^2 + x + 6$.

Solution

First factor the common factor of -1 from each term in the trinomial. Factoring -1 changes the signs of each term in the expression.

$$-x^2 + x + 6 = -(x^2 - x - 6)$$

We are looking for an answer that is a product of two parentheses $(x \pm \underline{\quad})(x \pm \underline{\quad})$

Now our job is to factor $x^2 - x - 6$.

The number -6 can be written as the product of the following numbers.

$$\begin{array}{llll}
 -6 = -1 \cdot 6 & \text{and} & -1 + 6 = 5 & \\
 -6 = 1 \cdot (-6) & \text{and} & 1 + (-6) = -5 & \\
 -6 = -2 \cdot 3 & \text{and} & -2 + 3 = 1 & \\
 -6 = 2 \cdot (-3) & \text{and} & 2 + (-3) = -1 & \leftarrow \text{ This is the correct choice.}
 \end{array}$$

The answer is $-(x-3)(x+2)$.

To Summarize,

A quadratic of the form $x^2 + bx + c$ factors as a product of two parenthesis $(x+m)(x+n)$.

- If b and c are positive then both m and n are positive
 - Example $x^2 + 8x + 12$ factors as $(x+6)(x+2)$.
- If b is negative and c is positive then both m and n are negative.
 - Example $x^2 - 6x + 8$ factors as $(x-2)(x-4)$.
- If c is negative then either m is positive and n is negative or vice-versa
 - Example $x^2 + 2x - 15$ factors as $(x+5)(x-3)$.
 - Example $x^2 + 34x - 35$ factors as $(x+35)(x-1)$.
- If $a = -1$, factor a common factor of -1 from each term in the trinomial and then factor as usual. The answer will have the form $-(x+m)(x+n)$.
 - Example $-x^2 + x + 6$ factors as $-(x-3)(x+2)$.

Review Questions

Factor the following quadratic polynomials.

1. $x^2 + 10x + 9$
2. $x^2 + 15x + 50$
3. $x^2 + 10x + 21$
4. $x^2 + 16x + 48$
5. $x^2 - 11x + 24$
6. $x^2 - 13x + 42$
7. $x^2 - 14x + 33$
8. $x^2 - 9x + 20$
9. $x^2 + 5x - 14$
10. $x^2 + 6x - 27$
11. $x^2 + 7x - 78$
12. $x^2 + 4x - 32$
13. $x^2 - 12x - 45$
14. $x^2 - 5x - 50$
15. $x^2 - 3x - 40$
16. $x^2 - x - 56$
17. $-x^2 - 2x - 1$
18. $-x^2 - 5x + 24$
19. $-x^2 + 18x - 72$
20. $-x^2 + 25x - 150$
21. $x^2 + 21x + 108$

22. $-x^2 + 11x - 30$
23. $x^2 + 12x - 64$
24. $x^2 - 17x - 60$

Review Answers

1. $(x+1)(x+9)$
2. $(x+5)(x+10)$
3. $(x+7)(x+3)$
4. $(x+12)(x+4)$
5. $(x-3)(x-8)$
6. $(x-7)(x-6)$
7. $(x-11)(x-3)$
8. $(x-5)(x-4)$
9. $(x-2)(x+7)$
10. $(x-3)(x+9)$
11. $(x-6)(x+13)$
12. $(x-4)(x+8)$
13. $(x-15)(x+3)$
14. $(x-10)(x+5)$
15. $(x-8)(x+5)$
16. $(x-8)(x+7)$
17. $-(x+1)(x+1)$
18. $-(x-3)(x+8)$
19. $-(x-6)(x-12)$
20. $-(x-15)(x-10)$
21. $(x+9)(x+12)$
22. $-(x-5)(x-6)$
23. $(x-4)(x+16)$
24. $(x-20)(x+3)$

9.3 Factoring Special Products

Learning Objectives

- Factor the difference of two squares.
- Factor perfect square trinomials.
- Solve quadratic polynomial equation by factoring.

Introduction

When you learned how to multiply binomials we talked about two special products.

The Sum and Difference Formula	$(a + b)(a - b) = a^2 - b^2$
The Square of a Binomial Sormula	$(a + b)^2 = a^2 + 2ab + b^2$
	$(a - b)^2 = a^2 - 2ab + b^2$

In this section we will learn how to recognize and factor these special products.

Factor the Difference of Two Squares

We use the sum and difference formula to factor a difference of two squares. A difference of two squares can be a quadratic polynomial in this form.

$$a^2 - b^2$$

Both terms in the polynomial are perfect squares. In a case like this, the polynomial factors into the sum and difference of the square root of each term.

$$a^2 - b^2 = (a + b)(a - b)$$

In these problems, the key is figuring out what the a and b terms are. Lets do some examples of this type.

Example 1

Factor the difference of squares.

a) $x^2 - 9$

b) $x^2 - 100$

c) $x^2 - 1$

Solution

a) Rewrite as $x^2 - 9$ as $x^2 - 3^2$. Now it is obvious that it is a difference of squares.

The difference of squares formula is $a^2 - b^2 = (a + b)(a - b)$

Let's see how our problem matches with the formula $x^2 - 3^2 = (x + 3)(x - 3)$

The answer is $x^2 - 9 = (x + 3)(x - 3)$.

We can check to see if this is correct by multiplying $(x + 3)(x - 3)$.

$$\begin{array}{r}
 x + 3 \\
 x - 3 \\
 \hline
 -3x - 9 \\
 x^2 + 3x \\
 \hline
 x^2 + 0x - 9
 \end{array}$$

The answer checks out.

We could factor this polynomial without recognizing that it is a difference of squares. With the methods we learned in the last section we know that a quadratic polynomial factors into the product of two binomials.

$$(x \pm \underline{\quad})(x \pm \underline{\quad})$$

We need to find two numbers that multiply to -9 and add to 0 , since the middle term is missing.

We can write -9 as the following products

$$\begin{array}{llll}
 -9 = -1 \cdot 9 & \text{and} & -1 + 9 = 8 & \\
 -9 = 1 \cdot (-9) & \text{and} & 1 + (-9) = -8 & \\
 -9 = 3 \cdot (-3) & \text{and} & 3 + (-3) = 0 & \leftarrow \text{ This is the correct choice}
 \end{array}$$

We can factor $x^2 - 9$ as $(x + 3)(x - 3)$, which is the same answer as before.

You can always factor using methods for factoring trinomials, but it is faster if you can recognize special products such as the difference of squares.

b) Rewrite $x^2 - 100$ as $x^2 - 10^2$. This factors as $(x + 10)(x - 10)$.

c) Rewrite $x^2 - 1$ as $x^2 - 1^2$. This factors as $(x + 1)(x - 1)$.

Example 2

Factor the difference of squares.

a) $16x^2 - 25$

b) $4x^2 - 81$

c) $49x^2 - 64$

Solution

- a) Rewrite $16x^2 - 25$ as $(4x)^2 - 5^2$. This factors as $(4x + 5)(4x - 5)$.
 b) Rewrite $4x^2 - 81$ as $(2x)^2 - 9^2$. This factors as $(2x + 9)(2x - 9)$.
 c) Rewrite $49x^2 - 64$ as $(7x)^2 - 8^2$. This factors as $(7x + 8)(7x - 8)$.

Example 3

Factor the difference of squares:

- a) $x^2 - y^2$
 b) $9x^2 - 4y^2$
 c) $x^2y^2 - 1$

Solution

- a) $x^2 - y^2$ factors as $(x + y)(x - y)$.
 b) Rewrite $9x^2 - 4y^2$ as $(3x)^2 - (2y)^2$. This factors as $(3x + 2y)(3x - 2y)$.
 c) Rewrite as $x^2y^2 - 1$ as $(xy)^2 - 1^2$. This factors as $(xy + 1)(xy - 1)$.

Example 4

Factor the difference of squares.

- a) $x^4 - 25$
 b) $16x^4 - y^2$
 c) $x^2y^8 - 64z^2$

Solution

- a) Rewrite $x^4 - 25$ as $(x^2)^2 - 5^2$. This factors as $(x^2 + 5)(x^2 - 5)$.
 b) Rewrite $16x^4 - y^2$ as $(4x^2)^2 - y^2$. This factors as $(4x^2 + y)(4x^2 - y)$.
 c) Rewrite $x^2y^8 - 64z^2$ as $(xy^2)^2 - (8z)^2$. This factors as $(xy^2 + 8z)(xy^2 - 8z)$.

Factor Perfect Square Trinomials

We use the **Square of a Binomial Formula** to factor perfect square trinomials. A perfect square trinomial has the following form.

$$a^2 + 2ab + b^2 \qquad \text{or} \qquad a^2 - 2ab + b^2$$

In these special kinds of trinomials, the first and last terms are perfect squares and the middle term is twice the product of the square roots of the first and last terms. In a case like this, the polynomial factors into perfect squares.

$$\begin{aligned} a^2 + 2ab + b^2 &= (a + b)^2 \\ a^2 - 2ab + b^2 &= (a - b)^2 \end{aligned}$$

In these problems, the key is figuring out what the a and b terms are. Lets do some examples of this type.

Example 5

Factor the following perfect square trinomials.

- a) $x^2 + 8x + 16$

b) $x^2 - 4x + 4$

c) $x^2 + 14x + 49$

Solution

a) $x^2 + 8x + 16$

The first step is to recognize that this expression is actually perfect square trinomials.

1. Check that the first term and the last term are perfect squares. They are indeed because we can re-write:

$$x^2 + 8x + 16 \qquad \qquad \text{as} \qquad \qquad x^2 + 8x + 4^2.$$

2. Check that the middle term is twice the product of the square roots of the first and the last terms. This is true also since we can rewrite them.

$$x^2 + 8x + 16 \qquad \qquad \text{as} \qquad \qquad x^2 + 2 \cdot 4 \cdot x + 4^2$$

This means we can factor $x^2 + 8x + 16$ as $(x + 4)^2$.

We can check to see if this is correct by multiplying $(x + 4)(x + 4)$.

$$\begin{array}{r} x + 4 \\ x + 4 \\ \hline 4x + 16 \\ x^2 + 4x \\ \hline x^2 + 8x + 16 \end{array}$$

The answer checks out.

We could factor this trinomial without recognizing it as a perfect square. With the methods we learned in the last section we know that a trinomial factors as a product of the two binomials in parentheses.

$$(x \pm \underline{\quad})(x \pm \underline{\quad})$$

We need to find two numbers that multiply to 16 and add to 8. We can write 16 as the following products.

$$\begin{array}{lll} 16 = 1 \cdot 16 & \text{and} & 1 + 16 = 17 \\ 16 = 2 \cdot 8 & \text{and} & 2 + 8 = 10 \\ 16 = 4 \cdot 4 & \text{and} & 4 + 4 = 8 \end{array} \quad \leftarrow \quad \text{This is the correct choice.}$$

We can factor $x^2 + 8x + 16$ as $(x + 4)(x + 4)$ which is the same as $(x + 4)^2$.

You can always factor by the methods you have learned for factoring trinomials but it is faster if you can recognize special products.

b) Rewrite $x^2 - 4x + 4$ as $x^2 + 2 \cdot (-2) \cdot x + (-2)^2$.

We notice that this is a perfect square trinomial and we can factor it as: $(x - 2)^2$.

c) Rewrite $x^2 + 14x + 49$ as $x^2 + 2 \cdot 7 \cdot x + 7^2$.

We notice that this is a perfect square trinomial as we can factor it as: $(x + 7)^2$.

Example 6

Factor the following perfect square trinomials.

a) $4x^2 + 20x + 25$

b) $9x^2 - 24x + 16$

c) $x + 2xy + y^2$

Solution

a) Rewrite $4x^2 + 20x + 25$ as $(2x)^2 + 2 \cdot 5 \cdot (2x) + 5^2$

We notice that this is a perfect square trinomial and we can factor it as $(2x + 5)^2$.

b) Rewrite $9x^2 - 24x + 16$ as $(3x)^2 + 2 \cdot (-4) \cdot (3x) + (-4)^2$.

We notice that this is a perfect square trinomial as we can factor it as $(3x - 4)^2$.

We can check to see if this is correct by multiplying $(3x - 4)^2 = (3x - 4)(3x - 4)$.

$$\begin{array}{r}
 3x + 4 \\
 3x - 4 \\
 \hline
 -12x + 16 \\
 9x^2 - 12x \\
 \hline
 9x^2 - 24x + 16
 \end{array}$$

The answer checks out.

c) $x + 2xy + y^2$

We notice that this is a perfect square trinomial as we can factor it as $(x + y)^2$.

Solve Quadratic Polynomial Equations by Factoring

With the methods we learned in the last two sections, we can factor many kinds of quadratic polynomials. This is very helpful when we want to solve polynomial equations such as

$$ax^2 + bx + c = 0$$

Remember that to solve polynomials in expanded form we use the following steps:

Step 1

If necessary, **rewrite** the equation in standard form so that

Polynomial expression = 0.

Step 2

Factor the polynomial completely.

Step 3

Use the Zero-Product rule to **set each factor equal to zero**.

Step 4

Solve each equation from Step 3.

Step 5

Check your answers by substituting your solutions into the original equation.

We will do a few examples that show how to solve quadratic polynomials using the factoring methods we just learned.

Example 7

Solve the following polynomial equations.

a) $x^2 + 7x + 6 = 0$

b) $x^2 - 8x = -12$

c) $x^2 = 2x + 15$

Solution

a) $x^2 + 7x + 6 = 0$

Rewrite This is not necessary since the equation is in the correct form already.

Factor We can write 6 as a product of the following numbers.

$$\begin{array}{llll} 6 = 1 \cdot 6 & \text{and} & 1 + 6 = 7 & \leftarrow \text{ This is the correct choice.} \\ 6 = 2 \cdot 3 & \text{and} & 2 + 3 = 5 & \end{array}$$

$x^2 + 7x + 6 = 0$ factors as $(x + 1)(x + 6) = 0$

Set each factor equal to zero

$$x + 1 = 0 \qquad \text{or} \qquad x + 6 = 0$$

Solve

$$x = -1 \qquad \text{or} \qquad x = -6$$

Check Substitute each solution back into the original equation.

$$x = -1 \qquad (-1)^2 + 7(-1) + 6 = 1 - 7 + 6 = 0 \qquad \text{Checks out.}$$

$$x = -6 \qquad (-6)^2 + 7(-6) + 6 = 36 - 42 + 6 = 0 \qquad \text{Checks out.}$$

b) $x^2 - 8x = -12$

Rewrite $x^2 - 8x = -12$ is rewritten as $x^2 - 8x + 12 = 0$.

Factor We can write 12 as a product of the following numbers.

$12 = 1 \cdot 12$	and	$1 + 12 = 13$	
$12 = -1 \cdot (-12)$	and	$-1 + (-12) = -13$	
$12 = 2 \cdot 6$	and	$2 + 6 = 8$	
$12 = -2 \cdot (-6)$	and	$-2 + (-6) = -8$	← This is the correct choice.
$12 = 3 \cdot 4$	and	$3 + 4 = 7$	
$12 = -3 \cdot (-4)$	and	$-3 + (-4) = -7$	

$x^2 - 8x + 12 = 0$ factors as $(x - 2)(x - 6) = 0$

Set each factor equal to zero.

$$x - 2 = 0 \qquad \qquad \qquad \text{or} \qquad \qquad \qquad x - 6 = 0$$

Solve.

$$x = 2 \qquad \qquad \qquad \text{or} \qquad \qquad \qquad x = 6$$

Check Substitute each solution back into the original equation.

$x = 2$	$(2)^2 - 8(2) = 4 - 16 = -12$	Checks out.
$x = 6$	$(6)^2 - 8(6) = 36 - 48 = -12$	Checks out.

c) $x^2 = 2x + 15$

Rewrite $x^2 = 2x + 15$ is re-written as $x^2 - 2x - 15 = 0$.

Factor We can write -15 as a product of the following numbers.

$-15 = 1 \cdot (-15)$	and	$1 + (-15) = -14$	
$-15 = -1 \cdot (15)$	and	$-1 + (15) = 14$	
$-15 = -3 \cdot 5$	and	$-3 + 5 = 2$	
$-15 = 3 \cdot (-5)$	and	$3 + (-5) = -2$	← This is the correct choice.

$x^2 - 2x - 15 = 0$ factors as $(x + 3)(x - 5) = 0$.

Set each factor equal to zero

$$x + 3 = 0 \qquad \qquad \qquad \text{or} \qquad \qquad \qquad x - 5 = 0$$

Solve

$$x = -3 \qquad \qquad \qquad \text{or} \qquad \qquad \qquad x = 5$$

Check Substitute each solution back into the original equation.

$$\begin{array}{lll} x = -3 & (-3)^2 = 2(-3) + 15 \Rightarrow 9 = 0 & \text{Checks out.} \\ x = 5 & (5)^2 = 2(5) + 15 \Rightarrow 25 = 25 & \text{Checks out.} \end{array}$$

Example 8

Solve the following polynomial equations.

a) $x^2 - 12x + 36 = 0$

b) $x^2 - 81 = 0$

c) $x^2 + 20x + 100 = 0$

Solution

a) $x^2 - 12x + 36 = 0$

Rewrite This is not necessary since the equation is in the correct form already.

Factor: Re-write $x^2 - 12x + 36$ as $x^2 - 2 \cdot (-6)x + (-6)^2$.

We recognize this as a difference of squares. This factors as $(x - 6)^2 = 0$ or $(x - 6)(x - 6) = 0$.

Set each factor equal to zero

$$x - 6 = 0 \qquad \qquad \qquad \text{or} \qquad \qquad \qquad x - 6 = 0$$

Solve

$$x = 6 \qquad \qquad \qquad \text{or} \qquad \qquad \qquad x = 6$$

Notice that for a perfect square the two solutions are the same. This is called a **double root**.

Check Substitute each solution back into the original equation.

$$x = 6 \qquad \qquad 6^2 - 12(6) + 36 = 36 - 72 + 36 = 0 \qquad \qquad \text{Checks out.}$$

b) $x^2 - 81 = 0$

Rewrite This is not necessary since the equation is in the correct form already

Factor Rewrite $x^2 - 81 = 0$ as $x^2 - 9^2 = 0$.

We recognize this as a difference of squares. This factors as $(x - 9)(x + 9) = 0$.

Set each factor equal to zero.

$$x - 9 = 0 \qquad \qquad \qquad \text{or} \qquad \qquad \qquad x + 9 = 0$$

Solve:

$$x = 9 \qquad \qquad \qquad \text{or} \qquad \qquad \qquad x = -9$$

Check: Substitute each solution back into the original equation.

$x = 9$	$9^2 - 81 = 81 - 81 = 0$	Checks out.
$x = -9$	$(-9)^2 - 81 = 81 - 81 = 0$	Checks out.

c) $x^2 + 20x + 100 = 0$

Rewrite This is not necessary since the equation is in the correct form already.**Factor** Rewrite $x^2 + 20x + 100 = 0$ as $x^2 + 2 \cdot 10 \cdot x + 10^2$ We recognize this as a perfect square. This factors as: $(x + 10)^2 = 0$ or $(x + 10)(x + 10) = 0$.**Set each factor equal to zero.**

$$x + 10 = 0 \qquad \qquad \qquad \text{or} \qquad \qquad \qquad x + 10 = 0$$

Solve.

$$x = -10 \qquad \qquad \qquad \text{or} \qquad \qquad \qquad x = -10 \qquad \qquad \qquad \text{This is a double root.}$$

Check Substitute each solution back into the original equation.

$x = 10$	$(-10)^2 + 20(-10) + 100 = 100 - 200 + 100 = 0$	Checks out.
----------	---	-------------

Review Questions

Factor the following perfect square trinomials.

1. $x^2 + 8x + 16$
2. $x^2 - 18x + 81$
3. $-x^2 + 24x - 144$
4. $x^2 + 14x + 49$
5. $4x^2 - 4x + 1$
6. $25x^2 + 60x + 36$
7. $4x^2 - 12xy + 9y^2$
8. $x^4 + 22x^2 + 121$

Factor the following difference of squares.

9. $x^2 - 4$
10. $x^2 - 36$

11. $-x^2 + 100$
12. $x^2 - 400$
13. $9x^2 - 4$
14. $25x^2 - 49$
15. $-36x^2 + 25$
16. $16x^2 - 81y^2$

Solve the following quadratic equation using factoring.

17. $x^2 - 11x + 30 = 0$
18. $x^2 + 4x = 21$
19. $x^2 + 49 = 14x$
20. $x^2 - 64 = 0$
21. $x^2 - 24x + 144 = 0$
22. $4x^2 - 25 = 0$
23. $x^2 + 26x = -169$
24. $-x^2 - 16x - 60 = 0$

Review Answers

1. $(x + 4)^2$
2. $(x - 9)^2$
3. $-(x - 12)^2$
4. $(x + 7)^2$
5. $(2x - 1)^2$
6. $(5x + 6)^2$
7. $(2x - 3y)^2$
8. $(x^2 + 11)^2$
9. $(x + 2)(x - 2)$
10. $(x + 6)(x - 6)$
11. $-(x + 10)(x - 10)$
12. $(x + 20)(x - 20)$
13. $(3x + 2)(3x - 2)$
14. $(5x + 7)(5x - 7)$
15. $-(6x + 5)(6x - 5)$
16. $(4x + 9y)(4x - 9y)$
17. $x = 5, x = 6$
18. $x = -7, x = 3$
19. $x = 7$
20. $x = -8, x = 8$
21. $x = 12$
22. $x = 5/2, x = -5/2$
23. $x = -13$
24. $x = -10, x = -6$

9.4 Factoring Polynomials Completely

Learning Objectives

- Factor out a common binomial.
- Factor by grouping.
- Factor a quadratic trinomial where $a \neq 1$.
- Solve real world problems using polynomial equations.

Introduction

We say that a polynomial is **factored completely** when we factor as much as we can and we can't factor any more. Here are some suggestions that you should follow to make sure that you factor completely.

- Factor all common monomials first.
- Identify special products such as difference of squares or the square of a binomial. Factor according to their formulas.
- If there are no special products, factor using the methods we learned in the previous sections.
- Look at each factor and see if any of these can be factored further.

Here are some examples

Example 1

Factor the following polynomials completely.

a) $6x^2 - 30x + 24$

b) $2x^2 - 8$

c) $x^3 + 6x^2 + 9x$

Solution

a) $6x^2 - 30x + 24$

Factor the common monomial. In this case 6 can be factored from each term.

$$6(x^2 - 5x + 6)$$

There are no special products. We factor $x^2 - 5x + 6$ as a product of two binomials $(x \pm \underline{\quad})(x \pm \underline{\quad})$.

The two numbers that multiply to 6 and add to -5 are -2 and -3 . Let's substitute them into the two parentheses. The 6 is outside because it is factored out.

$$6(x^2 - 5x + 6) = 6(x - 2)(x - 3)$$

If we look at each factor we see that we can't factor anything else.

The answer is $6(x - 2)(x - 3)$

b) $2x^2 - 8$

Factor common monomials $2x^2 - 8 = 2(x^2 - 4)$.

We recognize $x^2 - 4$ as a difference of squares. We factor as $2(x^2 - 4) = 2(x + 2)(x - 2)$.

If we look at each factor we see that we can't factor anything else.

The answer is $2(x + 2)(x - 2)$.

c) $x^3 + 6x^2 + 9x$

Factor common monomials $x^3 + 6x^2 + 9x = x(x^2 + 6x + 9)$.

We recognize as a perfect square and factor as $x(x + 3)^2$.

If we look at each factor we see that we can't factor anything else.

The answer is $x(x + 3)^2$.

Example 2

Factor the following polynomials completely.

a) $-2x^4 + 162$

b) $x^5 - 8x^3 + 16x$

Solution

a) $-2x^4 + 162$

Factor the common monomial. In this case, factor -2 rather than 2 . It is always easier to factor the negative number so that the leading term is positive.

$$-2x^4 + 162 = -2(x^4 - 81)$$

We recognize expression in parenthesis as a difference of squares. We factor and get this result.

$$-2(x^2 - 9)(x^2 + 9)$$

If we look at each factor, we see that the first parenthesis is a difference of squares. We factor and get this answers.

$$-2(x + 3)(x - 3)(x^2 + 9)$$

If we look at each factor, we see that we can factor no more.

The answer is $-2(x + 3)(x - 3)(x^2 + 9)$

b) $x^5 - 8x^3 + 16x$

Factor out the common monomial $x^5 - 8x^3 + 16x = x(x^4 - 8x^2 + 16)$.

We recognize $x^4 - 8x^2 + 16$ as a perfect square and we factor it as $x(x^2 - 4)^2$.

We look at each term and recognize that the term in parenthesis is a difference of squares.

We factor and get: $x[(x + 2)^2(x - 2)]^2 = x(x + 2)^2(x - 2)^2$.

We use square brackets "[" and "]" in this expression because x is multiplied by the expression $(x + 2)^2(x - 2)$. When we have "nested" grouping symbols we use brackets "[" and "]" to show the levels of nesting.

If we look at each factor now we see that we can't factor anything else.

The answer is: $x(x + 2)^2(x - 2)^2$.

Factor out a Common Binomial

The first step in the factoring process is often factoring the common monomials from a polynomial. Sometimes polynomials have common terms that are binomials. For example, consider the following expression.

$$x(3x + 2) - 5(3x + 2)$$

You can see that the term $(3x + 2)$ appears in both term of the polynomial. This common term can be factored by writing it in front of a parenthesis. Inside the parenthesis, we write all the terms that are left over when we divide them by the common factor.

$$(3x + 2)(x - 5)$$

This expression is now completely factored.

Lets look at some more examples.

Example 3

Factor the common binomials.

a) $3x(x - 1) + 4(x - 1)$

b) $x(4x + 5) + (4x + 5)$

Solution

a) $3x(x - 1) + 4(x - 1)$ has a common binomial of $(x - 1)$.

When we factor the common binomial, we get $(x - 1)(3x + 4)$.

b) $x(4x + 5) + (4x + 5)$ has a common binomial of $(4x + 5)$.

When we factor the common binomial, we get $(4x + 5)(x + 1)$.

Factor by Grouping

It may be possible to factor a polynomial containing four or more terms by factoring common monomials from groups of terms. This method is called **factor by grouping**.

The next example illustrates how this process works.

Example 4

Factor $2x + 2y + ax + ay$.

Solution

There isn't a common factor for all four terms in this example. However, there is a factor of 2 that is common to the first two terms and there is a factor of a that is common to the last two terms. Factor 2 from the first two terms and factor a from the last two terms.

$$2x + 2y + ax + ay = 2(x + y) + a(x + y)$$

Now we notice that the binomial $(x + y)$ is common to both terms. We factor the common binomial and get.

$$(x + y)(2 + a)$$

Our polynomial is now factored completely.

Example 5

Factor $3x^2 + 6x + 4x + 8$.

Solution

We factor $3x$ from the first two terms and factor 4 from the last two terms.

$$3x(x + 2) + 4(x + 2)$$

Now factor $(x + 2)$ from both terms.

$$(x + 2)(3x + 4).$$

Now the polynomial is factored completely.

Factor Quadratic Trinomials Where $a \neq 1$

Factoring by grouping is a very useful method for factoring quadratic trinomials where $a \neq 1$. A quadratic polynomial such as this one.

$$ax^2 + bx + c$$

This does not factor as $(x \pm m)(x \pm n)$, so it is not as simple as looking for two numbers that multiply to give c and add to give b . In this case, we must take into account the coefficient that appears in the first term.

To factor a quadratic polynomial where $a \neq 1$, we follow the following steps.

1. We find the product ac .
2. We look for two numbers that multiply to give ac and add to give b .
3. We rewrite the middle term using the two numbers we just found.
4. We factor the expression by grouping.

Lets apply this method to the following examples.

Example 6

Factor the following quadratic trinomials by grouping.

a) $3x^2 + 8x + 4$

b) $6x^2 - 11x + 4$

c) $5x^2 - 6x + 1$

Solution

Lets follow the steps outlined above.

a) $3x^2 + 8x + 4$

Step 1 $ac = 3 \cdot 4 = 12$

Step 2 The number 12 can be written as a product of two numbers in any of these ways:

$$\begin{array}{lll} 12 = 1 \cdot 12 & \text{and} & 1 + 12 = 13 \\ 12 = 2 \cdot 6 & \text{and} & 2 + 6 = 8 \\ 12 = 3 \cdot 4 & \text{and} & 3 + 4 = 7 \end{array} \quad \text{This is the correct choice.}$$

Step 3 Re-write the middle term as: $8x = 2x + 6x$, so the problem becomes the following.

$$3x^2 + 8x + 4 = 3x^2 + 2x + 6x + 4$$

Step 4: Factor an x from the first two terms and 2 from the last two terms.

$$x(3x + 2) + 2(3x + 2)$$

Now factor the common binomial $(3x + 2)$.

$$(3x + 2)(x + 2)$$

Our answer is $(3x + 2)(x + 2)$.

To check if this is correct we multiply $(3x + 2)(x + 2)$.

$$\begin{array}{r} 3x + 2 \\ x + 2 \\ \hline 6x + 4 \\ 3x^2 + 2x \\ \hline 3x^2 + 8x + 4 \end{array}$$

The answer checks out.

b) $6x^2 - 11x + 4$

Step 1 $ac = 6 \cdot 4 = 24$

Step 2 The number 24 can be written as a product of two numbers in any of these ways.

$24 = 1 \cdot 24$	and	$1 + 24 = 25$	
$24 = -1 \cdot (-24)$	and	$-1 + (-24) = -25$	
$24 = 2 \cdot 12$	and	$2 + 12 = 14$	
$24 = -2 \cdot (-12)$	and	$-2 + (-12) = -14$	
$24 = 3 \cdot 8$	and	$3 + 8 = 11$	
$24 = -3 \cdot (-8)$	and	$-3 + (-8) = -11$	← This is the correct choice.
$24 = 4 \cdot 6$	and	$4 + 6 = 10$	
$24 = -4 \cdot (-6)$	and	$-4 + (-6) = -10$	

Step 3 Re-write the middle term as $-11x = -3x - 8x$, so the problem becomes

$$6x^2 - 11x + 4 = 6x^2 - 3x - 8x + 4$$

Step 4 Factor by grouping. Factor a $3x$ from the first two terms and factor -4 from the last two terms.

$$3x(2x - 1) - 4(2x - 1)$$

Now factor the common binomial $(2x - 1)$.

$$(2x - 1)(3x - 4)$$

Our answer is $(2x - 1)(3x - 4)$.

c) $5x^2 - 6x + 1$

Step 1 $ac = 5 \cdot 1 = 5$

Step 2 The number 5 can be written as a product of two numbers in any of these ways.

$5 = 1 \cdot 5$	and	$1 + 5 = 6$	
$5 = -1 \cdot (-5)$	and	$-1 + (-5) = -6$	← This is the correct choice

Step 3 Rewrite the middle term as $-6x = -x - 5x$. The problem becomes

$$5x^2 - 6x + 1 = 5x^2 - x - 5x + 1$$

Step 4 Factor by grouping: factor an x from the first two terms and a factor of -1 from the last two terms

$$x(5x - 1) - 1(5x - 1)$$

Now factor the common binomial $(5x - 1)$.

$$(5x - 1)(x - 1).$$

Our answer is $(5x - 1)(x - 1)$.


Solve Real-World Problems Using Polynomial Equations

Now that we know most of the factoring strategies for quadratic polynomials we can see how these methods apply to solving real world problems.

Example 7 Pythagorean Theorem

One leg of a right triangle is 3 feet longer than the other leg. The hypotenuse is 15 feet. Find the dimensions of the right triangle.

Solution



Let x = the length of one leg of the triangle, then the other leg will measure $x + 3$.

Let's draw a diagram.

Use the Pythagorean Theorem $(\text{leg}_1)^2 + (\text{leg}_2)^2 = (\text{hypotenuse})^2$ or $a^2 + b^2 = c^2$.

Here a and b are the lengths of the legs and c is the length of the hypotenuse.

Let's substitute the values from the diagram.

$$a^2 + b^2 = c^2$$

$$x^2 + (x + 3)^2 = 15^2$$

In order to solve, we need to get the polynomial in standard form. We must first distribute, collect like terms and **re-write** in the form polynomial = 0.

$$x^2 + x^2 + 6x + 9 = 225$$

$$2x^2 + 6x + 9 = 225$$

$$2x^2 + 6x - 216 = 0$$

Factor the common monomial $2(x + 3x - 108) = 0$.

To factor the trinomial inside the parenthesis we need to numbers that multiply to -108 and add to 3 . It would take a long time to go through all the options so let's try some of the bigger factors.

$$\begin{array}{ll} -108 = -12 \cdot \quad \quad \quad \text{and} \quad \quad \quad -12 + 9 = -3 \\ -108 = 12 \cdot (-9) \quad \quad \quad \text{and} \quad \quad \quad 12 + (-9) = 3 \end{array} \quad \leftarrow \text{This is the correct choice.}$$

We factor as: $2(x - 9)(x + 12) = 0$.

Set each term equal to zero and solve

$$x - 9 = 0$$

or

$$x = 9$$

$$x + 12 = 0$$

$$x = -12$$

It makes no sense to have a negative answer for the length of a side of the triangle, so the answer must be the following.

Answer $x = 9$ for one leg, and $x + 3 = 12$ for the other leg.

Check $9^2 + 12^2 = 81 + 144 = 225 = 15^2$ so the answer checks.

Example 8 Number Problems

The product of two positive numbers is 60. Find the two numbers if one of the numbers is 4 more than the other.

Solution

Let $x =$ one of the numbers and $x + 4$ equals the other number.

The product of these two numbers equals 60. We can write the equation.

$$x(x + 4) = 60$$

In order to solve we must write the polynomial in standard form. Distribute, collect like terms and re-write in the form polynomial = 0.

$$x^2 + 4x = 60$$

$$x^2 + 4x - 60 = 0$$

Factor by finding two numbers that multiply to -60 and add to 4. List some numbers that multiply to -60 :

$-60 = -4 \cdot 15$	and	$-4 + 15 = 11$	
$-60 = 4 \cdot (-15)$	and	$4 + (-15) = -11$	
$-60 = -5 \cdot 12$	and	$-5 + 12 = 7$	
$-60 = 5 \cdot (-12)$	and	$5 + (-12) = -7$	
$-60 = -6 \cdot 10$	and	$-6 + 10 = 4$	← This is the correct choice
$-60 = 6 \cdot (-10)$	and	$6 + (-10) = -4$	

The expression factors as $(x + 10)(x - 6) = 0$.

Set each term equal to zero and solve.

$$x + 10 = 0$$

or

$$x = -10$$

$$x - 6 = 0$$

$$x = 6$$


Since we are looking for positive numbers, the answer must be the following.

Answer $x = 6$ for one number, and $x + 4 = 10$ for the other number.

Check $6 \cdot 10 = 60$ so the answer checks.

Example 9 Area of a rectangle

A rectangle has sides of $x + 5$ and $x - 3$. What value of x gives an area of 48?



Solution:

Make a sketch of this situation.

Area of the rectangle = length \times width

$$(x + 5)(x - 3) = 48$$

In order to solve, we must write the polynomial in standard form. Distribute, collect like terms and **rewrite** in the form polynomial = 0.

$$x^2 + 2x - 15 = 48$$

$$x^2 + 2x - 63 = 0$$

Factor by finding two numbers that multiply to -63 and add to 2. List some numbers that multiply to -63 .

$$-63 = -7 \cdot 9$$

and

$$-7 + 9 = 2$$

← This is the correct choice

$$-63 = 7 \cdot (-9)$$

and

$$7 + (-9) = -2$$

The expression factors as $(x + 9)(x - 7) = 0$.

Set each term equal to zero and solve.

$$x + 9 = 0$$

$$x - 7 = 0$$

or

$$x = -9$$

$$x = 7$$

Since we are looking for positive numbers the answer must be $x = 7$.

Answer The width is $x - 3 = 4$ and the length is $x + 5 = 12$.

Check $4 \cdot 12 = 48$ so the answer checks out.

Review Questions

Factor completely.

- $2x^2 + 16x + 30$
- $-x^3 + 17x^2 - 70x$
- $2x^2 - 512$
- $12x^3 + 12x^2 + 3x$

Factor by grouping.

- $6x^2 - 9x + 10x - 15$
- $5x^2 - 35x + x - 7$
- $9x^2 - 9x - x + 1$
- $4x^2 + 32x - 5x - 40$

Factor the following quadratic binomials by grouping.

- $4x^2 + 25x - 21$
- $6x^2 + 7x + 1$
- $4x^2 + 8x - 5$
- $3x^2 + 16x + 21$

Solve the following application problems:

- One leg of a right triangle is 7 feet longer than the other leg. The hypotenuse is 13 feet. Find the dimensions of the right triangle.
- A rectangle has sides of $x + 2$ and $x - 1$. What value of x gives an area of 108?
- The product of two positive numbers is 120. Find the two numbers if one number is 7 more than the other.
- Framing Warehouse offers a picture framing service. The cost for framing a picture is made up of two parts. The cost of glass is \$1 per square foot. The cost of the frame is \$2 per linear foot. If the frame is a square, what size picture can you get framed for \$20?

Review Answers

- $2(x + 3)(x + 5)$
- $-x(x - 7)(x - 10)$
- $2(x - 4)(x + 4)(x^2 + 16)$
- $3x(2x + 1)^2$
- $(2x - 3)(3x + 5)$
- $(x - 7)(5x + 1)$
- $(9x - 1)(x - 1)$
- $(x + 8)(4x - 5)$
- $(4x - 3)(x + 7)$
- $(6x + 1)(x + 1)$
- $(2x - 1)(2x + 5)$
- $(x + 3)(3x + 7)$
- Leg1 = 5, Leg2 = 12
- $x = 10$
- Numbers are 8 and 15.
- You can frame a 2 foot \times 2 foot picture.